

Computing certain topological indices of thorny graphs

Seyyedeh Tahereh Jalali *

Masoud Ghods[†]

Department of Mathematics, Statistics and Computer Science

Semnan University

Semnan 35131-19111

Iran

Abstract

The topological index has transformed the chemical structure into a numeric number and they make a connection between chemistry and mathematics.

In this paper, we compute some certain topological indices like First Zagreb index, First Leap Zagreb index and F-Leap index of certain generalized graphs such as special families of thorny graphs.

Subject Classification: (2010) 05C40, 05C12, 05C07, 05C35.

Keywords: Topological index, Thorny graph.

Introduction

We recommend [5, 6, 7, 8] for general symbols and ideas in graph theory. In this paper, we have assumed the graphs that are simple, connected with no directed or weighted edges. For an instance $G = (V, E)$, we assume the number of vertices and edges are n, m and denote by $|V| = n$, $|E| = m$. The symbols $d_G(v)$ represents the degree of a vertex v which is the number of vertices are adjacent to v .

Consider two vertices u and v of G and the shortest path which connect these two vertices, the distance $d(u, v)$ denotes the number of edges in this minimum path. In graph G , for a $k \in \mathbb{Z}^+$ and a $v \in V(G)$, we define $N_k(v)$, the open k -distance neighbourhood of v , as $N_k(v) = \{u \in V(G) : d(u, v) = k\}$.

*E-mail: st_jalali@semnan.ac.ir (Corresponding Author)

[†]E-mail: mghods@semnan.ac.ir

Also, we define $d_k(v)$, the k -distance degree of a vertex v , as $d_k(v) = |N_k(v)|$. For further information, see [9].

There are a lot of topological indices that are studied.

Scientists are interested to find these topological indices to know the chemical molecular behavior, to see more details one can refer to [1, 2, 11, 13].

Among these topological indices, the most significant in which scientists were interested in studying, is Zagreb indices that correlates with molecules and their chemical properties. Gutman and Trinajstić [8] introduced the First and Second Zagreb index which were defined as

$$M_1(G) = \sum_{v \in V(G)} d_G(v)^2 = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]$$

In [10], F-index was introduced and defined as

$$F(G) = \sum_{u \in V(G)} d^3(u) = \sum_{uv \in E(G)} (d^2(u) + d^2(v)).$$

In [9], by depending on second vertex degrees, Gutman introduced the First Leap Zagreb index and defined as

$$LM_1(G) = \sum_{u \in V(G)} d_2^2(u).$$

In [12], Kulli introduced the F-Leap index of a graph as follows:

$$FL(G) = \sum_{u \in V(G)} d_2^3(u).$$

Consider a graph $G = (V, E)$, with $|V| = n, |E| = m$. In this paper we define thorn graph. The thorn graph of G , G_p with n non-negative integers like $p_1, p_2, \dots, p_n$, is achieved by connecting p_i vertices of degree one to the vertex v_i of G , for $i = 1, \dots, n$. Thorn vertices of v_i are the p_i vertices of degree one that have been joined to the vertex of v_i . Obviously, $V(G_p) = V(G) \cup V_1 \cup V_2 \cup \dots \cup V_n$. The idea of introducing the thorn graph belongs to Gutman and finally the difference of applications in chemistry was found, see [3, 4, 14].

Now we classify thorn graphs by different types of graphs like path, cycle, complete graph, complete bipartite graph, etc.

First type

Consider a path P of order greater than 2 and let the vertex v_i be the beginning vertex of such paths in graph G . The thorn path graph, G_p is achieved by joining p_i , sample of such paths at each vertex v_i of G .

Second type

Consider a cycle C of length p and let the vertex v_i be a vertex in each cycle in graph G . The thorn cycle graph, G_C is achieved by joining p_i , sample of such cycle at each vertex v_i of G .

Third type

Consider a complete graph K_p of order greater than 3 and let the vertex v_i be a vertex in K_p in graph G . The thorn complete graph, G_K is achieved by joining p_i , sample of such complete graph K_p at each vertex v_i of G

Fourth type

Consider a complete bipartite graph $K_{r,s}$ and let the vertex v_i be a vertex in a partition of $K_{r,s}$ in graph G . The thorn complete bipartite graph, G_A is achieved by joining p_i sample of such complete bipartite graph $K_{r,s}$ at each vertex v_i of G .

Fifth type

Consider a graph K_p and let the vertex v_i be a vertex in graph G . The thorn graph, G'_K is achieved by attaching p_i sample of K_p at each edge of vertex v_i of G .

Sixth type

Consider a complete bipartite graph $K_{r,s}$ and let the vertex v_i be a vertex in graph G . The thorn graph, G''_K is achieved by attaching p_i sample of $K_{r,s}$ at each edge of vertex v_i of G .

In this paper, we like to compute the First Zagreb index, First Leap Zagreb index and F-Leap index for some classifications of thorn graphs.

Results for first zagreb index

We consider the graph $G=(V,E)$. It has an order n and size m , $|V|=n$, $|E|=m$. For thorn graphs, we compute the First and Second Zagreb index which were defined as

$$M_1(G) = \sum_{v \in V(G)} d_G(v)^2 = \sum_{uv \in E(G)} [d_G(u) + d_G(v)].$$

Theorem 1 : Consider a graph G that is connected. Then we have:

$$M_1(G_p) = M_1(G) + 2 \sum_{i=1}^n d_G(v_i) p_i + \sum_{i=1}^n p_i^2 + (4p-7) \sum_{i=1}^n p_i.$$

Proof : Commonly, consider $V(G) = \{v_1, v_2, \dots, v_n\}$. Denoting the K^{th} path of order greater than 2 by $P^{(k)}$ and let the vertex v_i be the beginning vertex of a path $P^{(k)}$. Now we join $P^{(k)}$ at each vertex v_i of G . Suppose in G_p by considering $v_i = v_{i1}^{(k)}$, we have $V(P^{(k)}) = \{v_{i1}^{(k)}, v_{i2}^{(k)}, \dots, v_{ip}^{(k)}\}$, for $1 \leq i \leq n$, $1 \leq k \leq p_i$.

With a little care we notice that:

- 1) $d_{G_p}(v_i) = d_G(v_i) + p_i$,
- 2) $d_{G_p}(v_{ij}^{(k)}) = 2$ and $d_{G_p}(v_{ip}^{(k)}) = 1$; $1 \leq k \leq p_i$, $2 \leq j \leq p_i - 1$, $1 \leq i \leq n$.

So we have:

$$\begin{aligned} M_1(G_p) &= \sum_{v \in V(G_p)} d_{G_p}^2(v) \\ &= \sum_{i=1}^n [d_G(v_i) + p_i]^2 + (4(p-2)+1) \sum_{i=1}^n p_i \\ &= M_1(G) + 2 \sum_{i=1}^n d_G(v_i) p_i + \sum_{i=1}^n p_i^2 + (4p-7) \sum_{i=1}^n p_i. \end{aligned}$$

Note 1 : Let consider $p_i = b$ for all $1 \leq i \leq n$, then:

$$M_1(G_p) = M_1(G) + 4nb + nb^2 + (4p-7)nb.$$

Theorem 2 : Consider a graph G that is connected. Then we have:

$$M_1(G_C) = M_1(G) + 4 \sum_{i=1}^n d_G(v_i) p_i + 4 \sum_{i=1}^n p_i^2 + 4(p-1) \sum_{i=1}^n p_i.$$

Proof : Commonly, consider $V(G) = \{v_1, v_2, \dots, v_n\}$. Denoting the K^{th} sample of a cycle of length p by $C^{(k)}$ and let v_i is a vertex of G . Now we join $C^{(k)}$ at each vertex v_i of G . Suppose in G_C by considering $v_i = v_{i1}^{(k)}$, we have $V(C^{(k)}) = \{v_{i1}^{(k)}, v_{i2}^{(k)}, \dots, v_{ip}^{(k)}\}$, for $1 \leq i \leq n$, $1 \leq k \leq p_i$.

With a little care we notice that:

- 1) $d_{G_C}(v_i) = d_G(v_i) + 2p_i$,
- 2) $d_{G_C}(v_{ij}^{(k)}) = 2$ for all $1 \leq k \leq p_i$, $2 \leq j \leq p$, $1 \leq i \leq n$.

So we have:

$$\begin{aligned}
 M_1(G_C) &= \sum_{v \in V(G_C)} d_{G_C}^2(v) \\
 &= \sum_{i=1}^n (d_G(v_i) + 2p_i)^2 + 4(p-1) \sum_{i=1}^n p_i \\
 &= M_1(G) + 4 \sum_{i=1}^n d_G(v_i) p_i + 4 \sum_{i=1}^n p_i^2 + 4(p-1) \sum_{i=1}^n p_i.
 \end{aligned}$$

Note 2 : Let consider $p_i = b$ for all $1 \leq i \leq n$, then:

$$M_1(G_C) = M_1(G) + 8mb + 4nb^2 + 4(p-1)nb.$$

Example 1 : Consider the path P_n as graph G and let $P_i = b$ for all $1 \leq i \leq n$, then we have:

- 1) $M_1(G_p) = 4n + 4mb + nb^2 + (4p-7)nb - 6$,
- 2) $M_1(G_C) = 4n + 8mb + 4nb^2 + 4(p-1)nb - 6$.

Example 2 : Consider the cycle C_n as a graph G and let $P_i = b$ for all $1 \leq i \leq n$, then we have:

- 1) $M_1(G_p) = 4n + 4mb + nb^2 + (4p-7)nb$,
- 2) $M_1(G_C) = 4n + 8mb + 4nb^2 + 4(p-1)nb$.

Theorem 3 : Consider a graph G that is connected. Then we have:

$$M_1(G_K) = M_1(G) + 2(p-1) \sum_{i=1}^n d_G(v_i) p_i + (p-1)^2 \sum_{i=1}^n p_i^2 + (p-1)^3 \sum_{i=1}^n p_i.$$

Proof : Commonly, consider $V(G) = \{v_1, v_2, \dots, v_n\}$. Denoting the K^{th} sample of a complete graph K_p by $K_p^{(k)}$ and let v_i is a vertex of G . Then we join $K_p^{(k)}$ at each vertex v_i of G . Suppose in G_K by considering $v_i = v_{i1}^{(k)}$, we have $V(K_p^{(k)}) = \{v_{i1}^{(k)}, v_{i2}^{(k)}, \dots, v_{ip}^{(k)}\}$, for $1 \leq i \leq n$, $1 \leq k \leq p_i$.

With a little care we notice that:

- 1) $d_{G_K}(v_i) = d_G(v_i) + (p-1)p_i$,
- 2) $d_{G_K}(v_{ij}^{(k)}) = p-1$ for all $1 \leq k \leq p_i$, $2 \leq j \leq m$ and $1 \leq i \leq n$.

So we have:

$$\begin{aligned}
 M_1(G_K) &= \sum_{v \in V(G_K)} d_{G_K}^2(v) \\
 &= \sum_{i=1}^n (d_G(v_i) + (p-1)p_i)^2 + (p-1)^3 \sum_{i=1}^n p_i \\
 &= M_1(G) + 2(p-1) \sum_{i=1}^n d_G(v_i)p_i + (p-1)^2 \sum_{i=1}^n p_i^2 + (p-1)^3 \sum_{i=1}^n p_i.
 \end{aligned}$$

Note 3 : Let consider $p_i = b$ for all $1 \leq i \leq n$, then:

$$M_1(G_K) = M_1(G) + 4mb(p-1) + (p-1)^2 nb^2 + (p-1)^3 nb.$$

Theorem 4 : Consider a graph G that is connected. Then we have:

$$M_1(G_A) = M_1(G) + 2s \sum_{i=1}^n d_G(v_i)p_i + s^2 \sum_{i=1}^n p_i^2 + (r-1)s^2 \sum_{i=1}^n p_i + sr^2 \sum_{i=1}^n p_i.$$

Proof : Commonly, consider $V(G) = \{v_1, v_2, \dots, v_n\}$. Denoting the K^{th} sample of a complete bipartite graph $K_{r,s}$ by $K_{r,s}^{(k)}$ and let v_i is a vertex of G . Now we join $K_{r,s}^{(k)}$ at each vertex v_i of G . Consider the $K_{r,s}^{(k)}$ has the bipartition of $(X^{(k)}, Y^{(k)})$

Suppose in G_A by considering $v_i = v_{i1}^{(k)}$, we have $V(X^{(k)}) = \{v_{i1}^{(k)}, v_{i2}^{(k)}, \dots, v_{ir}^{(k)}\}$ and $V(Y^{(k)}) = \{u_{i1}^{(k)}, u_{i2}^{(k)}, \dots, u_{is}^{(k)}\}$ for $1 \leq i \leq n$, $1 \leq k \leq p_i$.

With a little care we notice that:

- 1) $d_{G_A}(v_i) = d_G(v_i) + sp_i$,
- 2) $d_{G_A}(v_{ij}^{(k)}) = s$ $2 \leq j \leq r$,
- 3) $d_{G_A}(u_{ij}^{(k)}) = r$ $1 \leq j \leq s$.

So we have:

$$\begin{aligned}
 M_1(G_A) &= \sum_{v \in V(G_A)} d_{G_A}^2(v) \\
 &= \sum_{i=1}^n [d_G(v_i) + sp_i]^2 + (r-1)s^2 \sum_{i=1}^n p_i + sr^2 \sum_{i=1}^n p_i
 \end{aligned}$$

$$= M_1(G) + 2s \sum_{i=1}^n d_G(v_i) p_i + s^2 \sum_{i=1}^n p_i^2 + (r-1)s^2 \sum_{i=1}^n p_i + sr^2 \sum_{i=1}^n p_i.$$

Theorem 5 : Consider a graph G that is connected. Then we have:

$$M_1(G'_K) = M_1(G) + 2 \sum_{i=1}^n d_G(v_i) p_i + \sum_{i=1}^n p_i^2 + (p-1)^3 \sum_{i=1}^n p_i + p^2 \sum_{i=1}^n p_i.$$

Proof : Commonly, consider $V(G) = \{v_1, v_2, \dots, v_n\}$. Denoting the K^{th} sample of a complete graph K_p of order p by K_p^k and let v_i be a vertex of G . Then we join $v_{i1}^{(k)}$ to the vertex v_i of G and we obtain G'_K . We have $V(K_p^{(k)}) = \{v_{i1}^{(k)}, v_{i2}^{(k)}, \dots, v_{ip}^{(k)}\}$, for $1 \leq i \leq n$, $1 \leq k \leq p$.

With a little care we notice that in G'_K :

- 1) $d_{G'_K}(v_i) = d_G(v_i) + p_i$,
- 2) $d_{G'_K}(v_{ij}^{(k)}) = p-1$ for $2 \leq j \leq p$.
- 3) $d_{G'_K}(v_{i1}^{(k)}) = p$,

So we have:

$$\begin{aligned} M_1(G'_K) &= \sum_{v \in V(G'_K)} d_{G'_K}^2(v) \\ &= \sum_{i=1}^n (d_G(v_i) + p_i)^2 + (p-1)^3 \sum_{i=1}^n p_i + p^2 \sum_{i=1}^n p_i \\ &= M_1(G) + 2 \sum_{i=1}^n d_G(v_i) p_i + \sum_{i=1}^n p_i^2 + (p-1)^3 \sum_{i=1}^n p_i + p^2 \sum_{i=1}^n p_i. \end{aligned}$$

Note 4 : Let consider $p_i = b$ for all $1 \leq i \leq n$, then:

$$M_1(G'_K) = M_1(G) + 4mb + nb^2 + (p-1)^3 nb + p^2 nb.$$

Theorem 6 : Consider a graph G that is connected. Then we have:

$$\begin{aligned} M_1(G''_K) &= M_1(G) + 2 \sum_{i=1}^n d_G(v_i) p_i + \sum_{i=1}^n p_i^2 + (s+1)^2 \sum_{i=1}^n p_i \\ &\quad + (r-1)s^2 \sum_{i=1}^n p_i + sr^2 \sum_{i=1}^n p_i. \end{aligned}$$

Proof : Commonly, consider $V(G) = \{v_1, v_2, \dots, v_n\}$. Denoting the K^{th} sample of a complete bipartite graph $K_{r,s}$ by $K_{r,s}^{(k)}$ and let v_i is a vertex of G . Now we join $v_{i1}^{(k)}$ to the vertex v_i of G in G_K'' . Consider the $K_{r,s}^{(k)}$ has the bipartition of $(X^{(k)}, Y^{(k)})$.

We have $V(X^{(k)}) = \{v_{i1}^{(k)}, v_{i2}^{(k)}, \dots, v_{ir}^{(k)}\}$ and $V(Y^{(k)}) = \{u_{i1}^{(k)}, u_{i2}^{(k)}, \dots, u_{is}^{(k)}\}$ for $1 \leq i \leq n, 1 \leq k \leq p_i$.

With a little care we notice that:

- 1) $d_{G_K''}(v_i) = d_G(v_i) + p_i \quad 1 \leq i \leq n,$
- 2) $d_{G_K''}(v_{i1}^{(k)}) = s + 1,$
- 3) $d_{G_K''}(v_{ij}^{(k)}) = s \quad 2 \leq j \leq r,$
- 4) $d_{G_K''}(u_{ij}^{(k)}) = r \quad 1 \leq j \leq s.$

So we have:

$$\begin{aligned}
 M_1(G_K'') &= \sum_{v \in V(G_K'')} d_{G_K''}^2(v) \\
 &= \sum_{i=1}^n [d_G(v_i) + p_i]^2 + (s+1)^2 \sum_{i=1}^n p_i + (r-1)s^2 \sum_{i=1}^n p_i + sr^2 \sum_{i=1}^n p_i \\
 &= M_1(G) + 2 \sum_{i=1}^n d_G(v_i) p_i + \sum_{i=1}^n p_i^2 + (s+1)^2 \sum_{i=1}^n p_i + (r-1)s^2 \sum_{i=1}^n p_i + sr^2 \sum_{i=1}^n p_i.
 \end{aligned}$$

Note 5 : Let consider $p_i = b$ for all $1 \leq i \leq n$, then:

$$M_1(G_K'') = M_1(G) + 4mb + nb^2 + (s+1)^2 nb + (r-1)s^2 nb + sr^2 nb.$$

Results for first leap zagreb index and f-leap index

We consider the First Leap Zagreb index as defined $LM_1(G) = \sum_{u \in V(G)} d_2^2(u)$ and F-Leap index as defined $FL(G) = \sum_{u \in V(G)} d_2^3(u)$. Now we compute them for some thorn graphs.

It's good to review the definition of thorn graph. Commonly, let $G = (V, E)$ be a simple connected graph in which $V = \{v_1, v_2, \dots, v_n\}$. For nonnegative integers like $p_1, p_2, \dots, p_n$, a thorn graph is a graph achieved by joining p_i vertices with degree one to each vertex v_i of G . Specifically, if $p_i = b$, then we have a b-thorny graph and denote it by G^b .

Consider the path P_n with n vertices, by joining b thorns (vertex with degree one) to each vertex of P_n , we achieved a b -thorny path graph P_n^b .

Before we state the theorems, it's better to give a vertex partition of P_n^b according to 2-degree.

Table 1
Vertex partition of P_n^b according to 2-degree

n	2-degree	Number of vertices
2	b	2b+2
3	b	2b
$n \geq 4$	b+1	b+2
	2b	1
	b	2b
	b+1	(n-2)b+2
	2b+1	2
	2b+2	n-4

Theorem 7 : Consider the b -thorny path graph P_n^b , the First Leap Zagreb index is computed as follows:

$$\begin{aligned}
 LM_1(P_2^b) &= (2b+2)b^2, \\
 LM_1(P_3^b) &= 2b^3 + (b(b+1)+2)^2 + 4b^2, \\
 LM_1(P_n^b) &= 2b^3 + [(n(b+1)-2)b(b+1)+2]^2 + 2(2b+1)^2 \\
 &\quad + (n(2b+2)-4)^2.
 \end{aligned}$$

Proof : By considering the table 1, we have:

$$\begin{aligned}
 LM_1(P_2^b) &= \sum_{v \in V(P_2^b)} d_2(v)^2 = (2b+2)b^2, \\
 LM_1(P_3^b) &= \sum_{v \in V(P_3^b)} d_2(v)^2 = 2b(b^2) + (b+2)(b+1)^2 + (2b)^2 = 2b^3 \\
 &\quad + (b(b+1)+2)^2 + 4b^2, \\
 LM_1(P_n^b) &= \sum_{v \in V(P_n^b)} d_2(v)^2 = 2b.b^2 + [(n-2)b+2](b+1)^2 + 2(2b+1)^2 \\
 &\quad + (n-4)(2b+2)^2 \\
 &= 2b^3 + [(n(b+1)-2)b(b+1)+2]^2 + 2(2b+1)^2 + (n(2b+2)-4)^2.
 \end{aligned}$$

Theorem 8 : Consider the b -thorny path graph P_n^b , the F-Leap index is computed as follows:

$$FL(P_2^b) = (2b+2)b^3,$$

$$FL(P_3^b) = 2b^4 + (b+2)(b+1)^3 + 8b^3,$$

$$FL(P_n^b) = 2b^4 + [(n-2)b+2](b+1)^3 + 2(2b+1)^3 + (n-4)(2b+2)^3.$$

Proof : By considering the definition of F-Leap index, $FL(G) = \sum_{u \in V(G)} d_2^3(u)$ and table 1 the proof is obvious.

Consider the cycle C_n with n vertices, by joining b thorns (a vertex with degree one) to each vertex of C_n , we achieved a b -thorny cycle graph C_n^b .

Before we state the theorems, it's better to give a vertex partition of C_n^b according to 2-degree.

Table 2
Vertex partition of C_n^b according to 2-degree

n	2-degree	Number of vertices
3	$b+1$	$3b$
	$2b$	3
4	$b+1$	$4b$
	$2b+1$	4
$n \geq 5$	$b+1$	nb
	$2b+2$	n

Theorem 9 : Consider the b -thorny cycle graph C_n^b , the First Leap Zagreb index is computed as follows:

$$LM_1(C_3^b) = 3b(b+1)^2 + 12b^2,$$

$$LM_1(C_4^b) = 4b(b+1)^2 + 4(2b+1)^2,$$

$$LM_1(C_n^b) = nb(b+1)^2 + n(2b+2)^2.$$

Proof : By considering the table 2, we have:

$$LM_1(C_3^b) = \sum_{v \in V(C_2^b)} d_2(v)^2 = 3b(b+1)^2 + 12b^2,$$

$$LM_1(C_4^b) = \sum_{v \in V(C_3^b)} d_2(v)^2 = 4b(b+1)^2 + 4(2b+1)^2,$$

$$LM_1(C_n^b) = \sum_{v \in V(C_n^b)} d_2(v)^2 = nb(b+1)^2 + n(2b+2)^2.$$

Theorem 10 : Consider the b -thorny cycle graph C_n^b , the F -Leap index is computed as follows:

$$FL(C_3^b) = 3b(b+1)^3 + 24b^3,$$

$$FL(C_4^b) = 4b(b+1)^3 + 4(2b+1)^3,$$

$$FL(C_n^b) = nb(b+1)^3 + n(2b+2)^3.$$

Proof : By considering the definition of F -Leap index, $FL(G) = \sum_{u \in V(G)} d_2^3(u)$ and table 2 the proof is obvious.

Consider the star S_{n+1} with $n+1$ vertices, by joining b thorns(vertex with degree one) to each vertex of S_{n+1} , we achieved a b -thorny star graph S_{n+1}^b .

Before we state the theorems, it's better to give a vertex partition of S_{n+1}^b according to 2-degree.

Table 3
Vertex partition of S_{n+1}^b according to 2-degree

2- degree	Number of vertices
b	nb
$(n-1+b)$	$n+b$
nb	1

Theorem 11 : Consider the b -thorny star graph S_{n+1}^b , the First Leap Zagreb index is computed as follows:

$$LM_1(S_{n+1}^b) = nb^3 + (n+b)(n-1+b)^2 + n^2b^2.$$

Proof : By considering the table 3 and the definition of First Leap Zagreb index, one can prove it easily.

Theorem 12 : Consider the b -thorny star graph S_{n+1}^b , the F -Leap index is computed as follows:

$$FL(S_{n+1}^b) = nb^4 + (n+b)(n-1+b)^3 + n^3b^3.$$

Proof : By considering the table 3 and the definition of F-Leap index, one can prove it easily.

Consider the complete graph K_n with n vertices, by joining b thorns (vertex with degree one) to each vertex of K_{n+1} , we achieved a b -thorny complete graph K_n^b .

Before we state the theorems, it's better to give a vertex partition of K_n^b according to 2-degree.

Table 4

Vertex partition of K_n^b according to 2-degree

2-degree	Number of vertices
$b+n-1$	nb
$(n-1)b$	n

Theorem 13 : Consider the b -thorny complete graph K_n^b , the First Leap Zagreb index is computed as follows:

$$LM_1(K_n^b) = nb(b+n-1)^2 + n((n-1)b)^2.$$

Proof : By considering the table 4 and the definition of First Leap Zagreb index, one can prove it easily.

Theorem 14 : Consider the b -thorny complete graph K_n^b , the F-Leap index is computed as follows:

$$FL(K_n^b) = nb(b+n-1)^3 + n((n-1)b)^3.$$

Proof : By considering the table 4 and the definition of F-Leap index, one can prove it easily.

Consider the complete bipartite graphs $K_{r,s}$, by joining b thorns (a vertex with degree one) to its each vertex of $K_{r,s}$, we achieved a b -thorny complete bipartite graphs $K_{r,s}^b$.

Before we state the theorems, it's better to give a vertex partition of $K_{r,s}^b$ according to 2-degree.

Table 5
Vertex partition of $K_{r,s}^b$ according to 2-degree

2-degree	Number of vertices
$b+s-1$	rb
$b+r-1$	rs
$rb+s-1$	s
$sb+r-1$	r

Theorem 15 : Consider the b -thorny complete bipartite graph $K_{r,s}^b$, the First Leap Zagreb index is computed as follows:

$$LM_1(K_{r,s}^b) = br(b+s-1)^2 + bs(b+r-1)^2 + s(br+s-1)^2 + r(bs+r-1)^2.$$

Proof : By considering the table 5 and the definition of First Leap Zagreb index, one can prove it easily.

Theorem 16 : Consider the b -thorny complete bipartite graph $K_{r,s}^b$, the F-Leap index is computed as follows:

$$FL(K_{r,s}^b) = br(b+s-1)^3 + bs(b+r-1)^3 + s(br+s-1)^3 + r(bs+r-1)^3.$$

Proof : By considering the table 5 and the definition of F-Leap index, one can prove it easily.

Conclusion

In this paper, we compute some topological indices like First Zagreb index, First Leap Zagreb index and F-Leap index and give some examples for some kind of particular structure in chemical graph recognized as thorn graph.

References

- [1] A. Faisal, S. Muhammad, A. Hafeez & Sh. Huang, "On computation of some distance-based topological indices on circulant networks-II," *Journal of Information and Optimization Sciences*, 39:4, pp. 759-782, 2018, DOI: 10.1080/02522667.2016.1223588.

- [2] A. Ahmad, K. Elahi, R. Hasni & M. F. Nadeem, "Computing the degree based topological indices of line graph of benzene ring embedded in P-type-surface in 2D network," *Journal of Information and Optimization Sciences*, 2019. 40:7, 1511-1528, DOI: 10.1080/02522667.2018.1552411
- [3] A. Heydari and I. Gutman, "On the terminal Wiener index of thorn graphs," *Kragujevac J. Sci*, no. 32, pp. 57-64, 2010.
- [4] Bytautas, L., D. Bonchev, and D. J. Klein. "On the generation of mean Wiener numbers of thorny graphs," *MATCH Commun. Math. Comput. Chem*, no. 44, pp. 31-40, 2001.
- [5] D. B. West, "Introduction to graph theory," Upper Saddle River, NJ: Prentice hall, 1996.
- [6] F. Harary, "Graph theory," Narosa Publication, New Delhi, 2001.
- [7] J. A. Bondy and R. M. Uppaluri Siva, "Graph theory with applications," London: Macmillan, Vol. 290. 1976.
- [8] I. Gutman and T. Nenad, "Graph theory and molecular orbitals. Total ϕ -electron energy of alternant hydrocarbons," *Chemical Physics Letters* 17. no. 4, pp. 535-538, 1972.
- [9] I. Gutman., A. M. Naji, and N. D. Soner. "On Leap Zagreb indices of graphs," *Communication in Combinatorics and Optimization* 2, no. 2, pp. 99-117, 2017.
- [10] I. Z. Milovanović, M. M. Matejić and R.I. Milovanović, "Remark on Forgotten Topological Index of a Line Graph," *Bulletin of international mathematical virtual* 7, no. 3, pp. 473-478, 2017.
- [11] M. N. Jahangeer Baig, Ch. Yong Jung, N. Ahmad & Sh. Min Kang, "On the M-polynomials and degree-based topological indices of an important class of graphs," *Journal of Discrete Mathematical Sciences and Cryptography*, 2019. 22:7, 1281-1288, DOI: 10.1080/09720529.2019.1691327
- [12] V. R. Kulli, "On F-Leap indices and F-Leap polynomials of some graphs," *International Journal of Mathematical Archive* 9. no. 12, pp. 41-49, 2018.

- [13] W. Gao., and M. R. Farahani, “ The Zagreb topological indices for a type of Benzenoid systems jagged-rectangle,” *Journal of Interdisciplinary Mathematics*, 20:5, pp. 1341-1348, 2017.DOI:0.1080/09720502.2016.1232037
- [14] Y. Alizadeh, A. Iranmanesh, D.Tomislav et al., “The edge Wiener index of suspensions, bottlenecks, and thorny graphs,” *Glasnik matematički*, no. 49, pp.1-12, 2014.

Received September, 2020

Revised January, 2021