

Norms over bifuzzy submodules

Rasul Rasuli

Department of Mathematics

Payame Noor University (PNU)

Tehran

Iran

Abstract

In this work, we introduce the concepts of bifuzzy submodules under norms and consider some of their properties and structured characteristics. Also, we investigate the intersection and direct product of them and obtain some new results. Finally, we investigate their properties under homomorphisms of modules.

Subject Classification: 03E72, 47A30, 06F25, 55N45, 13B10.

Keywords: Fuzzy set theory, Bifuzzy set, Norms, Modules, Products and intersections, Morphisms.

1. Introduction

Fuzzy sets were introduced by Zadeh[48] and Dieter Klaua [9] in 1965 as an extension of the classical notion of set. As a generalization of a fuzzy set, the concept of a bifuzzy set was introduced by Atanassov [3, 4]. Many researches introduced and presented various types of norms (T and C) for particular purposes [2, 7, 8, 33]. Malakooti Rad and Mahdavi [11] investigated a note on the intersection graph of submodules of a module. The author by using norms, investigated some properties of fuzzy algebraic structures [12-46]. In this work as Section 2, we define bifuzzy submodules under norms and obtain some conditions that bifuzzy submodules will be submodules. In Section 3, we introduce the concepts of intersection and direct product of bifuzzy submodules under norms and investigate some of their properties. Next, we consider some of their properties and

E-mail: rasulirasul@yahoo.com

structures of bifuzzy submodules under norms by homomorphisms of R -submodules.

2. Bifuzzy submodules under norms

We have following definition from [5].

Definition 2.1 : Let X a non-empty sets. A fuzzy subset μ of X is a function $\mu : X \rightarrow [0, 1]$. Denote by $[0, 1]^X$, the set of all fuzzy subset of X .

We give following definition from [3, 6].

Definition 2.2 : Let X be a nonempty set. An intuitionistic fuzzy set is an object of the form: $A = \{(\mu_A(x), \nu_A(x)) : x \in X\}$ such that $\mu_A, \nu_A \in [0, 1]^X$ and for all $x \in X$ we have $\mu_A(x) + \nu_A(x) \in [0, 1]$. This object is also called a bifuzzy set. For the sake of simplicity, we shall use the symbol $A = (\mu_A, \nu_A)$ for the $A = \{(\mu_A(x), \nu_A(x)) : x \in X\}$.

Let X be a nonempty set and let $A = (\mu_A, \nu_A)$ and $B = (\mu_B, \nu_B)$ be IFSs in X . Then

- (1) $A \subseteq B$ iff $\mu_A \leq \mu_B$ and $\nu_A \geq \nu_B$.
- (2) $A = B$ iff $A \subseteq B$ and $B \subseteq A$.

We have following definition from [5].

Definition 2.3 : A t -norm T is a function $T : [0, 1] \times [0, 1] \rightarrow [0, 1]$ having the following four properties:

- (T1) $T(x, 1) = x$ (neutral element),
 - (T2) $T(x, y) \leq T(x, z)$ if $y \leq z$ (monotonicity),
 - (T3) $T(x, y) = T(y, x)$ (commutativity),
 - (T4) $T(x, T(y, z)) = T(T(x, y), z)$ (associativity),
- for all $x, y, z \in [0, 1]$.

It is clear that if $x_1 \geq x_2$ and $y_1 \geq y_2$, then $T(x_1, y_1) \geq T(x_2, y_2)$.

Example 2.4 :

- (1) Standard intersection t -norm $T_m(x, y) = \min\{x, y\}$.
- (2) Bounded sum t -norm $T_b(x, y) = \max\{0, x + y - 1\}$.
- (3) algebraic product t -norm $T_p(x, y) = xy$.
- (4) Drastic t -norm

$$T_D(x, y) = \begin{cases} y & \text{if } x = 1 \\ x & \text{if } y = 1 \\ 0 & \text{otherwise.} \end{cases}$$

(5) Nilpotent minimum t -norm

$$T_{nM}(x, y) = \begin{cases} \min\{x, y\} & \text{if } x + y > 1 \\ 0 & \text{otherwise.} \end{cases}$$

(6) Hamacher product t -norm

$$T_{H_0}(x, y) = \begin{cases} 0 & \text{if } x = y = 0 \\ \frac{xy}{x + y - xy} & \text{otherwise.} \end{cases}$$

The drastic t -norm is the pointwise smallest t -norm and the minimum is the pointwise largest t -norm: $T_D(x, y) \leq T(x, y) \leq T_{\min}(x, y)$ for all $x, y \in [0, 1]$.

We have following definition from [5].

Definition 2.5 : A t -conorm C is a function $C : [0, 1] \times [0, 1] \rightarrow [0, 1]$ having the following four properties:

- (C1) $C(x, 0) = x$,
- (C2) $C(x, y) \leq C(x, z)$ if $y \leq z$,
- (C3) $C(x, y) = C(y, x)$,
- (C4) $C(x, C(y, z)) = C(C(x, y), z)$,

for all $x, y, z \in [0, 1]$.

- (1) Standard union t -conorm $C_m(x, y) = \max\{x, y\}$.
- (2) Bounded sum t -conorm $C_b(x, y) = \min\{1, x + y\}$.
- (3) Algebraic sum t -conorm $C_p(x, y) = x + y - xy$.
- (4) Drastic T -conorm

$$C_D(x, y) = \begin{cases} y & \text{if } x = 0 \\ x & \text{if } y = 0 \\ 1 & \text{otherwise,} \end{cases}$$

dual to the drastic t -norm.

- (5) Nilpotent maximum t -conorm, dual to the nilpotent minimum t -norm:

$$C_{nM}(x, y) = \begin{cases} \max\{x, y\} & \text{if } x + y < 1 \\ 1 & \text{otherwise.} \end{cases}$$

- (6) Einstein sum (compare the velocity-addition formula under special relativity) $C_{H_2}(x, y) = \frac{x+y}{1+xy}$ is dual to one of the Hamacher t -norms. Note that all t -conorms are bounded by the maximum and the drastic t -conorm: $C_{\max}(x, y) \leq C(x, y) \leq C_D(x, y)$ for any t -conorm C and all $x, y \in [0, 1]$.

Recall that t -norm T (t -conorm C) is idempotent if for all $x \in [0, 1]$, $T(x, x) = x$ ($C(x, x) = x$).

Following lemma is from [1].

Lemma 2.7 : Let T be a t -norm and C be a t -conorm. Then

$$T(T(x, y), T(w, z)) = T(T(x, w), T(y, z))$$

and

$$C(C(x, y), C(w, z)) = C(C(x, w), C(y, z)),$$

for all $x, y, w, z \in [0, 1]$.

we have the following definition in [10].

Definition 2.8 : Let R be a ring. A commutative group $(M, +)$ is called a left R -module or a left module over R with respect to a mapping

$$\therefore R \times M \rightarrow M$$

if for all $r, s \in R$ and $m, n \in M$,

$$(1) \quad r.(m + n) = r.m + r.n,$$

$$(2) \quad r.(s.m) = (rs).m,$$

$$(3) \quad (r + s).m = r.m + s.m.$$

If R has an identity 1 and if $1.m = m$ for all $m \in M$, then M is called a unitary or unital left R -module. A right R -module can be defined in a similar fashion.

Let M be an R -module and S be a nonempty subset of M . Then S is called a submodule of M if S is a subgroup of M and for all $r \in R, a \in S$, we have $ra \in S$.

Now we define bifuzzy submodule of R -module M under norms.

Definition 2.9 : A bifuzzy set $M^* = (\mu_M, \nu_M)$ in R -module M is called bifuzzy submodule of M with respect to a t -norm T and a t -conorm C , if it satisfies:

- (1) $M^*(x+y) \supseteq (T(\mu_M(x), \mu_M(y)), C(\nu_M(x), \nu_M(y)))$,
- (2) $M^*(rx) \supseteq M^*(x)$,
- (3) $M^*(0) = (1, 0)$,

for all $x, y \in M$ and $r \in R$.

Denote by $BFN(M)$, the set of all bifuzzy submodules of M with respect to norms $(T$ and $C)$.

Example 2.10 : Let $R = (\mathbb{Z}, +, -)$ be a ring of integers and $M = \mathbb{Z}$ be an R -module. Define $\mu_{\mathbb{Z}} : \mathbb{Z} \rightarrow [0, 1]$ as

$$\mu_{\mathbb{Z}}(z) = \begin{cases} 0.45 & \text{if } z \in \{\pm 1, \pm 3, \dots\} \\ 0 & \text{if } z = 0 \\ 0.35 & \text{if } z \in \{\pm 2, \pm 4, \dots\} \end{cases}$$

and

$\nu_{\mathbb{Z}} : \mathbb{Z} \rightarrow [0, 1]$ as

$$\nu_{\mathbb{Z}}(z) = \begin{cases} 0.55 & \text{if } z \in \{\pm 1, \pm 3, \dots\} \\ 0 & \text{if } z = 0 \\ 0.65 & \text{if } z \in \{\pm 2, \pm 4, \dots\}. \end{cases}$$

For all $x, y \in [0, 1]$ if $C(x, y) = C_p(x, y) = x + y - xy$ and $T(x, y) = T_p(x, y) = xy$, then $\mathbb{Z}^* = (\mu_{\mathbb{Z}}, \nu_{\mathbb{Z}}) \in BFN(\mathbb{Z})$.

Proposition 2.11 : Let R a ring with identity 1, and T and C be idempotent. Then for any $r \in R$ and $x \in M$ the condition (2) in Definition 2.9 is equivalent to the condition $M^*(rx) = M^*(x)$.

Proof : As $M^*(rx) = (\mu_M(rx), \nu_M(rx))$ so from condition (2) we get that

- (a) $\mu_M(rx) \geq \mu_M(x)$ and
- (b) $\nu_M(rx) \leq \nu_M(x)$. Now

$$\begin{aligned}
\mu_M(x) &= \mu_M(rx + (1-r)x) \geq T(\mu_M(rx), \mu_M((1-r)x)) = T(\mu_M(rx), \mu_M(x-rx)) \\
&\geq T(\mu_M(rx), T(\mu_M(x), \mu_M(-rx))) \geq T(\mu_M(x), T(\mu_M(x), \mu_M(-rx))) \\
&= T(T(\mu_M(x), \mu_M(x)), \mu_M(-rx)) = T(\mu_M(x), \mu_M(-rx)) \\
&= T(\mu_M(x), \mu_M(-1rx)) \geq T(\mu_M(x), \mu_M(rx))
\end{aligned}$$

then we obtain $\mu_M(x) = T(\mu_M(x), \mu_M(x)) \geq T(\mu_M(x), \mu_M(rx))$ and so

$$(c) \quad \mu_M(x) \geq \mu_M(rx).$$

Also

$$\begin{aligned}
\nu_M(x) &= \nu_M(rx + (1-r)x) \leq C(\nu_M(rx), \nu_M((1-r)x)) = C(\nu_M(rx), \nu_M(x-rx)) \\
&\leq C(\nu_M(rx), C(\nu_M(x), \nu_M(-rx))) \leq C(\nu_M(x), C(\nu_M(x), \nu_M(-rx))) \\
&= C(C(\nu_M(x), \nu_M(x)), \nu_M(-rx)) = C(\nu_M(x), \nu_M(-rx)) \\
&= C(\nu_M(x), \nu_M(-1rx)) \leq C(\nu_M(x), \nu_M(rx))
\end{aligned}$$

then we get $\nu_M(x) = C(\nu_M(x), \nu_M(x)) \leq C(\nu_M(x), \nu_M(rx))$ and then

$$(d) \quad \nu_M(x) \leq \nu_M(rx). \text{ Then from (a) and (c) we get that } \mu_M(rx) = \mu_M(x) \text{ and from (b) and (d) we have } \nu_M(rx) = \nu_M(x). \text{ Thus } M^*(rx) = M^*(x).$$

□

Proposition 2.12 : Let R be a field. Then the condition (2) in Definition 2.9 for all $r \in R, r \neq 0$, will be equivalent to condition $M^*(rx) = M^*(x)$.

Proof: If the the condition (2) be satisfied, then $\mu_M(rx) \geq \mu_M(x) = \mu_M(\frac{1}{r}(rx)) \geq \mu_M(rx)$ and so $\mu_M(rx) = \mu_M(x)$. Also $\nu_M(rx) \leq \nu_M(x) = \nu_M(\frac{1}{r}(rx)) \leq \nu_M(rx)$ and so $\nu_M(rx) = \nu_M(x)$. Therefore $M^*(rx) = M^*(x)$. □

Proposition 2.13 : Let there exists $x \in M$ such that $M^*(x) = (1, 0)$. If R a ring with identity 1, then the condition (3) in Definition 2.9 follows from conditions (1) and (2).

Proof: Let conditions (1) and (2) in Definition 2.9 are hold and there exists $x \in M$ such that $M^*(x) = (\mu_M(x), \nu_M(x)) = (1, 0)$. Then for such x , we have

$$\mu_M(0) = \mu_M(x-x) \geq T(\mu_M(x), \mu_M(-x)) = T(1, \mu_M(-x)) = \mu_M(-x) = \mu_M(-1x) \geq \mu_M(x) = 1$$

and so $\mu_M(0) = 1$. Also

$$\nu_M(0) = \nu_M(x-x) \leq C(\nu_M(x), \nu_M(-x)) = C(0, \nu_M(-x)) = \nu_M(-x) = \nu_M(-1x) \leq \nu_M(x) = 0$$

and then $\nu_M(0) = 0$.

Thus $M^*(x) = (1, 0)$ and thus condition (3) will be hold. $\square$

The following propositions investigate the relationships between bifuzzy submodules of R -module M under norms and submodules of R -modules M .

Proposition 2.14 : Let $M^* = (\mu_M, \nu_M) \in BFN(M)$. Then $S = \{x \mid x \in M, M^*(x) = (1, 0)\}$ is a submodule of the module M and $M^* \in BFN(S)$.

Proof : Let $x, y \in S$ and $r \in R$. Then $\mu_M(x) = 1 = \mu_M(y)$ and $\nu_M(x) = 0 = \nu_M(y)$. Let $M^* = (\mu_M, \nu_M) \in BFN(M)$.

(1) As $\mu_M(x+y) \geq T(\mu_M(x), \mu_M(y)) = T(1, 1) = 1$ so $\mu_M(x+y) = 1$.

(2) As $\nu_M(x+y) \leq C(\nu_M(x), \nu_M(y)) = C(0, 0) = 0$ and $\nu_M(x+y) = 0$.

Now from (1) and (2) we get that $x+y \in S$.

(3) As $\mu_M(rx) \geq \mu_M(x) = 1$ so $\mu_M(rx) = 1$.

(4) As $\nu_M(rx) \leq \nu_M(x) = 0$ so $\nu_M(rx) = 0$.

Then from (3) and (4) we obtain that $rx \in S$.

(5) Also since $M^*(0) = (1, 0)$ so $0 \in S$.

Now from (1)-(5) we get that S will be a submodule of the module M .

The second part of the statement is obvious. $\square$

Proposition 2.15 : Let $M^* = (\mu_M, \nu_M) \in BFN(M)$. If T and C be idempotent, then

$$S_t = \{x \mid x \in M, M^*(x) \supseteq M^*(t)\}$$

will be a submodule of the module M for any $t \in [0, 1]$.

Proof : Let $x, y \in S_t$ and $r \in R$. Then $\mu_M(x) \geq t$ and $\mu_M(y) \geq t$. Also $\nu_M(x) \leq t$ and $\nu_M(y) \leq t$. Let $A = (\mu_M, \nu_M) \in BFN(M)$.

(1) Since

$$\mu_M(x+y) \geq T(\mu_M(x), \mu_M(y)) \geq T(t, t) = t$$

and

$$\nu_M(x+y) \leq C(\nu_M(x), \nu_M(y)) \leq C(t, t) = t$$

so $\mu_M(x+y) \geq t$ and $\nu_M(x+y) \leq t$. Then $x+y \in S_t$.

- (2) As $\mu_M(rx) \geq \mu_M(x) = t$ and $\nu_M(rx) \leq \nu_M(x) = t$ so $\mu_M(rx) \geq t$ and $\nu_M(rx) \leq t$.

Thus $rx \in S_t$.

- (3) Also since $M^*(0) = (1, 0) \geq (\mu_M(t), \nu_M(t)) = M^*(t)$ for any $t \in [0, 1]$ so $0 \in S_t$.

Now from (1)-(3) we get that S_t will be a submodule of the module M . $\square$

Proposition 2.16 : Let M be an R -module and S be a subset of M . Let

$\mu_M : S \rightarrow \{0, 1\}$ be the characteristic function of S as

$$\mu_M(x) = \begin{cases} 1 & \text{if } x \in S \\ 0 & \text{if } x \notin S \end{cases}$$

and $\nu_M : S \rightarrow \{0, 1\}$ be the anti characteristic function of S as

$$\nu_M(x) = \begin{cases} 0 & \text{if } x \in S \\ 1 & \text{if } x \notin S. \end{cases}$$

Then $M^* = (\mu_M, \nu_M) \in BFN(M)$ if and only if S is a submodule of M .

Proof : Let $M^* = (\mu_M, \nu_M) \in BFN(M)$ and we prove that S is a submodule of M . Let $x, y \in S \subseteq M$ and $r \in R$. Since $\mu_M(x+y) \geq T(\mu_M(x), \mu_M(y)) = T(1, 1) = 1$ and $\nu_M(x+y) \leq C(\nu_M(x), \nu_M(y)) = C(0, 0) = 0$ so $\mu_M(x+y) = 1$ and $\nu_M(x+y) = 0$. Thus $x+y \in S$. Also $\mu_M(rx) \geq \mu_M(x) = 1$ and $\nu_M(rx) \leq \nu_M(x) = 0$ and then $rx \in S$. Finally $M^*(0) = (1, 0)$ means that $0 \in S$. Thus S is a submodule of M .

Conversely, let S is a submodule of M and we prove that $M^* = (\mu_M, \nu_M) \in BFN(M)$. Let $x, y \in M$.

- (1) If $x, y \in S$, then $x+y \in S$. Now

$$\mu_M(x+y) = 1 \geq 1 = T(1, 1) = T(\mu_M(x), \mu_M(y))$$

and

$$\nu_M(x+y) = 0 \leq 0 = C(0,0) = C(\nu_M(x), \nu_M(y)).$$

(2) If $x \in S$ and $y \notin S$, then $x+y \notin S$. Then

$$\mu_M(x+y) = 0 \geq 0 = T(1,0) = T(\mu_M(x), \mu_M(y))$$

and

$$\nu_M(x+y) = 1 \leq 1 = C(0,1) = C(\nu_M(x), \nu_M(y)).$$

(3) If $x, y \notin S$, then we have the following conditions.

(a) If $x+y \notin S$, then

$$\mu_M(x+y) = 0 \geq 0 = T(0,0) = T(\mu_M(x), \mu_M(y))$$

and

$$\nu_M(x+y) = 1 \leq 1 = C(1,1) = C(\nu_M(x), \nu_M(y)).$$

(b) If $x+y \in S$, then

$$\mu_M(x+y) = 1 \geq 0 = T(0,0) = T(\mu_M(x), \mu_M(y))$$

and

$$\nu_M(x+y) = 0 \leq 1 = C(1,1) = C(\nu_M(x), \nu_M(y)).$$

Therefore from (1)-(3) we obtain that

$$M^*(x+y) \supseteq (T(\mu_M(x), \mu_M(y)), C(\nu_M(x), \nu_M(y))).$$

Now let $x \in M$ and $r \in R$.

(1) if $x \in S$, then $rx \in S$ and so $\mu_M(rx) = 1 \geq 1 = \mu_M(x)$ and $\nu_M(rx) = 0 \leq 0 = \nu_M(x)$.

(2) if $x \notin S$, then $rx \notin S$ and so $\mu_M(rx) = 0 \geq 0 = \mu_M(x)$ and $\nu_M(rx) = 1 \leq 1 = \nu_M(x)$.

Thus (1) and (2) give us that $M^*(rx) \supseteq M^*(x)$.

Finally, as $0 \in S$ so $\mu_M(0) = 1$ and $\nu_M(0) = 0$ and then $M^*(0) = (\mu_M(0), \nu_M(0)) = (1, 0)$.

Hence $M^* = (\mu_M, \nu_M) \in \text{BFN}(M)$. □

3. Intersections, direct products and homomorphisms of bifuzzy submodules under norms

Definition 3.1 : (See [14]) Let $A = (\mu_A, \nu_A)$ and $B = (\mu_B, \nu_B)$ be two intuitionistic fuzzy sets of X . Define intersection of A and B as $A \cap B = (\mu_{A \cap B}, \nu_{A \cap B})$ by

$$\mu_{A \cap B}(x) = T(\mu_A(x), \mu_B(x))$$

and

$$\nu_{A \cap B}(x) = C(\nu_A(x), \nu_B(x))$$

for all $x \in X$.

Proposition 3.2 : Let $M_1^* = (\mu_{M_1^*}, \nu_{M_1^*})$ and $M_2^* = (\mu_{M_2^*}, \nu_{M_2^*})$ be two intuitionistic fuzzy sets in R -module M . If $M_1^*, M_2^* \in BFN(M)$, then

$$M_1^* \cap M_2^* = (\mu_{M_1^* \cap M_2^*}, \nu_{M_1^* \cap M_2^*}) \in BFN(M).$$

Proof : Let $x, y \in M$. Since

$$\begin{aligned} \mu_{M_1^* \cap M_2^*}(x+y) &= T(\mu_{M_1^*}(x+y), \mu_{M_2^*}(x+y)) \geq T(T(\mu_{M_1^*}(x), \mu_{M_1^*}(y)), T(\mu_{M_2^*}(x), \mu_{M_2^*}(y))) \\ &= T(T(\mu_{M_1^*}(x), \mu_{M_2^*}(x)), T(\mu_{M_1^*}(y), \mu_{M_2^*}(y))) = T(\mu_{M_1^* \cap M_2^*}(x), \mu_{M_1^* \cap M_2^*}(y)) \end{aligned}$$

so

$$\mu_{M_1^* \cap M_2^*}(x+y) \geq T(\mu_{M_1^* \cap M_2^*}(x), \mu_{M_1^* \cap M_2^*}(y)).$$

Also

$$\begin{aligned} \nu_{M_1^* \cap M_2^*}(x+y) &= C(\nu_{M_1^*}(x+y), \nu_{M_2^*}(x+y)) \leq C(C(\nu_{M_1^*}(x), \nu_{M_1^*}(y)), C(\nu_{M_2^*}(x), \nu_{M_2^*}(y))) \\ &= C(C(\nu_{M_1^*}(x), \nu_{M_2^*}(x)), C(\nu_{M_1^*}(y), \nu_{M_2^*}(y))) = C(\nu_{M_1^* \cap M_2^*}(x), \nu_{M_1^* \cap M_2^*}(y)) \end{aligned}$$

and then

$$\nu_{M_1^* \cap M_2^*}(x+y) \leq C(\nu_{M_1^* \cap M_2^*}(x), \nu_{M_1^* \cap M_2^*}(y)).$$

Thus

$$(M_1^* \cap M_2^*)(x+y) \supseteq (T(\mu_{M_1^* \cap M_2^*}(x), \mu_{M_1^* \cap M_2^*}(y)), C(\nu_{M_1^* \cap M_2^*}(x), \nu_{M_1^* \cap M_2^*}(y))).$$

Let $x \in M$ and $r \in R$. As

$$\mu_{M_1^* \cap M_2^*}(rx) = T(\mu_{M_1^*}(rx), \mu_{M_2^*}(rx)) \geq T(\mu_{M_1^*}(x), \mu_{M_2^*}(x)) = \mu_{M_1^* \cap M_2^*}(x)$$

and

$$\nu_{M_1^* \cap M_2^*}(rx) = C(\nu_{M_1^*}(rx), \nu_{M_2^*}(rx)) \leq C(\nu_{M_1^*}(x), \nu_{M_2^*}(x)) = \nu_{M_1^* \cap M_2^*}(x)$$

so $(M_1^* \cap M_2^*)(rx) \supseteq (M_1^* \cap M_2^*)(x)$.

Finally, from

$$\mu_{M_1^* \cap M_2^*}(0) = T(\mu_{M_1^*}(0), \mu_{M_2^*}(0)) = T(1, 1) = 1$$

and

$$\nu_{M_1^* \cap M_2^*}(0) = C(\nu_{M_1^*}(0), \nu_{M_2^*}(0)) = C(0, 0) = 0$$

we obtain that

$$(M_1^* \cap M_2^*)(0) = (\mu_{M_1^* \cap M_2^*}(0), \nu_{M_1^* \cap M_2^*}(0)) = (1, 0).$$

Hence

$$M_1^* \cap M_2^* = (\mu_{M_1^* \cap M_2^*}, \nu_{M_1^* \cap M_2^*}) \in BFN(M).$$

□

Corollary 3.3 : Let $\{M_i = (\mu_{M_i}, \nu_{M_i}) \mid i = 1, 2, 3, \dots, n\} \subseteq BFN(M)$. Then $\cap_{M_i} \in BFN(M)$.

Definition 3.4 : (See [14]) Let $A = (\mu_A, \nu_A)$ and $B = (\mu_B, \nu_B)$ be two intuitionistic fuzzy sets in X and Y , respectively. The direct product of A and B , denoted by $A \times B = (\mu_{A \times B}, \nu_{A \times B})$, is an intuitionistic fuzzy set in $X \times Y$ such that for all x in X and y in Y , define

$$\mu_{A \times B}(x, y) = T(\mu_A(x), \mu_B(y))$$

and

$$\nu_{A \times B}(x, y) = C(\nu_A(x), \nu_B(y)).$$

Proposition 3.5 : Let M_i be R -modules and $M_i^* = (\mu_{M_i^*}, \nu_{M_i^*}) \in BFN(M)$ for $i = 1, 2$. Then

$$M_1^* \times M_2^* = (\mu_{M_1^* \times M_2^*}, \nu_{M_1^* \times M_2^*}) \in BFN(M_1 \times M_2).$$

Proof:

(1) Let $(x_1, y_1), (x_2, y_2) \in M_1 \times M_2$. Then

$$\begin{aligned} \mu_{M_1^* \times M_2^*}((x_1, y_1) + (x_2, y_2)) &= \mu_{M_1^* \times M_2^*}(x_1 + x_2, y_1 + y_2) \\ &= T(\mu_{M_1^*}(x_1 + x_2), \mu_{M_2^*}(y_1 + y_2)) \geq T(T(\mu_{M_1^*}(x_1), \mu_{M_1^*}(x_2)), T(\mu_{M_2^*}(y_1), \mu_{M_2^*}(y_2))) \\ &= T(T(\mu_{M_1^*}(x_1), \mu_{M_2^*}(y_1)), T(\mu_{M_1^*}(x_2), \mu_{M_2^*}(y_2))) = T(\mu_{M_1^* \times M_2^*}(x_1, y_1), \mu_{M_1^* \times M_2^*}(x_2, y_2)). \end{aligned}$$

Also

$$\begin{aligned} \nu_{M_1^* \times M_2^*}((x_1, y_1) + (x_2, y_2)) &= \nu_{M_1^* \times M_2^*}(x_1 + x_2, y_1 + y_2) \\ &= C(\nu_{M_1^*}(x_1 + x_2), \nu_{M_2^*}(y_1 + y_2)) \leq C(C(\nu_{M_1^*}(x_1), \nu_{M_1^*}(x_2)), C(\nu_{M_2^*}(y_1), \nu_{M_2^*}(y_2))) \\ &= C(C(\nu_{M_1^*}(x_1), \nu_{M_2^*}(y_1)), C(\nu_{M_1^*}(x_2), \nu_{M_2^*}(y_2))) = C(\nu_{M_1^* \times M_2^*}(x_1, y_1), \nu_{M_1^* \times M_2^*}(x_2, y_2)). \end{aligned}$$

Thus

$$\begin{aligned} &(M_1^* \times M_2^*)((x_1, y_1) + (x_2, y_2)) \\ &\supseteq (T(\mu_{M_1^* \times M_2^*}(x_1, y_1), \mu_{M_1^* \times M_2^*}(x_2, y_2)), C(\nu_{M_1^* \times M_2^*}(x_1, y_1), \nu_{M_1^* \times M_2^*}(x_2, y_2))). \end{aligned}$$

(2) Let $(x, y) \in M_1 \times M_2$ and $r \in R$. Now

$$\begin{aligned} \mu_{M_1^* \times M_2^*}(r(x, y)) &= \mu_{M_1^* \times M_2^*}(rx, ry) = T(\mu_{M_1^*}(rx), \mu_{M_2^*}(ry)) \\ &\geq T(\mu_{M_1^*}(x), \mu_{M_2^*}(y)) = \mu_{M_1^* \times M_2^*}(x, y). \end{aligned}$$

Also

$$\begin{aligned} \nu_{M_1^* \times M_2^*}(r(x, y)) &= \nu_{M_1^* \times M_2^*}(rx, ry) = C(\nu_{M_1^*}(rx), \nu_{M_2^*}(ry)) \\ &\leq C(\nu_{M_1^*}(x), \nu_{M_2^*}(y)) = \nu_{M_1^* \times M_2^*}(x, y). \end{aligned}$$

Therefore $(M_1^* \times M_2^*)(r(x, y)) \supseteq (M_1^* \times M_2^*)(x, y)$.

(3) Let $(0_{M_1}, 0_{M_2}) \in M_1 \times M_2$. Then

$$\mu_{M_1^* \times M_2^*}(0_{M_1}, 0_{M_2}) = T(\mu_{M_1^*}(0_{M_1}), \mu_{M_2^*}(0_{M_2})) = T(1, 1) = 1$$

and

$$\nu_{M_1^* \times M_2^*}(0_{M_1}, 0_{M_2}) = C(\nu_{M_1^*}(0_{M_1}), \nu_{M_2^*}(0_{M_2})) = C(0, 0) = 0.$$

$$\text{Then } (M_1^* \times M_2^*)(0_{M_1}, 0_{M_2}) = (1, 0).$$

$$\text{Hence (1)-(3) give us that } M_1^* \times M_2^* = (\mu_{M_1^* \times M_2^*}, \nu_{M_1^* \times M_2^*}) \in BFN(M_1 \times M_2).$$

□

Corollary 3.6 : Let M_i be R -modules and $M_i^* = (\mu_{M_i^*}, \nu_{M_i^*}) \in BFN(M)$ for $i = 1, 2, \dots, n$. Then

$$M_1^* \times M_2^* \times \dots \times M_n^* \in BFN(M_1 \times M_2 \times \dots \times M_n).$$

Definition 3.7 : (See [14]) Let φ be a morphism from set X into set Y such that $A = (\mu_A, \nu_A)$ and $B = (\mu_B, \nu_B)$ be two intuitionistic fuzzy sets in X and Y respectively. For all $x \in X, y \in Y$, we define

$$\varphi(A)(y) = (\varphi(\mu_A)(y), \varphi(\nu_A)(y))$$

$$= \begin{cases} (\sup\{\mu_A(x) \mid x \in X, \varphi(x) = y\}, \inf\{\nu_A(x) \mid x \in X, \varphi(x) = y\}) & \text{if } \varphi^{-1}(y) \neq \emptyset \\ (0, 1) & \text{if } \varphi^{-1}(y) = \emptyset. \end{cases}$$

$$\text{Also } \varphi^{-1}(B)(x) = (\varphi^{-1}(\mu_B)(x), \varphi^{-1}(\nu_B)(x)) = (\mu_B(\varphi(x)), \nu_B(\varphi(x))).$$

Proposition 3.8 : Let φ be an epimorphism from R -module M into R -module S . If $M^* = (\mu_{M^*}, \nu_{M^*}) \in BFN(M)$, then $\varphi(M^*) = (\mu_{\varphi(M^*)}, \nu_{\varphi(M^*)}) \in BFN(S)$.

Proof : (1) Let $y_1, y_2 \in S$. Then

$$\begin{aligned} \mu_{\varphi(M^*)}(y_1 + y_2) &= \sup\{\mu_{M^*}(x_1 + x_2) \mid x_1, x_2 \in M, \varphi(x_1) = y_1, \varphi(x_2) = y_2\} \\ &\geq \sup\{T(\mu_{M^*}(x_1), \mu_{M^*}(x_2)) \mid x_1, x_2 \in M, \varphi(x_1) = y_1, \varphi(x_2) = y_2\} \\ &= T(\sup\{\mu_{M^*}(x_1) \mid x_1 \in M, \varphi(x_1) = y_1\}, \sup\{\mu_{M^*}(x_2) \mid x_2 \in M, \varphi(x_2) = y_2\}) \\ &= T(\mu_{\varphi(M^*)}(y_1), \mu_{\varphi(M^*)}(y_2)). \end{aligned}$$

Also

$$\nu_{\varphi(M^*)}(y_1 + y_2) = \inf\{\nu_{M^*}(x_1 + x_2) \mid x_1, x_2 \in M, \varphi(x_1) = y_1, \varphi(x_2) = y_2\}$$

$$\begin{aligned}
&\leq \inf\{C(\nu_{M^*}(x_1), \nu_{M^*}(x_2)) \mid x_1, x_2 \in M, \varphi(x_1) = y_1, \varphi(x_2) = y_2\} \\
&= C(\inf\{\nu_{M^*}(x_1) \mid x_1 \in M, \varphi(x_1) = y_1\}, \inf\{\nu_{M^*}(x_2) \mid x_2 \in M, \varphi(x_2) = y_2\}) \\
&= C(\nu_{\varphi(M^*)}(y_1), \nu_{\varphi(M^*)}(y_2)).
\end{aligned}$$

Therefore

$$\varphi(M^*)(y_1 + y_2) \supseteq (T(\mu_{\varphi(M^*)}(y_1), \mu_{\varphi(M^*)}(y_2)), C(\nu_{\varphi(M^*)}(y_1), \nu_{\varphi(M^*)}(y_2))).$$

(2) Let $y \in S$ and $r \in R$. Thus

$$\begin{aligned}
\mu_{\varphi(M^*)}(ry) &= \sup\{\mu_{M^*}(rx) \mid x \in M, \varphi(rx) = ry\} \\
&\geq \sup\{\mu_{M^*}(x) \mid x \in M, \varphi(x) = y\} = \mu_{\varphi(M^*)}(y)
\end{aligned}$$

and

$$\begin{aligned}
\nu_{\varphi(M^*)}(ry) &= \inf\{\nu_{M^*}(rx) \mid x \in M, \varphi(rx) = ry\} \\
&\leq \inf\{\nu_{M^*}(x) \mid x \in M, \varphi(x) = y\} = \nu_{\varphi(M^*)}(y).
\end{aligned}$$

Then $\varphi(M^*)(ry) \supseteq \varphi(M^*)(y)$.

(3) As $\mu_{M^*}(0_M) = 1$ and $\nu_{M^*}(0_M) = 0$ so we get that

$$\mu_{\varphi(M^*)}(0_S) = \sup\{\mu_{M^*}(0_M) \mid 0_M \in M, \varphi(0_M) = 0_S\} = \sup\{1 \mid 0_M \in M, \varphi(0_M) = 0_S\} = 1$$

and

$$\nu_{\varphi(M^*)}(0_S) = \inf\{\nu_{M^*}(0_M) \mid 0_M \in M, \varphi(0_M) = 0_S\} = \inf\{0 \mid 0_M \in M, \varphi(0_M) = 0_S\} = 0$$

and then $\varphi(M^*)(0_S) = (1, 0)$.

Hence from (1) to (3) we get that $\varphi(M^*) = (\mu_{\varphi(M^*)}, \nu_{\varphi(M^*)}) \in BFN(S)$. $\square$

Proposition 3.9: Let φ be a morphism from R -module M into R -module S . If $S^* = (\mu_{S^*}, \nu_{S^*}) \in BFN(S)$, then $\varphi^{-1}(S^*) = (\mu_{\varphi^{-1}(S^*)}, \nu_{\varphi^{-1}(S^*)}) \in BFN(M)$.

Proof: (1) Let $x_1, x_2 \in M$.

As

$$\mu_{\varphi^{-1}(S^*)}(x_1 + x_2) = \mu_{S^*}(\varphi(x_1 + x_2)) = \mu_{S^*}(\varphi(x_1) + \varphi(x_2))$$

$$\geq T(\mu_{S^*}(\varphi(x_1)), \mu_{S^*}(\varphi(x_2))) = T(\mu_{\varphi^{-1}(S^*)}(x_1), \mu_{\varphi^{-1}(S^*)}(x_2))$$

and

$$\nu_{\varphi^{-1}(S^*)}(x_1 + x_2) = \nu_{S^*}(\varphi(x_1 + x_2)) = \nu_{S^*}(\varphi(x_1) + \varphi(x_2))$$

$$\leq C(\nu_{S^*}(\varphi(x_1)), \nu_{S^*}(\varphi(x_2))) = C(\nu_{\varphi^{-1}(S^*)}(x_1), \nu_{\varphi^{-1}(S^*)}(x_2))$$

$$\text{so } \varphi^{-1}(S^*)(x_1 + x_2) \supseteq (T(\mu_{\varphi^{-1}(S^*)}(x_1), \mu_{\varphi^{-1}(S^*)}(x_2)), C(\nu_{\varphi^{-1}(S^*)}(x_1), \nu_{\varphi^{-1}(S^*)}(x_2))).$$

(2) Let $x \in M$ and $r \in R$. Then

$$\mu_{\varphi^{-1}(S^*)}(rx) = \mu_{S^*}(\varphi(rx)) = \mu_{S^*}(r\varphi(x)) \geq \mu_{S^*}(\varphi(x)) = \mu_{\varphi^{-1}(S^*)}(x)$$

and

$$\nu_{\varphi^{-1}(S^*)}(rx) = \nu_{S^*}(\varphi(rx)) = \nu_{S^*}(r\varphi(x)) \leq \nu_{S^*}(\varphi(x)) = \nu_{\varphi^{-1}(S^*)}(x).$$

$$\text{Then } \varphi^{-1}(S^*)(rx) \supseteq \varphi^{-1}(S^*)(x).$$

(3) Since

$$\mu_{\varphi^{-1}(S^*)}(0_M) = \mu_{S^*}(\varphi(0_M)) = \mu_{S^*}(0_S) = 1$$

and

$$\nu_{\varphi^{-1}(S^*)}(0_M) = \nu_{S^*}(\varphi(0_M)) = \nu_{S^*}(0_S) = 0$$

$$\text{so } \varphi^{-1}(S^*)(0_M) = (1, 0).$$

Thus (1) to (3) will give us that $\varphi^{-1}(S^*) = (\mu_{\varphi^{-1}(S^*)}, \nu_{\varphi^{-1}(S^*)}) \in BFN(M)$.

References

- [1] M. T. Abu Osman, On some products of fuzzy subgroups, *Fuzzy Sets and Systems*, 24 (1987), 79-86.
- [2] C. Alsina et al., On some logical connectives for fuzzy set theory, *J. Math. Anal. Appl.*, 93(1983), 15-26.
- [3] K. Atanassov, Intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 20 (1986), 87-96.
- [4] K. T. Atanassov, New operations defined over the intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 61(1994), 137-142.
- [5] J. J. Buckley and E. Eslami, An introduction to fuzzy logic and fuzzy sets, Springer-Verlag Berlin Heidelberg GmbH, (2002).

- [6] T. Gerstenkorn and J. Manko, Bifuzzy probabilistic sets, *Fuzzy Sets and Systems*, 71 (1995), 207-214.
- [7] M. M. Gupta and J. Qi, Theory of T-norms and fuzzy inference methods, *Fuzzy Sets and Systems*, 40(1991), 431-450.
- [8] U. Hohle, Probabilistic uniformization of fuzzy topologies, *Fuzzy Sets and Systems*, 1(1978), 311-332.
- [9] D. Klaua, An early approach toward graded identity and graded membership in set theory, *Fuzzy Sets and Systems*, 161(2010), 2369-2379.
- [10] D. S. Malik, J. N. Mordeson and M. K. Sen, Fundamentals of Abstract Algebra, McGraw Hill, (1997).
- [11] P. M. Rad and L. A. Mahdavi, A note on the intersection graph of submodules of a module, *Journal of Interdisciplinary Mathematics*, 22(4) (2019), 493-502.
- [12] R. Rasuli, Fuzzy Ideals of Subtraction Semigroups with Respect to A t-norm and A t-conorm, *The Journal of Fuzzy Mathematics Los Angeles*, 24 (4) (2016), 881-892.
- [13] R. Rasuli, Fuzzy modules over a t-norm, *Int. J. Open Problems Compt. Math.*, 9 (3) (2016), 12-18.
- [14] R. Rasuli, Fuzzy Subrings over a t-norm, *The Journal of Fuzzy Mathematics Los Angeles*, 24 (4) (2016), 995-1000.
- [15] R. Rasuli, Norms over intuitionistic fuzzy subrings and ideals of a ring, *Notes on Intuitionistic Fuzzy Sets*, 22 (5) (2016), 72-83.
- [16] R. Rasuli, Norms over fuzzy Lie algebra, *Journal of New Theory*, 15(2017), 32-38.
- [17] R. Rasuli, Fuzzy subgroups on direct product of groups over a t-norm, *Journal of Fuzzy Set Valued Analysis*, 3(2017), 96-101.
- [18] R. Rasuli, Characterizations of intuitionistic fuzzy subsemirings of semirings and their homomorphisms by norms, *Journal of New Theory*, 18(2017), 39-52.
- [19] R. Rasuli, Intuitionistic fuzzy subrings and ideals of a ring under norms, LAP LAMBERT Academic publishing, 2017, ISBN: 978-620-2-06926-7.
- [20] R. Rasuli, Characterization of Q-Fuzzy subrings (Anti Q-Fuzzy Subrings) with respect to a T-norm (T-Conorms), *Journal of Information and Optimization Science*, 31(2018), 1-11.

- [21] R. Rasuli, T-Fuzzy Submodules of $R \times M$, *Journal of New Theory*, 22(2018), 92-102.
- [22] R. Rasuli, Fuzzy subgroups over a T-norm, *Journal of Information and Optimization Science*, 39(2018), 1757-1765.
- [23] R. Rasuli, Fuzzy Sub-vector Spaces and Sub-bivector Spaces under t-Norms, *General Letters in Mathematics*, 5(2018), 47-57.
- [24] R. Rasuli, Anti Fuzzy Submodules over A t -conorm and Some of Their Properties, *The Journal of Fuzzy Mathematics Los Angeles*, 27(2019), 229-236.
- [25] R. Rasuli, Artinian and Noetherian Fuzzy Rings, *Int. J. Open Problems Compt. Math.*, 12(2019), 1-7.
- [26] R. Rasuli and H. Narghi, T-Norms Over Q-Fuzzy Subgroups of Group, *Jordan Journal of Mathematics and Statistics (JJMS)*, 12(2019), 1-13.
- [27] R. Rasuli, Fuzzy equivalence relation, fuzzy congruence relation and fuzzy normal subgroups on group G over t-norms, *Asian Journal of Fuzzy and Applied Mathematics*, 7(2019), 14-28.
- [28] R. Rasuli, Norms over anti fuzzy G-submodules, *MathLAB Journal*, 2(2019), 56-64.
- [29] R. Rasuli, Norms over bifuzzy bi-ideals with operators in semigroups, *Notes on Intuitionistic Fuzzy Sets*, 25(2019), 1-11.
- [30] R. Rasuli, Norms Over Basic Operations on Intuitionistic Fuzzy Sets, *The Journal of Fuzzy Mathematics Los Angeles*, 27(3)(2019), 561-582.
- [31] R. Rasuli, T-fuzzy Bi-ideals in Semirings, *Earthline Journal of Mathematical Sciences*, 27(1)(2019), 241-263.
- [32] R. Rasuli, Norms Over Intuitionistic Fuzzy Vector Spaces, *Algebra Letters*, 1(1)(2019), 1-19.
- [33] R. Rasuli, Some Results of Anti Fuzzy Subrings Over tConorms, *MathLAB Journal*, 1(4)(2019), 25-32.
- [34] R. Rasuli, Anti Fuzzy Equivalence Relation on Rings with respect to t-conorm C, *Earthline Journal of Mathematical Sciences*, 3(1)(2020), 1-19.
- [35] R. Rasuli, Anti Fuzzy Subbigroups of Bigroups under t -conorms, *The Journal of Fuzzy Mathematics Los Angeles*, 28(1)(2020), 181-200.
- [36] R. Rasuli, t-norms over Fuzzy Multigroups, *Earthline Journal of Mathematical Sciences*, 3(2)(2020), 207-228.

- [37] R. Rasuli, Anti Q-fuzzy subgroups under t-conorms, *Earthline Journal of Mathematical Sciences*, 4(1)(2020), 13-28.
- [38] R. Rasuli, Anti Fuzzy Congruence on Product Lattices with respect to S-norms, The Second National Congress on Mathematics and Statistics Conbad Kavous University, 2020.
- [39] R. Rasuli, Direct product of fuzzy multigroups under t-norms, *Open Journal of Discrete Applied Mathematics (ODAM)*, 3(1)(2020), 75-85.
- [40] R. Rasuli, Level subsets and translations of QF ST(G), *MathLAB Journal*, 5(1)(2020), 1-11.
- [41] R. Rasuli, Conorms over anti fuzzy vector spaces, *Open Journal of Mathematical Sciences*, 4(2020), 158-167.
- [42] R. Rasuli, Intuitionistic fuzzy subgroups with respect to norms (T, S), *Eng. Appl. Sci. Lett. (EASL)*, 3(2)(2020), 40-53.
- [43] Rasul Rasuli, Mohammad Mehde Moatamedi nezhad and Hossien Naraghi, Characterization of TF(G) and direct product of it, The First National Conference on Soft Computing and Cognitive Science Conbad Kavous University, July 2020.
- [44] R. Rasuli, Anti complex fuzzy subgroups under s-norms, *Eng. Appl. Sci. Lett. (EASL)*, 3(4)(2020), 1-10.
- [45] Rasul Rasuli and Mohammad Mehde Moatamedi nezhad, Characterization of Fuzzy modules and anti fuzzy modules under norms, The First International Conference on Basic Sciences, Tehran, October 21, 2020.
- [46] Rasul Rasuli and Mohammad Mehde Moatamedi nezhad, Fuzzy subrings and anti fuzzy subrings under norms, The First International Conference on Basic Sciences, Tehran, October 21, 2020.
- [47] Yu yandong, Triangular norms and TNF-sigma algebras, *Fuzzy Sets and Systems*, 16(1985), 251-264.
- [48] L. A. Zadeh, Fuzzy sets, *Inform. Control.*, 8(1965), 338-353.

Received September, 2020

Revised January, 2021