

## Orthogonality of generalized $\Gamma$ -m-derivations in semi-prime $\Gamma$ -near-rings

Hiba A. Ahmed  
Department of Mathematics  
College of Science  
Baghdad University  
Baghdad  
Iraq

---

### Abstract

We introduced the nomenclature of orthogonal  $\Gamma$ -m-derivations and orthogonal generalized  $\Gamma$ -m-derivations in semi-prime  $\Gamma$ -near-rings and provide a few essentials and enough provision for generalized  $\Gamma$ -n-derivations in semi-prime  $\Gamma$ -near-rings by orthogonal.

---

**Subject Classification:** Primary 93A30, Secondary 49K15.

**Keywords:** Semi-prime  $\Gamma$ -near-rings,  $\Gamma$ -m-derivations, Generalized  $\Gamma$ -m-derivations, Orthogonal Generalized  $\Gamma$ -m-derivations, Orthogonal  $\Gamma$ -m-derivations.

### 1. Introduction

In this work. A  $\Gamma$ -near-ring was submitted in [3, 4, 5]. A  $\Gamma$ -near-ring  $N$  possesses the condition for  $s, r \in N$ ,  $s \Gamma N \Gamma r = 0$  gives  $s = 0$  or  $r = 0$  is said prime  $\Gamma$ -near-ring and called semi-prime if  $N$  possesses the property  $t \Gamma G \Gamma t = 0$  gives  $t = 0$  for every  $t \in G$  [4, 5]. The additional commutators are as follows;  $[s, r]_p = spr - rps$  and  $(s, r) = s+r-s-r$  designate the commutator for additive groups [2, 4]. If  $(N, +)$  is abelian,  $N$  is said to be commutative and 2-torsion-free if  $2s = 0$  gives  $s = 0$ . [1].

An additive mapping  $h: N \times N \times \dots \times N \rightarrow N$  called  $\Gamma$ -m-derivation if the relationships are:

---

E-mail: [hiba.ahmed@sc.uobaghdad.edu.iq](mailto:hiba.ahmed@sc.uobaghdad.edu.iq)

$$\begin{aligned}
h(k_1 \gamma k_1', k_2, k_3, \dots, k_m) &= h(k_1, k_2, \dots, k_m) \gamma k_1' + k_1 \gamma h(k_1', k_2, k_3, \dots, k_m) \\
h(k_1, k_2 \gamma k_2', \dots, k_m) &= h(k_1, k_2, \dots, k_m) \gamma k_2' + k_2 \gamma h(k_1, k_2', \dots, k_m) \\
h(k_1, k_2, \dots, k_m \gamma k_m') &= h(k_1, k_2, \dots, k_m) \gamma k_m' + k_m \gamma h(k_1, k_2, k_3, \dots, k_m'), \text{ for} \\
\text{every } k_{1i}, k_1', k_{2i}, k_2', \dots, k_{mi}, k_m' &\in N, \gamma \in \Gamma. [3, 4]
\end{aligned}$$

The mapping is additive  $g: N \times N \times \dots \times N \rightarrow N$  called right generalize  $\Gamma$ -m-derivation of  $N$  linked with  $\Gamma$ -m-derivation  $h$  if the eq.

$$\begin{aligned}
g(k_1 \gamma k_1', k_2, \dots, k_m) &= g(k_1, k_2, \dots, k_m) \gamma k_1' + k_1 \gamma h(k_1', k_2, \dots, k_m) \\
g(k_1, k_2 \gamma k_2', \dots, k_m) &= g(k_1, k_2, \dots, k_m) \gamma k_2' + k_2 \gamma h(k_1, k_2', \dots, k_m) \\
&\vdots \\
g(k_1, k_2, \dots, k_m \gamma k_m') &= g(k_1, k_2, \dots, k_m) \gamma k_m' + k_m \gamma h(k_1, k_2, \dots, k_m')
\end{aligned}$$

Hold for every  $k_1, k_1', k_2, k_2', \dots, k_m, k_m' \in N$  and  $\gamma \in \Gamma$ . [4]. Lastly an  $\Gamma$ -m-additive mapping  $h: N \times N \times \dots \times N \rightarrow N$  called generalized  $\Gamma$ -m-derivation of  $N$  linked with  $\Gamma$ -m-derivation  $h$  if a right and left generalized  $\Gamma$ -m-derivation of  $N$  linked with  $\Gamma$ -m-additive  $h$ . [4].

Ashraf [2] introduces the orthogonal derivations in  $\Gamma$ -ring. Park [7] introduced in semi-prime near-rings, the concept of orthogonal generalized derivations is found. Kalyan and Akhil [5] insert in semi-prime gamma near-rings the concept of orthogonal generalized derivations. In the work, we introduced the definition of orthogonal of two generalized  $\Gamma$ -m-derivations in  $\Gamma$ -near-ring and orthogonal of two  $\Gamma$ -m-derivations in semi-prime  $\Gamma$ -near-ring and the current number of essential and sufficient circumstances for generalized  $\Gamma$ -m-derivations on semi-prime  $\Gamma$ -near-rings to be orthogonal. Also, we discuss some other related results.

## 2. Preliminaries

The following lemmas are required for the development of our primary proofs. We recall  $\Gamma$ -near-ring by  $\Gamma$ -(n-r) and  $\Gamma$ -m-derivations by  $\Gamma$ -m-der.

**Lemma 2.1[5]:** When  $N$  is a 2-torsion-free semi-prime  $\Gamma$ -(n-r) and  $s, k \in N$ , then The criteria listed below equal:

- (a)  $s\alpha t\beta k = 0$  every  $t \in N, \alpha, \beta \in \Gamma$ .
- (b)  $k\alpha t\beta s = 0$  every  $t \in N, \alpha, \beta \in \Gamma$ .

(c)  $s\alpha t\beta k + k\alpha t\beta s = 0$  every  $t \in N$ ,  $\alpha, \beta \in \Gamma$ .

If the above of the three conditions listed are met, thus  $s\Gamma k = k\Gamma s = 0$ .

**Lemma 2.2 [3, 4]:** When  $N$  be a  $\Gamma$ -( $n$ - $r$ ) and  $h$  be  $\Gamma$ - $m$ -der., thereafter for every  $k_1, k'_1, \dots, k_m, s \in N$  and  $\gamma, \beta \in \Gamma$ .

$$\begin{aligned} & (h(ik'_1, k_2, \dots, k_m)\gamma k'_1 + k_1\gamma h(k'_1, ik_2, \dots, ik_m))\beta s = \\ & h(ik_1, k_2, \dots, k_m)\gamma k'_1\beta s + k_1\gamma h(k'_1, k_2, \dots, k_m)\beta s \end{aligned}$$

**Lemma 2.3 [4]:** When  $N$  is a  $\Gamma$ -( $n$ - $r$ ),  $g$  be a right generalized  $\Gamma$ - $m$ -der. linked with  $\Gamma$ - $m$ -der.  $h$  of  $N$  and for every  $k_1, k_2, \dots, k_m, k, r \in N$  and  $\gamma, \beta \in \Gamma$ . Then

$$\begin{aligned} & (g(ik_1, k_2, \dots, ik_m)\gamma k + k_1\gamma h(ik_1, k_2, \dots, k_m))\beta r = \\ & g(k_1, ik_2, k_3, \dots, k_m)\gamma k\beta r + k_1\gamma h(k_1, k_2, \dots, k_m)\beta r \end{aligned}$$

**Definition 2.4:** Let  $h_1$  and  $h_2$  be two  $\Gamma$ - $n$ -derivations of  $\Gamma$ -( $n$ - $r$ ).  $h_1$  and  $h_2$  are called orthogonal if  $h_1(k_1, k_2, \dots, k_m)\Gamma N\Gamma h_2(v_1, v_2, \dots, v_m) = h_2(v_1, v_2, \dots, v_m)\Gamma N\Gamma h_1(k_1, k_2, k_3, \dots, k_m) = 0$  for every  $k_1, k_2, k_3, \dots, k_m, v_1, v_2, \dots, v_m \in N$ .

**Definition 2.5:** Let  $g_1$  and  $g_2$  be two generalized  $\Gamma$ - $m$ -der. of  $\Gamma$ -( $n$ - $r$ ).  $g_1$  and  $g_2$  are called orthogonal if  $g_1(k_1, k_2, \dots, k_m)\Gamma N\Gamma g_2(v_1, v_2, \dots, v_m) = g_2(v_1, v_2, \dots, v_m)\Gamma N\Gamma g_1(ik_1, k_2, k_3, \dots, k_m) = 0$  for every  $ik_1, k_2, k_3, \dots, k_m, v_1, v_2, \dots, v_m \in N$ .

### 3. Main Results

**Lemma 3.1:** When  $N$  is 2- torsion-free semi-prime  $\Gamma$ -( $n$ - $r$ ),  $g_1, g_2$  be two generalized  $\Gamma$ - $m$ -der. of  $N$  linked with  $\Gamma$ - $m$ -der.  $h_1$  and  $h_2$  of  $N$  progression. If  $g_1$  and  $g_2$  are orthogonal, thus go to the next conditions are:

- (a)  $g_1(k_1, k_2, \dots, k_m)\Gamma g_2(v_1, v_2, \dots, v_m) = g_2(k_1, k_2, \dots, k_m)\Gamma g_1(v_1, v_2, \dots, v_m) = 0$  for every  $k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N$ .
- (b)  $h_1$  and  $g_2$  are orthogonal and  $h_1(k_1, k_2, \dots, k_m)\Gamma g_2(v_1, v_2, \dots, v_m) = g_2(v_1, v_2, \dots, v_m)\Gamma h_1(ik_1, k_2, \dots, k_m) = 0$  for every  $ik_1, k_2, k_3, \dots, k_m, v_1, v_2, \dots, v_m \in N$ .
- (c)  $h_2$  and  $g_1$  are orthogonal and  $h_2(k_1, k_2, \dots, k_m)\Gamma g_1(v_1, v_2, \dots, v_m) = g_1(v_1, v_2, \dots, v_m)\Gamma h_2(ik_1, k_2, \dots, k_m) = 0$  for every  $ik_1, k_2, k_3, \dots, k_m, v_1, v_2, \dots, v_m \in N$ .

**Proof:** (a) Since  $g_1$  and  $g_2$  orthogonality, obtain  $g_1(k_1, k_2, \dots, k_m)\alpha k\beta g_2(v_1, v_2, \dots, v_m) = 0$  for every  $k, k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N$  and  $\alpha, \beta \in \mu$ . By Lemma 2.1 obtain  $g_1(k_1, k_2, \dots, k_m)\alpha g_2(v_1, v_2, \dots, v_m) = g_2(k_1, k_2, \dots, k_m)\alpha g_1(v_1, v_2, \dots, v_m) = 0$  (b) By (a) we get  $g_1(k_1, k_2, \dots, k_m)\alpha g_2(v_1, v_2, \dots, v_m) = 0$  for every  $k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N$  and  $\alpha \in \Gamma$ . Replacing  $k_1$  by  $k_1\beta k_1'$ , where  $k_1' \in N$ , in above eq. and taking Lemma 2.3 gives

$$\begin{aligned} 0 &= g_1(k_1'\beta k_1, k_2, \dots, k_m)\alpha g_2(v_1, v_2, \dots, v_m) \\ &= g_1(k_1', k_2, \dots, k_m)\beta t_1\alpha g_2(v_1, v_2, \dots, v_m) + t_1'\beta h_1(t_1, t_2, \dots, t_n)\alpha g_2(v_1, v_2, \dots, v_m) \\ &= k_1'\beta h_1(k_1, k_2, \dots, k_m)\alpha g_2(v_1, v_2, \dots, v_m). \end{aligned}$$

It follows from the semiprimeness of  $N$  that  $h_1(k_1, k_2, \dots, k_m)\alpha g_2(v_1, v_2, \dots, v_m) = 0$  for every  $k_1, k_1', k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N$  and  $\alpha \in \Gamma$ .

Thus  $h_1(k_1, k_2, \dots, k_m)\Gamma g_2(v_1, v_2, \dots, v_m) = 0$ . This is the first part of (b).

To show the second part we replace  $k_1$  by  $k_1\alpha k_1'$ , where  $k_1' \in N$ , in above eq. and using Lemma 2.2, gives

$$\begin{aligned} 0 &= h_1(k_1\alpha k_1', k_2, \dots, k_m)\beta g_2(v_1, v_2, \dots, v_m) \\ &= h_1(k_1, k_2, \dots, k_m)\alpha k_1'\beta g_2(v_1, v_2, \dots, v_m) + k_1\alpha h_1(k_1', k_2, \dots, k_m)\beta g_2(v_1, v_2, \dots, v_m) \\ &= h_1(k_1, k_2, \dots, k_m)\alpha t_1'\beta g_2(v_1, v_2, \dots, v_m). \text{ By Lemma 2.1 we obtain} \\ &g_2(v_1, v_2, \dots, v_m)\gamma h_1(ik_1, tk_2, k_3, \dots, k_m) = 0 \text{ for every } ik_1, k_2, k_3, \dots, ik_m, v_1, v_2, \dots, v_m \in N \text{ and } \gamma \in \Gamma. \end{aligned}$$

**Theorem 3.2:** When  $N$  be a 2-torsion-free semi-prime  $\Gamma$ -( $n$ -r). Suppose that  $g_1$  and  $g_2$  are two generalized  $\Gamma$ - $n$ -der. of  $N$  linked with  $\Gamma$ - $m$ -der.  $h_1$  and  $h_2$  respectively, the next term are equivalent:

- (a)  $g_1$  and  $g_2$  are orthogonal.
- (b)  $g_1(ik_1, k_2, \dots, k_m)\gamma g_2(v_1, v_2, \dots, v_m) = h_1(k_1, k_2, \dots, k_m)\gamma g_2(v_1, v_2, \dots, v_m) = 0$  for every  $k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N, \gamma \in \Gamma$ .
- (c)  $g_1(k_1, k_2, \dots, k_m)\gamma g_2(v_1, v_2, \dots, v_m) = h_1(k_1, k_2, \dots, k_m)\gamma h_2(v_1, v_2, \dots, v_m) = 0$  and  $h_1(g_2(k_1, k_2, \dots, k_m), v_2, \dots, v_m) = h_1(h_2(k_1, k_2, \dots, k_m), v_2, \dots, v_m) = 0$

for every  $k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N$ .

**Proof:** The portion (a)  $\Rightarrow$  (b) and (a)  $\Rightarrow$  (c) are established by Lemma 3.1.

The portion, (a)  $\Rightarrow$  (c) is immediate from Lemma 3.1.

(b)  $\Rightarrow$  (a) We have, for every  $k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N$  and  $\alpha \in \Gamma$ .

$g_1(k_1, k_2, \dots, k_m) \gamma g_2(v_1, v_2, \dots, v_m) = h_1(k_1, k_2, \dots, k_m) \gamma h_2(v_1, v_2, \dots, v_m) = 0$   
 Replacing  $k_1$  by  $k_1 \alpha k_1'$  in the equation  $g_1(k_1, k_2, \dots, k_m) \gamma g_2(v_1, v_2, \dots, v_m) = 0$  gives

$g_1(t_1 \alpha t_1', t_2, \dots, t_n) \gamma g_2(r_1, r_2, \dots, r_n) = 0$ . Using hypothesis and Lemma 2.3 in last eq. obtain

$$\begin{aligned} 0 &= g_1(k_1, k_2, \dots, k_m) \alpha k_1' \gamma g_2(v_1, v_2, \dots, v_m) + k_1 \alpha h_1(k_1', k_2, \dots, k_m) \gamma g_2(v_1, v_2, \dots, v_m) \\ &= g_1(k_1, k_2, \dots, k_m) \alpha k_1' \gamma g_2(v_1, v_2, \dots, v_m) \text{ for every } k_1, k_1', k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N, \alpha, \gamma \in \Gamma. \end{aligned}$$

(c)  $\Rightarrow$  (a) Since  $h_1(g_2(k_1, k_2, \dots, k_m), v_2, \dots, v_m) = h_1(h_2(k_1, k_2, \dots, k_m), v_2, \dots, v_m) = 0$  for every  $k_1, k_2, \dots, k_m, v_2, \dots, v_m \in N$ . Replacing  $t_1$  by  $t_1 \alpha t_1'$  in the equation  $h_1(g_2(k_1, k_2, \dots, k_m), v_2, \dots, v_m) = 0$ , we get

$$\begin{aligned} 0 &= h_1(g_2(k_1 \alpha k_1', k_2, \dots, k_m), v_2, \dots, v_m) \\ &= h_1(g_2(k_1, k_2, \dots, k_m) \alpha k_1', v_2, \dots, v_m) + h_1(k_1 \alpha h_2(k_1', k_2, \dots, k_m), v_2, \dots, v_m) \\ &= h_1(g_2(k_1, k_2, \dots, k_m), v_2, \dots, v_m) \alpha k_1' + g_2(k_1, k_2, \dots, k_m) \alpha h_1(k_1', v_2, \dots, v_m) \\ &\quad + h_1(k_1, v_2, \dots, v_m) \alpha h_2(ik_1', ik_2, k_3, \dots, k_m) + k_1 \alpha h_1(h_2(ik_1', k_2, k_3, \dots, 0k_m), v_2, \dots, v_m). \end{aligned}$$

for every  $k_1, k_1', k_2, k_3, \dots, k_m, v_2, \dots, v_m \in N$ . Hypothesis is being used once more obtain  $g_2(k_1, k_2, \dots, k_m) \alpha h_1(k_1', v_2, \dots, v_m) = 0$ . Replacing  $k_1'$  by  $k_1' \gamma z$ , where  $z \in N$ , in above eq. and using it once again, gives

$$\begin{aligned} 0 &= g_2(k_1, k_2, \dots, k_m) \alpha h_1(k_1' \gamma z, v_2, \dots, v_m) \\ &= g_2(k_1, k_2, \dots, k_m) \alpha h_1(k_1', v_2, \dots, v_m) \gamma z + g_2(k_1, k_2, \dots, k_m) \alpha k_1' \gamma h_1(z, v_2, \dots, v_m) \\ &= g_2(k_1, k_2, \dots, k_m) \alpha k_1' \gamma h_1(z, v_2, \dots, v_m), \text{ for every } z, k_1, k_1', k_2, \dots, k_m, v_2, \dots, v_m \in N. \end{aligned}$$

Hence by Lemma 2.1 we get  $h_1(z, v_2, \dots, v_m) \alpha g_2(k_1, k_2, \dots, k_m) = 0$ , for every  $k_1, k_2, \dots, k_m, z, v_2, \dots, v_m \in N$ . Then (a) follows from (b)

**Theorem 3.3:** When  $N$  a 2-torsionfree semi-prime  $\Gamma$ -( $n$ - $r$ ),  $g$  be a generalized  $\Gamma$ - $m$ -der. of  $N$  linked with  $\Gamma$ - $m$ -der.  $h$  of  $N$ . If  $g(k_1, k_2, \dots, k_m) \Gamma g(v_1, v_2, \dots, v_m) = 0$  for every  $k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N$ , then  $g = h = 0$ .

**Proof:** By supposition have

$$g(k_1, k_2, \dots, k_m) \gamma g(v_1, v_2, \dots, v_m) = 0, \text{ for every } k_1, k_2, \dots, k_m, v_1, v_2, \dots, v_m \in N, \alpha \in \Gamma. (1)$$

$$\begin{aligned} &\text{Taking } v_1 \text{ by } v_1 \alpha v_1' \text{ in above eq. (1) and us it again, implies that} \\ 0 &= g(k_1, k_2, \dots, k_m) \gamma g(v_1 \alpha v_1', v_2, \dots, v_m) \\ &= g(k_1, k_2, \dots, k_m) \gamma g(v_1, v_2, \dots, v_m) \alpha v_1' + g(k_1, k_2, \dots, k_m) \gamma v_1 \alpha h(v_1', v_2, \dots, v_m) \\ &= g(k_1, k_2, \dots, k_m) \gamma v_1 \alpha h(v_1', v_2, \dots, v_m) \end{aligned}$$

$$\text{By Lemma 2.1 obtain } h(v_1', v_2, \dots, v_m) \gamma g(k_1, k_2, \dots, k_m) = 0 \quad (2)$$

Replacing  $k_1$  by  $k_1\beta v_1'$  in eq. (2) and when we use it again, give

$$\begin{aligned}
 0 &= h(v_1', v_2', \dots, v_m) \gamma g(k_1\beta v_1', k_2, \dots, k_m) \\
 &= h(v_1', v_2', \dots, v_m) \gamma g(k_1, k_2, \dots, k_m) \beta v_1' + h(v_1', \dots, v_m) \gamma k_1 \beta h(v_1', v_2, v_3, \dots, \\
 &\quad v_m) \\
 &= h(v_1', v_2', \dots, v_m) \gamma k_1 \beta h(v_1', k_2, \dots, k_m) \text{ for every } k_1, k_2, \dots, k_m, v_1', v_2', \dots, \\
 &\quad v_m \in N.
 \end{aligned}$$

In particular  $h(v_1', v_2', v_3, \dots, v_m) \gamma v_1 \beta h(v_1', v_2, v_3, \dots, v_m)$ . Semi-primeness of  $N$  give  $h = 0$ . By hypothesis and Lemma 2.3, obtain

$0 = g(k_1\beta v, k_2, \dots, k_m) \gamma g(v_1, v_2, \dots, v_m) = g(k_1, k_2, \dots, k_m) \beta v \gamma g(v_1, v_2, \dots, v_m) + k_1 \beta h(v, k_2, \dots, k_m) \gamma h(v_1, v_2, \dots, v_m)$ . By eq. (1) we conclude that  $g(k_1, k_2, \dots, k_m) \beta v \gamma g(v_1, v_2, \dots, v_m) = 0$  for every  $k_1, k_2, \dots, k_m, v, v_1, v_1', v_2, \dots, v_m \in N$ . Thus we get  $g = 0$  by the semi-primeness of  $N$ .

#### 4. Conclusion

In the present work, we defined two generalized orthogonality  $\Gamma$ - $m$ -der. In  $\Gamma$ -( $n$ - $r$ ) also orthogonal of two  $\Gamma$ - $m$ -der. in semi-prime  $\Gamma$ -( $n$ - $r$ ). And give characterizations of  $\Gamma$ - $m$ -der as a product of two  $\Gamma$ - $m$ -der. in 2-torsion-free semi-prime  $\Gamma$ -( $n$ - $r$ ).

#### References

- [1] Asraf, M. and Siddeeqe, M. A., "On permuting  $n$ -derivations in near-rings", *Commun. Kor. Math. Soc.*, Vol. 28, No. 4, pp. 697–707 (2013).
- [2] Ashraf M. and M.R.Jamal, Orthogonal Derivations in  $\Gamma$ -rings, *Advances in Algebra*, 3(1), 1-6 (2010).
- [3] Hiba A. Ahmed, Right  $G$ - $n$ -derivations in prime  $G$ -near-rings, *Int. J. Nonlinear Anal. Appl.* 12, No. 2, 1653–1658 (2021)
- [4] Hiba A. Ahmed, Abdulrahman H. M., Generalized  $G$ - $n$ -Derivations on Prime  $G$ -Near-Rings, *J. Phys.: Conf.* Vol. 1963, 012123 (2021).
- [5] Kalyan K., Akhil C. and Isamiddin S., Orthogonal Generalized derivations in semiprime gamma near-rings, *International Journal of Algebra*, Vol. 6, No. 23, 1127-1134 (2012).
- [6] Madhuchelvi L., On Generalized Derivations in  $G$ -Near-Ring *Int Jr.of Mathematical Sciences and Applications*, ISSN N0, 6 (1) 2230-9888 (2016).
- [7] Park, K.H. and Jung, Y.S. Semiprime near-rings with orthogonal derivations, *Korea Soc. J. Math.*, 13(40), pp.303-310 (2006).

- [8] Shubham Sharma & Ahmed J. Obaid, Mathematical modelling, analysis and design of fuzzy logic controller for the control of ventilation systems using MATLAB fuzzy logic toolbox, *Journal of Interdisciplinary Mathematics*, 23:4, 843-849 (2020), DOI: 10.1080/09720502.2020.1727611.
- [9] D. White. An orthogonal set of predictors for wins in major league baseball, *Journal of Interdisciplinary Mathematics*, 13:3, 309-319 (2010), DOI: 10.1080/09720502.2010.10700703.

*Received May, 2022*

*Revised June, 2022*