

Pre – supra v – open and some related concepts

Zaman R. Ghazi *

Luma S. Abdalbaqi †

*Department of Mathematics
College of Education for Women
Tikrit University
Salah Al-Din
Iraq*

Abstract

The main goals of the paper is presenting a comprehensive study about the concept of pre-supra v – open set and giving many new results. Studying the properties of some concepts relative to the supra v – open set such as semi – supra v – open, regular – supra v – open, α – supra v – open and β – supra v – open as well as these concepts are generalizations of a supra v – open set.

Subject Classification: 00A05, 00A06.

Keywords: Algebra, Topology, Cryptography, Open set.

1. Introduction

Originally coming from questions in analysis [1] and differential geometry by now topology permeates mostly every field of math including algebra [2],[3], combinatorics, logic and plays a fundamental role in algebraic geometry as we know it today.

The concept of semi-open set was studied by Levine [4] in 1963 which is a generalization of an open set.

Njastad in 1965 [5] was first introduced α – open set concept as a generalization of open set.

The concept of supra topological space was proposed by Mashhour [6] in 1983 as a generalization of the concept of topological space.

There are many authors are studied some concepts related to topological spaces for example Abdalbaqi, see [7] and [8] .

* E-mail: zaman.raeed23@st.tu.edu.iq (Corresponding Author)

† E-mail: lumahhany1977@tu.edu.iq

In 2023 Raoof and Abdalbaqi are introduced the concept of supra ν – open set where $\{\mathfrak{F}_\kappa\}_{\kappa \in J}$, $\kappa \geq 2$, supra topologies on $\mathcal{X}$ and $\mathcal{V} \subseteq \mathcal{X}$. Then $\mathcal{V}$ is supra ν – open set in $\mathcal{X}$ if $\exists T \in \bigcap_{\kappa \in J} \mathfrak{F}_\kappa \ni \phi \neq T \subseteq \mathcal{V}$ whenever $\mathcal{V} \neq \phi$ & $T = \phi$ when $\mathcal{V} = \phi$. $\mathcal{S}\nu O_{\mathcal{X}}$ denotes to the set of all supra ν – open sets.

2. The main results

Definition 2.1: Assume $(\mathcal{X}, \mathcal{S}\nu O_{\mathcal{X}})$ is a supra ν – space. Then a subset $\mathcal{V}$ of $\mathcal{X}$ is a pre – supra ν – open if $\mathcal{V} \subseteq \text{Int}_{\mathcal{S}\nu}(\text{cl}_{\mathcal{S}\nu}(\mathcal{V}))$.

$\mathcal{P}(\mathcal{S}\nu O_{\mathcal{X}})$ is a set of all pre – supra ν – opensets.

The complement of an pre – supra ν – open is called pre – supra ν – closed. $\mathcal{P}(\mathcal{S}\nu C_{\mathcal{X}})$ is a set of all pre – supra ν – closed.

Proposition 2.2: Every supra ν – open is a pre – supra ν – open.

Definition 2.3: Assume $(\mathcal{X}, \mathcal{S}\nu O_{\mathcal{X}})$ is a supra ν – space & $\mathcal{V} \subseteq \mathcal{X}$, then $\mathcal{V}$ is a regular – supra ν – open if $\mathcal{V} = \text{Int}_{\mathcal{S}\nu}(\text{cl}_{\mathcal{S}\nu}(\mathcal{V}))$.

- the set of all regular – supra ν – open is known as $\mathcal{R}(\mathcal{S}\nu O_{\mathcal{X}})$.
- the complement of regular – supra ν – open is called regular – supra ν – closed.
- the set of all regular – supra ν – closed is known as $\mathcal{R}(\mathcal{S}\nu C_{\mathcal{X}})$.

Proposition 2.4: Every regular – supra ν – open is a supra ν – open.

Proposition 2.5: Every regular – ν – open is a pre – supra ν – open.

Proof: Assume that $\mathcal{V}$ is a regular – ν – open, then by Proposition (2.4), we have $\mathcal{V}$ is a supra ν – open, hence by Proposition (2.2), we get $\mathcal{V}$ is a pre – supra ν – open.

Definition 2.6: Assume $(\mathcal{X}, \mathcal{S}\nu O_{\mathcal{X}})$ is a supra ν – space & $\mathcal{V} \subseteq \mathcal{X}$, then $\mathcal{V}$ is a β – supra ν – open if $\mathcal{V} \subseteq \text{cl}_{\mathcal{S}\nu}(\text{Int}_{\mathcal{S}\nu}(\text{cl}_{\mathcal{S}\nu}(\mathcal{V})))$.

- the set of all β – supra ν – open is denoted by $\beta(\mathcal{S}\nu O_{\mathcal{X}})$.
- the complement of an β – supra ν – open is said to be β – supra ν – closed.
- the set of all β – supra ν – closed is denoted by $\beta(\mathcal{S}\nu C_{\mathcal{X}})$.

Proposition 2.7: Every supra ν – open is a β – supra ν – open.

Proof: Assume that $\mathcal{V}$ is a supra ν – open. then $\mathcal{V} \subseteq \text{cl}_{\mathcal{S}\nu}(\mathcal{V})$, hence $\text{Int}_{\mathcal{S}\nu}(\mathcal{V}) \subseteq \text{Int}_{\mathcal{S}\nu}(\text{cl}_{\mathcal{S}\nu}(\mathcal{V}))$, but $\mathcal{V}$ is a supra ν – open, then $\text{Int}_{\mathcal{S}\nu}(\mathcal{V}) = \mathcal{V}$, Thus $\mathcal{V} \subseteq \text{Int}_{\mathcal{S}\nu}(\text{cl}_{\mathcal{S}\nu}(\mathcal{V}))$. Hence, $\text{cl}_{\mathcal{S}\nu}(\mathcal{V}) \subseteq \text{cl}_{\mathcal{S}\nu}(\text{Int}_{\mathcal{S}\nu}(\text{cl}_{\mathcal{S}\nu}(\mathcal{V})))$ by Theorem (1.39):(1). So, we have $\mathcal{V} \subseteq \text{cl}_{\mathcal{S}\nu}(\text{Int}_{\mathcal{S}\nu}(\text{cl}_{\mathcal{S}\nu}(\mathcal{V})))$.

Therefore, $\mathcal{V}$ is a β – supra ν – open.

Proposition 2.8: Every regular – $\mathcal{V}$ – open is a β – supra $\mathcal{V}$ – open.

Theorem 2.9: Assume $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space & $\phi \neq \mathcal{V} \subseteq X$. If $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq \phi$, then $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \mathcal{V}$.

Proof: When $\mathcal{V} = X$, we done. Thus, suppose $\mathcal{V} \neq X$.

Now, we have, $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})$ is the largest supra $\mathcal{V}$ – open set contained in $\mathcal{V}$. That is $\mathcal{V} \supseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \in \mathcal{S}\mathcal{V}O_X$. Hence, there exist $T \in \bigcap_{\kappa \in I} \mathfrak{F}_\kappa$ such that $\phi \neq T \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})$, thus $T \subseteq \mathcal{V}$, which implies that $\mathcal{V} \in \mathcal{S}\mathcal{V}O_X$.

Therefore, $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \mathcal{V}$.

Theorem 2.10: Suppose $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space & $\mathcal{V} \subset X$. If $\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq X$, then $\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \mathcal{V}$.

Proposition 2.11: If $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space, then every pre – supra $\mathcal{V}$ – open is a β – supra $\mathcal{V}$ – open.

Theorem 2.12: Assume $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space, then every β – supra $\mathcal{V}$ – open is a pre – supra $\mathcal{V}$ – open.

Proof: If $\mathcal{V}$ is β – supra $\mathcal{V}$ – open, so $\mathcal{V} \subseteq \text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V})))$. If $\mathcal{V} = \phi$ or $\mathcal{V} = X$, then $\mathcal{V}$ is a pre – supra $\mathcal{V}$ – open. Thus, let $\phi \neq \mathcal{V} \neq X$.

Now, either $\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = X$ or $\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq X$.

If $\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = X$, then $\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V})) = X$, hence $\mathcal{V} \subset \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))$, therefore $\mathcal{V}$ is a pre – supra $\mathcal{V}$ – open. If $\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq X$, since $\mathcal{V} \subset X$, then $\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \mathcal{V}$ by Theorem (2.10). hence $\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V})) = \text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})$.

Now, either $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \phi$ or $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq \phi$.

If $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \phi$, then $\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})) = \phi$, that is $\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))) = \phi$, but $\mathcal{V} \subseteq \text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V})))$ by hypothesis, then we get $\mathcal{V} \subseteq \phi$ this is contradiction. So, $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq \phi$, hence by Theorem (2.9) we get $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \mathcal{V}$. Now, since $\mathcal{V} \subseteq \text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V})))$, then we get $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))))$, but $\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V})) = \text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})$ and $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \mathcal{V}$, hence $\mathcal{V} \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))$, therefore $\mathcal{V}$ is a pre – supra $\mathcal{V}$ – open.

Proposition 2.13: $\mathcal{P}(\mathcal{S}\mathcal{V}O_X) = \beta(\mathcal{S}\mathcal{V}O_X)$.

Proposition 2.14: $\mathcal{R}(\mathcal{S}\mathcal{V}O_X) \subset \mathcal{P}(\mathcal{S}\mathcal{V}O_X)$.

Proposition 2.15: $\mathcal{R}(\mathcal{S}\mathcal{V}O_X) \subset \mathcal{S}\mathcal{V}O_X$.

Proposition 2.16: Assume that $(X, \mathcal{S}\mathcal{V}O_X)$ be a supra $\mathcal{V}$ – space. Then $\mathcal{S}\mathcal{V}O_X \subset \mathcal{P}(\mathcal{S}\mathcal{V}O_X)$.

Definition 2.17: Suppose that $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space. Then a subset $\mathcal{V}$ of X is said to be α – supra $\mathcal{V}$ – open if $\mathcal{V} \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})))$.

the set of all α – supra $\mathcal{V}$ – open is denoted by $\alpha(\mathcal{S}\mathcal{V}O_X)$.

complement for an α – supra $\mathcal{V}$ – open named α – supra $\mathcal{V}$ – closed.

A set for every α – supra $\mathcal{V}$ – closed is $\alpha(\mathcal{S}\mathcal{V}C_X)$.

Proposition 2.18: Every supra $\mathcal{V}$ – open is a α – supra $\mathcal{V}$ – open.

Theorem 2.19: Let $(X, \mathcal{S}\mathcal{V}O_X)$ be a supra $\mathcal{V}$ – space. Then every α – supra $\mathcal{V}$ – open is a supra $\mathcal{V}$ – open.

Proof: Assume $\mathcal{V}$ be an α – supra $\mathcal{V}$ – open, so

$$\mathcal{V} \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))).$$

when $\mathcal{V} = \phi$ or $\mathcal{V} = X$, hence $\mathcal{V}$ is supra $\mathcal{V}$ – open.

Thus, let $\phi \neq \mathcal{V} \neq X$.

Now, either $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \phi$ or $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq \phi$.

If $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \phi$, then $\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})) = \phi$, hence

$\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))) = \phi$, thus $\mathcal{V} \subseteq \phi$ contradiction.

Hence, $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \neq \phi$, since $\mathcal{V} \neq \phi$, then by Theorem (2.9) we get $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) = \mathcal{V}$. Therefore $\mathcal{V}$ is a supra $\mathcal{V}$ – open.

Proposition 2.20: $\alpha(\mathcal{S}\mathcal{V}O_X) = \mathcal{S}\mathcal{V}O_X$.

Proposition 2.21: Every regular – supra $\mathcal{V}$ – open is a α – supra $\mathcal{V}$ – open.

Proposition 2.22: Assume $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space. then $\mathcal{R}(\mathcal{S}\mathcal{V}O_X) \subset \alpha(\mathcal{S}\mathcal{V}O_X)$.

Proposition 2.23: Every α – supra $\mathcal{V}$ – open is a pre – supra $\mathcal{V}$ – open.

Proof: Assume that $\mathcal{V}$ is a α – supra $\mathcal{V}$ – open. Then $\mathcal{V} \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V})))$. Now, we have $\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}) \subseteq \mathcal{V}$, hence $\text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))) \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))$, thus $\mathcal{V} \subseteq \text{Int}_{\mathcal{S}\mathcal{V}}(\text{cl}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))$. Therefore, $\mathcal{V}$ is a pre – supra $\mathcal{V}$ – open.

Proposition 2.24: $\alpha(\mathcal{S}\mathcal{V}O_X) \subset \mathcal{P}(\mathcal{S}\mathcal{V}O_X)$.

Proposition 2.25: Every α – supra $\mathcal{V}$ – open is a β – supra $\mathcal{V}$ – open.

Proposition 2.26: Assume $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space. then $\alpha(\mathcal{S}\mathcal{V}O_X) \subset \beta(\mathcal{S}\mathcal{V}O_X)$.

Definition 2.27: Suppose that $(X, \mathcal{S}\mathcal{V}O_X)$ is a supra $\mathcal{V}$ – space. Then a subset $\mathcal{V}$ of X is called semi – supra $\mathcal{V}$ – open if $\mathcal{V} \subseteq \text{cl}_{\mathcal{S}\mathcal{V}}(\text{Int}_{\mathcal{S}\mathcal{V}}(\mathcal{V}))$.

- the set of all semi – supra $\mathcal{V}$ – open is denoted by $\mathcal{S}(\mathcal{S}\mathcal{V}O_X)$.

- the complement of semi – supra $\mathcal{V}$ – open is called semi – supra $\mathcal{V}$ – closed.
- the set of all semi – supra $\mathcal{V}$ – closed is denoted by $\mathcal{G}(\mathcal{SV}C_X)$.

Proposition 2.28: Every supra $\mathcal{V}$ – open is a semi – supra $\mathcal{V}$ – open.

Theorem 2.29: If $(X, \mathcal{SV}O_X)$ is a supra $\mathcal{V}$ – space, then every semi – supra $\mathcal{V}$ – open is a supra $\mathcal{V}$ – open.

Proof: Assume $\mathcal{V}$ is semi – supra $\mathcal{V}$ – open, so when $\mathcal{V} = \phi$ or $\mathcal{V} = X$, we get $\mathcal{V}$ is supra $\mathcal{V}$ – open.

Thus, let $\phi \neq \mathcal{V} \neq X$.

Now, either $\text{Int}_{\mathcal{SV}}(\mathcal{V}) = \phi$ or $\text{Int}_{\mathcal{SV}}(\mathcal{V}) \neq \phi$.

If $\text{Int}_{\mathcal{SV}}(\mathcal{V}) = \phi$, then $\text{cl}_{\mathcal{SV}}(\text{Int}_{\mathcal{SV}}(\mathcal{V})) = \phi$.

Hence $\mathcal{V} \subseteq \phi$ contradiction.

Thus, $\text{Int}_{\mathcal{SV}}(\mathcal{V}) \neq \phi$, since $\mathcal{V} \neq \phi$, then we get $\text{Int}_{\mathcal{SV}}(\mathcal{V}) = \mathcal{V}$. So $\mathcal{V}$ is supra $\mathcal{V}$ – open.

Proposition 2.30: $\mathcal{G}(\mathcal{SV}O_X) = \mathcal{SV}O_X$.

Theorem 2.31: Let $(X, \mathcal{SV}O_X)$ be a supra $\mathcal{V}$ – space & $\mathcal{V} \subseteq X$. Then $\mathcal{V}$ is a α – supra $\mathcal{V}$ – open iff $\mathcal{V}$ is a semi – supra $\mathcal{V}$ – open.

Conclusion

The main results are obtained are:

1. Every supra $\mathcal{V}$ – open is a pre – supra $\mathcal{V}$ – open.
2. Every β – supra $\mathcal{V}$ – open is a pre – supra $\mathcal{V}$ – open.
3. Every supra $\mathcal{V}$ – open is a α – supra $\mathcal{V}$ – open.
4. Every supra $\mathcal{V}$ – open is a semi – supra $\mathcal{V}$ – open.
5. Every regular – supra $\mathcal{V}$ – open is a supra $\mathcal{V}$ – open.
6. Assume $(X, \mathcal{SV}O_X)$ is a supra $\mathcal{V}$ – space & $\phi \neq \mathcal{V} \subseteq X$. If $\text{Int}_{\mathcal{SV}}(\mathcal{V}) \neq \phi$, then $\text{Int}_{\mathcal{SV}}(\mathcal{V}) = \mathcal{V}$.
7. Suppose $(X, \mathcal{SV}O_X)$ is a supra $\mathcal{V}$ – space & $\mathcal{V} \subset X$. If $\text{cl}_{\mathcal{SV}}(\mathcal{V}) \neq X$, then $\text{cl}_{\mathcal{SV}}(\mathcal{V}) = \mathcal{V}$.
8. Assume $(X, \mathcal{SV}O_X)$ is a supra $\mathcal{V}$ – space, then every β – supra $\mathcal{V}$ – open is a pre – supra $\mathcal{V}$ – open.
9. If $(X, \mathcal{SV}O_X)$ is a supra $\mathcal{V}$ – space, then every semi – supra $\mathcal{V}$ – open is a supra $\mathcal{V}$ – open.
10. Assume $(X, \mathcal{SV}O_X)$ is a supra $\mathcal{V}$ – space. Then $\mathcal{G}(\mathcal{SV}O_X) = \mathcal{SV}O_X$.
11. Let $(X, \mathcal{SV}O_X)$ be a supra $\mathcal{V}$ – space & $\mathcal{V} \subseteq X$. Then $\mathcal{V}$ is a α – supra $\mathcal{V}$ – open iff $\mathcal{V}$ is a semi – supra $\mathcal{V}$ – open.

References

- [1] M. H. Alobaidi, I. S. Ahmed, and S. H. Asaad, "Study of the generalized P-contractions on banach spaces and uniqueness for stability fixed points," *Int. J. Nonlinear Anal. Appl.*, vol. 12, no. 2, pp. 1903–1908 (2021), doi: 10.22075/IJNAA.2021.5324.
- [2] A. S. Mohammed, I. S. Ahmed, and S. H. Asaad, "Study of the rings in which each element express as the sum of an idempotent and pure," *Int. J. Nonlinear Anal. Appl.*, vol. 12, no. 2, pp. 1719–1724 (2021), doi: 10.22075/IJNAA.2021.5311.
- [3] S. H. Asaad, I. S. Ahmed, and A. S. Mohammed, "Generalization Fuzzy Completely Intra G-ideals in G-Semi Groups," *J. Phys. Conf. Ser.*, vol. 1664, no. 1 (2020), doi: 10.1088/1742-6596/1664/1/012028.
- [4] N. Levine, "Semi-Open Sets and Semi-Continuity in Topological Spaces," *Am. Math. Mon.*, vol. 70, no. 1, pp. 36–41 (1963), doi: 10.1080/00029890.1963.11990039.
- [5] O. Njastad, "On some classes of nearly open sets," *Pacific J. Math.*, vol. 15, no. 3, pp. 961–970 (1965).
- [6] F. H. Mashhour, A. S., Allam, A. A., Mahmoud, F. S., & Khedr, "on Supra Topological Spaces," *Indian J. pure appl. Math.*, vol. 14, no. 4, pp. 502–510 (1983).
- [7] Z. T. Abdulqader, S. H. Khalil, L. S. Abdalbaqi, and M. H. Hadi, "Hpre-closed graph in topological spaces," *J. Interdiscip. Math.*, vol. 25, no. 6, pp. 1823–1828 (2022), doi: 10.1080/09720502.2022.2052572.
- [8] Z. T. Sameer and L. S. Abdalbaqi, "Some properties of an v-open set," *J. Interdiscip. Math.*, vol. 26, no. 5, pp. 873–879 (2023), doi: 10.47974/JIM-1519.

Received January, 2024