

On z^S -ideals of commutative rings

Ahmed Maatallah
Department of Mathematics
Faculty of Sciences
University of Monastir
5000 Monastir
Tunisia

Abstract

Let R be a commutative ring with nonzero unity and S a multiplicatively closed subset (m.c.s) of R . An S -disjoint ideal of R is called z^S -ideal if there exists $s = s_i \in S$ and whenever $M_s(a) \subseteq M_s(b)$ for some $a \in I$ and $b \in R$, then $sb \in I$, where $M_s(a) = \{Max_s(R) : sa \in M \text{ for some } s \in S\}$, for each $a \in R$. In the present paper, we introduce and study the concept of z^S -ideal which is an extension of the notion of the z -ideals. We generalize some well-known results on z -ideal to z^S -ideal. In addition, many examples, characterizations, and properties of z^S -ideal are given. We also investigate the relation between z -ideals and z^S -ideals of a commutative ring.

Subject Classification: (2010) 13A15, 13B30, 13B10.

Keywords: Commutative ring, z -ideals, z^S -ideals, S -prime, S -maximal ideals.

1. Introduction

Throughout this article, the term “ring” refers to a commutative ring with nonzero unity. Recently, in 2002, Anderson and Dumitrescu introduced the notion of S -noetherian ring which is considered as an extension of the Noetherian concept (see [3]). After that, various authors focused on this generalization and gave the S -version of some well-known properties of these rings. After drawing attention, many concepts in commutative algebra were extended to their S -version as the S -noetherian, for example, we mention the prime ideals, maximal ideals, 2-absorbing ideals, multiplication modules (see [7, 2, 13]). Note that many other concepts in commutative algebra have also been recently generalized and

E-mail: ahmedmaatallahsd93@gmail.com

well studied, as the concept of r -ideals, 2-absorbing semiprimary ideals, etc. (see [10, 12]). In the light of the above, we define and explore the S -version of the notion of z -ideals. A proper ideal I of a ring R is referred to as a z -ideal if for every $x \in I$ and $y \in R$ such that x and y are contained in the same set of maximal ideals, then also $y \in I$. This concept was introduced by "Kohls" in 1957 (see [8]) during the examination of the ring known as $C(X)$, the ring of real-valued continuous functions on a topological space X , and it has been expanded, by Mason, to an arbitrary ring in 1973 (see [9]). We mention that, for years, there have been many generalizations on this topic. For example, see [1, 6]. Note that in this article, our focus is on investigating the notion of the z^S -ideal of an arbitrary ring, but the case of the ring $C(X)$ can be a very interesting topic to be studied. Presently, we give some notations which will be utilized all through this article and can be found in [4]. For an ideal I of a ring R , we denote by $U(R)$, $Max(R)$, $Spec(R)$, and $Min(I)$ the set of all units in R , maximal ideals, prime ideals, and minimal ideals over I , respectively. For every ideal K of R , we define the set $(I:K) = \{r \in R : rK \subseteq I\}$. Clearly, $(I:K)$ is an ideal of R . If $K = xR$, we use $(I:x)$ instead of $(I:K)$. For a m.c.s S of a ring R , we denote by $I:S = \{x \in R : ux \in I \text{ for some } u \in S\}$. For an element x of R , we define M_x (resp., $\mathcal{M}(x)$) as the intersection (resp., set) of the maximal ideals that contain the element x . Recall from [7] that an S -disjoint ideal P is termed S -prime, if there exists $u \in S$ such that for every $x, y \in R$ if $xy \in I$, we have $ux \in I$ or $uy \in I$. Also, recall from [14] that an S -disjoint ideal M of R is called an S -maximal ideal if there exists $s \in S$ such that for each S -disjoint ideal N containing M we have $sN \subseteq M$. By $Max_s(R)$ (resp., $Spec_s(R)$) we mean the set of the S -maximal (resp., S -prime) ideals of R . Let x be an element of R , we define $\mathcal{M}_s(x) = \{M \in Max_s(R) : sx \in M \text{ for some } s \in S\}$. Now, we will give the principal definition of this article. An S -disjoint ideal of R is said to be z^S -ideal if there exists $s = s_i \in S$ such that if $\mathcal{M}_s(a) \subseteq \mathcal{M}_s(b)$, where $a \in I$ and $b \in R$, then $sb \in I$, (see Proposition 2.5 for an equivalent definition). Among other results of this article, Section 2 considers the fundamental properties of z^S -ideals. It is well known that the assertion " I is a z -ideal" does not imply, in general, that " IR_s is a z -ideal" even if $I \cap S = \emptyset$. also " IR_s is a z -ideal" does not also imply that " I is a z -ideal". So, a question that naturally arises here is, what condition on I to get IR_s is a z -ideal of R_s . For this reason, we prove in Theorem 2.17 that I is a z^S -ideal of $R \Leftrightarrow IR_s$ is a z -ideal of R_s , and there exists $t \in S$ such that $I:S = I:t$. In section 3, we introduce the concept of S -radical ideals. In fact, an S -disjoint ideal I is called S -radical if there exists $s \in S$ such that $\sqrt{I}:S = I:s$. We show that

the z^S -ideals are S -radicals, (Proposition 3.4). We turn of this paper, by the section 4, with a study of the direct and inverse image of a z^S -ideal by a ring homomorphism. In fact, we prove that if f is a surjective homomorphism from R to R' and S' be a m.c.s of R' , then the inverse image of a $z^{S'}$ -ideal of R' by f is a z^S -ideal of R , where $S = f^{-1}(S') = \{r \in R : f(r) \in S'\}$. (see Proposition 4.3). Moreover, for the direct image, we start by defining the S -strict homomorphism. A ring homomorphism $f: R \rightarrow R'$ is called S -strict if it is surjective and if $a \in R \setminus S^*$ then $f(a) \in R' \setminus f(S)$. Then, it was shown that for a z^S -ideal I and a S -strict homomorphism $f: R \rightarrow R'$ with $S \cap \ker(f) = \emptyset$, we have $f(I)$ is a $z^{f(S)}$ -ideal of R' (see Proposition 4.6). Throughout this paper, R means a ring and S a m.c.s of the ring R .

2. Basic properties of z^S -ideals

We start with the following definition.

Definition 2.1 ([14]): Let M be an S -disjoint ideal of R , then M is called an S -maximal ideal if there exists $u \in S$ such that for each S -disjoint ideal N containing M we have $uN \subseteq M$.

Example 2.2: Let $R = \mathbb{Z} \times \mathbb{Z}$ and $S = \mathbb{Z} \times \mathbb{Z}^*$. Clearly, S is a m.c.s and every S -disjoint ideal is S -maximal. Indeed, every S -disjoint ideal I of R is of the form $n\mathbb{Z} \times (0)$ for some integer n . Take $s = (n, 0) \in S$ and assume that there exists an S -disjoint ideal J of R such that $I \subseteq J$. We can write $J = m\mathbb{Z} \times (0)$ for some integer m . Then we can easily get $sJ \subseteq I$. For example, we can see that $4\mathbb{Z} \times (0)$ is an S -maximal ideal that is not maximal.

It was proved that if Q is S -prime, then there exists $s \in S$ such that $(Q:s') \subseteq (Q:s)$ for each $s' \in S$ (or, equivalently, $Q:S = Q:s$ for some $s \in S$), that is to say, $(Q:s)$ is the maximal element of the family $\{(Q:s')\}_{s' \in S}$, ([11], Lemma 2.16(i)). Since every S -maximal ideal is S -prime ([14], Proposition 10), the previous property is also true for the S -maximal ideal. As noted in [14], for $Q \in \text{Spec}_S(R)$ (resp., $M \in \text{Max}_S(R)$), we denote this $s \in S$ by s_Q (resp., s_M). Also, we will show that this $s \in S$ is not necessarily unique even for the S -maximal ideals (see Remark 2.10) and the s_Q or s_M will be signified one of the $s \in S$ such that $(Q:s)$ is the maximal element of $\{(Q:s')\}_{s' \in S}$.

Notation 2.3: Let x be an element of a ring R . We denote by $\mathcal{M}_S^*(x) = \{(M:s_M) : M \in \mathcal{M}_S(x)\}$ and $M_x^{*S} = \cap \mathcal{M}_S^*(x)$. Also, by $H_S(R)$ we mean the set of all prime ideals P maximal with the property $P \cap S = \emptyset$. Furthermore,

we denote by $H_s(x)$ the set of elements of $H_s(R)$ containing x , and $H_{s,x} = \cap H_s(x)$.

Lemma 2.4: For every $a \in R$, $H_s(a) = \mathcal{M}_s^*(a)$, and so $M_a^{*s} = H_{s,a}$.

Proof: First, note that $Max(R_s) = \{PR_s : P \in H_s(R)\}$. Let $P \in H_s(a)$, it is enough to show that $P \in \mathcal{M}_s(a)$. Let J be an S -disjoint ideal containing P , then $PR_s \subseteq JR_s$. Because $PR_s \in Max(R_s)$ and JR_s is proper, we have $PR_s = JR_s$, then $P:S = J:S$, it results that $J \subseteq P : S = P$. Conversely, let $P \in \mathcal{M}_s^*(a)$, then $P = (M:s_M)$ for some $M \in \mathcal{M}_s(a)$. Clearly, $a \in P$. Since $P \cap S = \emptyset$, $PR_s \subseteq QR_s$, for some $Q \in H_s(R)$, then $M \subseteq Q$. Because $M \in Max_s(R)$, we have $s_M Q \subseteq M$, thus $Q = P$, and $P \in H_s(a)$.

Proposition 2.5: The following are equivalent for an S -disjoint ideal I of R :

1. $\exists s = s_I \in S$ such that if $M_a^{*s} = M_b^{*s}$, where $a \in I$ and $b \in R$, then $sb \in I$.
2. $\exists s = s_I \in S$ such that if $\mathcal{M}_s^*(a) = \mathcal{M}_s^*(b)$, where $a \in I$ and $b \in R$, then $sb \in I$.
3. $\exists s = s_I \in S$ such that if $\mathcal{M}_s(a) = \mathcal{M}_s(b)$, where $a \in I$ and $b \in R$, then $sb \in I$.
4. $\exists s = s_I \in S$ such that if $\mathcal{M}_s(a) \subseteq \mathcal{M}_s(b)$, where $a \in I$ and $b \in R$, then $sb \in I$.
5. $\exists s = s_I \in S$ such that if $\mathcal{M}_s^*(a) \subseteq \mathcal{M}_s^*(b)$, where $a \in I$ and $b \in R$, then $sb \in I$.
6. $\exists s = s_I \in S$ such that $sM_a^{*s} \subseteq I$, for all $a \in I$.
7. $\exists s = s_I \in S$ such that $sH_{s,a} \subseteq I$, for all $a \in I$.

Proof: 1) $\Rightarrow$ 2). It is readily seen that $\mathcal{M}_s^*(a) = \mathcal{M}_s^*(b) \Rightarrow M_a^{*s} = M_b^{*s}$.

2) $\Rightarrow$ 3). It is clear.

3) $\Rightarrow$ 4). If $\mathcal{M}_s(a) \subseteq \mathcal{M}_s(b)$, where $a \in I$ and $b \in R$, then $\mathcal{M}_s(ab) = \mathcal{M}_s(b)$. Indeed, let $M \in \mathcal{M}_s(ab)$, then $s'ab \in M$ for some $s' \in S$. Used ([11], Lemma 2.16(i)), either $s_M a \in M$ or $s_M b \in M$. Therefore, $M \in \mathcal{M}_s(a) \cup \mathcal{M}_s(b) = \mathcal{M}_s(b)$. The converse is clear. Now, by hypothesis we have $sb \in I$, as desired.

4) $\Rightarrow$ 5). If $\mathcal{M}_s^*(a) \subseteq \mathcal{M}_s^*(b)$, then $\mathcal{M}_s(a) \subseteq \mathcal{M}_s(b)$. To see this, take $Q \in \mathcal{M}_s(a)$, then $(Q:s_Q) \in \mathcal{M}_s^*(a)$ and thus $(Q:s_Q) = (N:s_N)$, for some $N \in \mathcal{M}_s(b)$. Therefore, $s'b \in N$ for some. But $s_Q N \subseteq Q$, then $s_Q s'b \in Q$, and $Q \in \mathcal{M}_s(b)$.

5) $\Rightarrow$ 6). Let $a \in I$ and $x \in M_a^{*S}$. First, we prove that $\mathcal{M}_S^*(a) \subseteq \mathcal{M}_S^*(x)$. For, let $(Q:s_Q) \in \mathcal{M}_S^*(a)$, where $Q \in \mathcal{M}_S(a)$, then $x \in (Q:s_Q)$ and so $s_Q x \in Q$, thus $Q \in \mathcal{M}_S(x)$, this implies that $(Q:s_Q) \in \mathcal{M}_S^*(x)$. Now, using hypothesis, we conclude that $sx \in I$. Therefore, $sM_a^{*S} \subseteq I$, as desired.

6) $\Rightarrow$ 7) $\Rightarrow$ 1). Using the previous lemma.

Definition 2.6: An S -disjoint ideal I of a ring R is called z^S -ideal if it satisfies one of the equivalent conditions of Proposition 2.5.

Remark 2.7:

- (i) Clearly, the S -disjoint maximal ideals are S -maximal. Also, the S -maximal ideals are a z^S -ideals.
- (ii) If $S \subseteq U(R)$, then the set of z -ideals is equal to the set of z^S -ideal. Indeed, it is not difficult to prove that

$$\text{Max}_S(R) = \text{Max}(R) \text{ and so } \mathcal{M}_S(a) = \mathcal{M}(a), \text{ for each } a \in R.$$

Example 2.8: In Example 2.2, we prove that every S -disjoint ideal of R is a S -maximal ideal, and so a z^S -ideal. Then, according to [5], $4\mathbb{Z} \times (0)$ is a z^S -ideal distinct from being a z -ideal.

Example 2.9: Let D be a principal ideal domain (PID), $\mathbb{P}(D)$ a system of irreducible elements, $p \in \mathbb{P}(D)$, and $S = \{p^n : n \in \mathbb{N}\}$ a m.c.s of D . Then:

1. $I = aD$ is S -disjoint, where $a = p^\alpha q_1^{n_1} \cdots q_s^{n_s}$, if and only if $s \geq 1$.
2. $\text{Max}_S(D) = \{p^\alpha qD : \alpha \in \mathbb{N}, q \in \mathbb{P}(D), \text{ and } p \neq q\}$.
3. The z^S -ideal of D are of the form aD , where $a = p^\alpha q_1 \cdots q_s$, $s \geq 1$, $\alpha \in \mathbb{N}$, and the q_i are in $\mathbb{P}(D) \setminus \{p\}$.

Proof:

1. Obvious.
2. Let $q \in \mathbb{P}(D)$ such that $q \neq p$, and $M = p^\alpha qD$. By 1., we have $M \cap S = \emptyset$. Now, assume that $M \subseteq J$ for some S -disjoint ideal J . Based on a irreducibility argument we can easily show that $J = p^\beta qD$ and $\beta \leq \alpha$, therefore $p^\alpha J \subseteq M$. Also, one can easily show that the S -maximal ideals are of the previous form.
3. First of all, by 2., we can easily see that for $a = p^\alpha q_1^{n_1} \cdots q_s^{n_s} \in D$ we have $\mathcal{M}_S(a) = \{p^\beta q_i D : \beta \in \mathbb{N}, \text{ and } 1 \leq i \leq s\}$. Also, for every S -maximal

ideal $M = p^\alpha qD$, we can suppose that $s_M = p^\alpha$, and so $(M : s_M) = qD$. Then, $M_a^{*S} = q_1 \cdots q_s D$.

Now, for the “if” part, let $I = p^\alpha q_1 \cdots q_s D$, where the $q_i \in \mathbb{P}(D) \setminus \{p\}$, and $\alpha \in \mathbb{N}$. Clearly, $I \cap S = \emptyset$. Take a nonzero element $a \in I$, then $a = p^\alpha q_1 \cdots q_s d$ for some $d \in D$. by the above, $M_a^{*S} = q_1 \cdots q_s rD$ for some $r \in D$, hence, $p^\alpha M_a^{*S} = p^\alpha q_1 \cdots q_s rD \subseteq I$, and consequently, it proves I is a z^S -ideal of D .

For the “only if” part, let $I = dD$ be a nonzero z^S -ideal of D , where $d = p^\alpha q_1^{n_1} \cdots q_s^{n_s}$, the $q_i \in \mathbb{P}(D) \setminus \{p\}$, $\alpha \in \mathbb{N}$, the $n_i \in \mathbb{N}^*$, and $s \geq 1$, because $I \cap S = \emptyset$. We will show that each $n_i = 1$. Since $d \in D$, we have $s_I M_d^{*S} \subseteq I$, then $s_I q_1 \cdots q_s \in I$, and consequently, $n_i = 1$ for each $1 \leq i \leq s$.

Remark 2.10: Now, we will show that the s_I or the s_M , as in the definition of the z^S -ideal and the S -maximal ideal, are not unique in general. Like in the previous example, let $M = p^\alpha qD$ be an S -maximal ideal. It is clear that $M : p^\alpha = qD$, also if $\alpha' \geq \alpha$, we have $(M : p^{\alpha'}) = (M : p^\alpha)$. Now, one can see that for each integer j , we obtain $(M : p^j) \subseteq (M : p^\alpha) = qD$.

Now, we focus on the intersection of z^S -ideals.

Proposition 2.11: Then the following statements hold:

1. If S is finite, then the intersection of z^S -ideals is also a z^S -ideal.
2. If $I_1, \dots, I_n$ are z^S -ideals, then $I = \bigcap_{i=1}^n I_i$ is a z^S -ideal.

Proof: It is straightforward.

Definition 2.12 ([7]): A m.c.s S is called strongly-multiplicatively set if for every family $(s_\alpha)_{\alpha \in \Lambda}$ of elements of S we have

$$(\bigcap_{\alpha \in \Lambda} s_\alpha R) \cap S \neq \emptyset$$

Proposition 2.13: If S is a strongly-multiplicatively closed subset, then the intersection of z^S -ideals is a z^S -ideal.

Proof: Let $(I_\alpha)_{\alpha \in \Lambda}$ be a family of z^S -ideals of R and $I = \bigcap_{\alpha \in \Lambda} I_\alpha$. For each $\alpha \in \Lambda$, there exists $s_\alpha \in S$ such that whenever $a \in I_\alpha$ we have $s_\alpha M_a^{*S} \subseteq I_\alpha$. By hypothesis, we have $(\bigcap_{\alpha \in \Lambda} s_\alpha R) \cap S \neq \emptyset$. Pick an element t of this intersection, and let $a \in I$. For each $\alpha \in \Lambda$ we have $s_\alpha M_a^{*S} \subseteq I_\alpha$, then $tM_a^{*S} \subseteq I_\alpha$. Thus $tM_a^{*S} \subseteq I$. Therefore, I is a z^S -ideal of R .

Example 2.14: The assumption “ S is a strongly-multiplicatively closed subset” is not necessary for the previous proposition. Indeed, let $S = \{2^i : i \in \mathbb{N}\}$. Then S is a m.c.s of $\mathbb{Z}$ but not even anti-Archimedean. In fact, suppose that $(\bigcap_{i \in \mathbb{N}} 2^i \mathbb{Z}) \cap S \neq \emptyset$ and let $j \in \mathbb{N}$ such that $2^j \in \bigcap_{i \in \mathbb{N}} 2^i \mathbb{Z}$. Now, take $j_0 > j$, then by hypothesis, $2^{j_0} \in 2^j \mathbb{Z}$, which is a contradiction. Also, put $I_p = p\mathbb{Z}$, where p is a prime number other than 2. By the preceding example, we have I_p is a z^S -ideal. Furthermore, $\bigcap_{p \geq 3, \text{prime}} p\mathbb{Z} = (0)$, and (0) is a z^S -ideal of $\mathbb{Z}$, since $M_0^{*S} = \bigcap_{M \in \text{Max}_S(R)} (M : s_M) \subseteq \bigcap_{p \geq 3, \text{prime}} p\mathbb{Z} = (0)$.

It was shown that, if I_1 and I_2 are z -ideals, $I_1 I_2$ is a z -ideal $\Leftrightarrow I_1 I_2 = I_1 \cap I_2$ (see [9]). The following proposition explores a result similar to the one mentioned.

Proposition 2.15: For two z^S -ideals I and J , we have: IJ is a z^S -ideal of $R \Leftrightarrow s(I \cap J) \subseteq IJ$ for some $s \in S$.

Proof: $(\Rightarrow)$ By hypothesis, there exists $s \in S$ such that $sM_a^{*S} \subseteq IJ$, for all $a \in IJ$. Let $a \in I \cap J$, then $a^2 \in IJ$. This signifies that $sM_{a^2}^{*S} \subseteq IJ$. But $M_{a^2}^{*S} = M_a^{*S}$, hence $sa \in IJ$.

$(\Leftarrow)$ Since I (resp., J) is a z^S -ideal, let $s_i \in S$ (resp., $s_j \in S$) such that for each $a \in I$ (resp., $a \in J$) we have $s_i M_a^{*S} \subseteq I$ (resp., $s_j M_a^{*S} \subseteq J$), and let $a \in IJ$, then $s_i s_j M_a^{*S} \subseteq I \cap J$, then $u M_a^{*S} \subseteq IJ$ with $u = s s_i s_j$. Therefore, IJ is a z^S -ideal.

Remember that the saturation S^* of S is characterized by $S^* = \{x \in R : \frac{x}{1} \text{ is invertible in } S^{-1}R\}$. One can easily see that S^* is a m.c.s, and $x \in S^* \Leftrightarrow \exists y \in R$ such that $xy \in S$.

Proposition 2.16: I is a z^S -ideal of $R \Leftrightarrow I$ is a z^{S^*} -ideal.

Proof: It is clear, since $R_S = R_{S^*}$, and the two rings have the same prime spectrum.

Question 1: Let $S_1 \subseteq S_2$ be two m.c.s and I is a z^{S_1} -ideal of R such that $I \cap S_2 = \emptyset$, it is true that I is a z^{S_2} -ideal?

In general, the assertion “ I is a z -ideal” does not imply that “ IR_S is a z -ideal” even if $I \cap S = \emptyset$. For example, clearly (0) is a z -ideal of $\mathbb{Z}$. Take a nonzero prime ideal P , and $S = A \setminus P$, then the nonzero ideal PR_p represent the unique z -ideal of R_p . Also, “ IR_S is a z -ideal” does not also imply that “ I is a z -ideal”. For example, let (R, M) be a local ring, and assume that $P \subset M$ is a prime ideal. Take $S = R \setminus P$, then PR_p is the

unique z -ideal of R_p however P is not a z -ideal of R . (see also Example 2.19). But we have

Theorem 2.17: I is a z^S -ideal of $R \Leftrightarrow IR_S$ is a z -ideal of R_S , and $I:S = I:t$ for some $t \in S$.

Proof: For the “only if” part, by assumption, let $s = s_I \in S$ such that for every $a \in I$, $sM_a^{*S} \subseteq I$. Let $a \in I:S$ and let $s \in S$, such that $as \in I$. Then, $s_I a \in s_I M_a^{*S} = s_I M_{sa}^{*S} \subseteq I$, thus $a \in I:s_I$. Now, we demonstrate that IR_S is a z -ideal of R_S . For, let $x \in IR_S$, thus $x = \frac{a}{s}$ for some $a \in I$ and $s \in S$.

Clearly $\mathcal{M}_{R_S}(\frac{a}{s}) = \{PR_S : P \in H_S(a)\}$. let $y = \frac{b}{u} \in \bigcap_{P \in H_S(a)} PR_S$, then $s_I b \in s_I$
 $\bigcap_{P \in H_S(a)} P = s_I H_{S,a} = s_I M_a^{*S} \subseteq I$. Consequently, $y = \frac{s_I b}{s_I u} \in IR_S$.

For the “if” part, let $a \in I$. One can easily show that $\mathcal{M}_{R_S}(\frac{a}{1}) = \{PR_S : P \in H_S(a)\}$. Since IR_S is a z -ideal of R_S , $\bigcap_{P \in H_S(a)} PR_S \subseteq IR_S$, it results that,
 $(\bigcap_{P \in H_S(a)} P) R_S \subseteq \bigcap_{P \in H_S(a)} PR_S \subseteq IR_S$. Then $\bigcap_{P \in H_S(a)} P \subseteq I:S = I:s$, and so
 $sM_a^{*S} \subseteq I$.

Corollary 2.18:

1. If S consists of nonzerodivisors and there exists $s \in S$ such that $IR_S \cap R = (I:s)$. Then I is a z^S -ideal of $R \Leftrightarrow IR_S$ is a z -ideal of R_S .
2. If S is finite, then I is a z^S -ideal of $R \Leftrightarrow IR_S$ is a z -ideal of R_S .
3. If P is S -prime, then P is a z^S -ideal of $R \Leftrightarrow PR_S$ is a z -ideal of R_S .

Proof: According to the previous theorem, it suffices to prove that there exists $s \in S$ such that $I:S = I:s$, that is evident in all cases.

Example 2.19: Let $P \in \text{Spec}(R)$ and $S = R \setminus P$, then P is a z^S -ideal. Indeed, since $PR_p \in \text{Max}(R_S)$, then it is a z -ideal. Also, since P is an S -prime ideal, then P is a z^S -ideal. For example, let (R, M) be an integral local domain which is not a field and $S = R \setminus (0)$, then (0) is a z^S -ideal of R that is not a z -ideal.

Remark 2.20: Using the S -maximal ideals, we can give a new description of maximal ideal of the ring R_S . In fact, we can prove that $\text{Max}(R_S) = \{MR_S : M \in \text{Max}_S(R)\}$.

3. S-Radical and z^S -ideals

From [14], recall that we denote by

$$\sqrt[n]{I} = \{x \in R : ux^m \in I \text{ for some } u \in S \text{ and an integer } m\}.$$

It is clear that $\sqrt[n]{I}$ is an ideal of R containing I and $\sqrt{I}$. Note that, as mention in [14], we have $\sqrt[n]{I} = \sqrt{(IR_S)^c}$, where $(IR_S)^c$ is the inverse image of IR_S by the natural homomorphism $\pi : R \rightarrow R_S$.

Also, one can easily see that $\sqrt[n]{I} = \sqrt{I : S}$.

Definition 3.1: An S -disjoint ideal I of a ring R is called S -radical if there exists $u \in S$ such that $u\sqrt[n]{I} \subseteq I$, or, equivalently, $\sqrt{I : S} = I : u$.

Example 3.2: Let $R = \mathbb{Z}[X]$ and $S = \{2^n : n \in \mathbb{N}\}$.

- (i) Put $I = 4XR$. Clearly, I is a non radical S -disjoint ideal. However, I is an S -radical ideal of R . Indeed, let $f \in \sqrt[n]{I}$, then $2^n f^m \in I$ for some $n, m \in \mathbb{N}$, thus $2^n f^m = 4Xh$ for some $h \in R$. It follows that $2^n f(0)^m = 0$, thus $f(0) = 0$, and so $f = Xf_1$ for some $f_1 \in R$. Consequently, $4f \in I$, and $4\sqrt[n]{I} \subseteq I$.
- (ii) Put $I = X^3R$, we have $X \in \sqrt[n]{I}$, but for each $n \in \mathbb{N}$, $2^n X \notin I$. Hence, I is not S -radical.

Remark 3.3: For a finite m.c.s S , every S -disjoint radical ideal is S -radical. It is well-known that the z -ideals are necessarily radical. Now,

Proposition 3.4: Every z^S -ideal is S -radical.

Proof: If L is a z^S -ideal, then by Theorem 2.17, LR_S is a z -ideal of R_S , and $L : S = L : u$ for some $u \in S$. Since every z -ideal is radical, then $LR_S = \sqrt{LR_S} = \sqrt{L}R_S$. It follows that $L : S = \sqrt{L} : S$. Thus, $\sqrt{L : S} = \sqrt{\sqrt{L} : S} = \sqrt{L} : S = L : S = L : u$. Consequently, L is S -radical ideals.

The converse of the previous result is not in general true.

Example 3.5: Let $M, Q \in \text{Spec}(R)$ such that $Q \subset M$, and $S = R \setminus M$. Clearly, Q is S -radical. Also, note that QR_M is not a z -ideal of R_M , because MR_M is the unique z -ideal of R_M . Now, using Theorem 2.17, Q is not z^S -ideal.

Example 3.6:

- (i) Let $R = \mathbb{Z} \times \mathbb{Z}$ and $S = \mathbb{Z} \times \mathbb{Z}^*$. We showed that each S -disjoint ideal is a z^S -ideal, and so S -radical, by the previous proposition.

- (ii) In Example 3.2, we proved that I is not S -radical, hence it is not z^S -ideal.

Example 3.7: If D is a PID, and $S = \{p^n : n \in \mathbb{N}\}$, then the concept of z^S -ideals and that of S -radical ideals coincides. In fact, as shown in Example 2.9, an S -radical ideal should be of the form $I = aD$, where $a = p^\alpha q_1^{n_1} \cdots q_s^{n_s}$, the $q_i \in \mathbb{P}(D) \setminus \{p\}$, the $n_i \geq 1$, and $s \geq 1$. Now, assume that $n_j \geq 2$ for some $j \in \{1, \dots, s\}$, then clearly $q_1 \cdots q_s \in \sqrt{I} : S$, but for every $s \in S$, $sq_1 \cdots q_s \notin I$, which is a contradiction. Thus, the S -radical ideal of D are of the form aD , where $a = p^\alpha q_1 \cdots q_s$, $s \geq 1$, α an integer and the $q_i \in \mathbb{P}(D)$ not equals to p , as the z^S -ideals.

Proposition 3.8: Every S -disjoint ideal minimal over a z^S -ideal is also a z^S -ideal.

Proof: Since P is minimal over I , then $PR_S \in \text{Min}(IR_S)$. Furthermore, I is a z^S -ideal, using Theorem 2.17, IR_S is a z -ideal of R_S . The conclusion is drawn from ([9], Theorem 1.1) and Theorem 2.17.

Corollary 3.9: If (0) is a z^S -ideals, then every minimal prime S -disjoint ideal is so.

Corollary 3.10: Let J and P are two S -disjoint ideals where $P \in \text{Spec}(R)$ and $J \cap P$ is a z^S -ideal, then one of the two ideals is a z^S -ideal.

Proof: If P is not a z^S -ideal, pick $Q \in \text{Min}(J \cap P)$ such that $J \cap P \subseteq Q \subseteq P$. By the previous proposition, $J \subseteq Q \subset P$. Hence, $J = J \cap P$ is a z^S -ideal of R .

Now, we give some properties of the S -radical ideals concept. In fact, a relation between S -radical ideals and radical ideals is given.

Proposition 3.11: If I is S -radical, then,

1. $\exists t \in S$ such that $(I : t)$ is a radical ideal.
2. $\exists t \in S$ such that $I : S = I : t$.
3. IR_S is a radical ideal of R_S .

Proof: Clear.

Proposition 3.12:

1. If R is an integral domain (or S consists of nonzerodivisors) and $JR_S \cap R = (J : u)$. Then J is S -radical $\Leftrightarrow J_S$ is radical in R_S .
2. If S is finite, then J is S -radical $\Leftrightarrow J_S$ is radical in R_S .

Proof: The “only if” part of 1. and 2. is ensured by the previous proposition.

1. Assume that JR_S is radical. We will show that $\sqrt{J:S} = J:u$. Indeed, clearly, $\sqrt{J:S} \subseteq \sqrt{JR_S} \cap R = JR_S \cap R = J:u$, by hypothesis, and we are done.
2. It is obvious.

Proposition 3.13: I is S -radical in $R \Leftrightarrow I$ is S^* -radical.

Proof: By ([14], Proposition 5) and the fact that $V_S(I) = V_{S^*}(I)$.

Question 2: If $S_1 \subseteq S_2$ be two m.c.s and I is an S_1 -radical ideal with $I \cap S_2 = \emptyset$, it is true that I is an S_2 -radical ideal?

4. Direct and inverse image of z^S -ideals

For $s \in S$, we denote by $\bar{s} = \pi(s)$ where $\pi: R \rightarrow R/I$ is the canonical homomorphism. We denote by $\bar{S} = \{\bar{u} \mid u \in S\}$. Clearly, if I is S -disjoint, then $\bar{S}$ is a m.c.s of R/I . Now, we will focus on the study of the inverse image of a z^S -ideal by a ring homomorphism. It is well known that M is a maximal ideal of a ring $R \Leftrightarrow R/M$ is a field. In the following, we will give the S -version of this property. But before that, we will recall from [14], the definition of the S -version of the field, namely S -field. In matter of fact, R is called an S -field, if the zero ideal is S -maximal.

Proposition 4.1: If M is an S -disjoint ideal of R . Then

$$M \text{ is } S\text{-maximal ideal} \Leftrightarrow R/M \text{ is } \bar{S}\text{-field.}$$

Proof: “ $\Rightarrow$ ” We prove that $(\bar{0})$ is $\bar{S}$ -maximal. For that, let $\mathcal{J} = I/M$ be an $\bar{S}$ -disjoint ideal of R/M , then I is an S -disjoint ideal. Since $M \subseteq I$, then by hypothesis, $s_M I \subseteq M$. Therefore, $s_M(I/M) = (\bar{0})$, as desired.

“ $\Leftarrow$ ” Clear.

Lemma 4.2: Let $\varphi: R \rightarrow R'$ be a ring isomorphism and S' a m.c.s of R' . Then $S = \varphi^{-1}(S') = \{\varphi^{-1}(s') : s' \in S'\}$ is a m.c.s of R and

$$R \text{ is an } S\text{-field} \Leftrightarrow R' \text{ is an } S'\text{-field}$$

Proof: It is straightforward.

Proposition 4.3: If g is a surjective homomorphism from R to R' and S' is a m.c.s of R' . Then the inverse image of a $z^{S'}$ -ideal of R' by g is a z^S -ideal of R , where $S = g^{-1}(S') = \{r \in R : g(r) \in S'\}$.

Proof: Clearly, S is a m.c.s of R . First, we will prove that the inverse image of an S' -maximal ideal of R' by g is an S -maximal ideal of R . Consider $\pi: R' \rightarrow R'/I$ the canonical homomorphism, where I an ideal of R' , then $\pi \circ g$ is a surjective homomorphism with $g^{-1}(I)$ as kernel. Therefore, $R/g^{-1}(I) \simeq R'/I$ by the following isomorphism

$$\begin{aligned} \varphi: R/g^{-1}(I) &\rightarrow R'/I \\ \tilde{a} &\mapsto \overline{g(a)} \end{aligned}$$

If $M \in \text{Max}_{S'}(R')$, then according to the previous proposition, R'/M is a $\overline{S'}$ -field. Using the previous lemma, we have $R/g^{-1}(M)$ is a $\varphi^{-1}(\overline{S'})$ -field. But it is easy to see that $\varphi^{-1}(\overline{S'}) = \tilde{S}$, this mean that $R/g^{-1}(M)$ is a $\tilde{S}$ -field, that is, $g^{-1}(M) \in \text{Max}_{\tilde{S}}(R)$. Now, let J be a $z^{S'}$ -ideal of R' . Clearly, $g^{-1}(J) \cap S = \emptyset$. Let $a \in g^{-1}(J)$ and $b \in R$ such that $\mathcal{M}_{\tilde{S}}(a) \subseteq \mathcal{M}_{\tilde{S}}(b)$. We have $\mathcal{M}_{\tilde{S}}(g(a)) \subseteq \mathcal{M}_{\tilde{S}}(g(b))$. Indeed, let $M \in \mathcal{M}_{\tilde{S}}(g(a))$, Since g is surjective, there exists then $s' = g(s) \in S'$ such that $g(s)g(a) \in M$, hence $sa \in g^{-1}(M)$, and so $g^{-1}(M) \in \mathcal{M}_{\tilde{S}}(a)$. By assumption, $ub \in g^{-1}(M)$ for some $u \in S$, this implies that $g(ub) = g(u)g(b) \in M$, and so $M \in \mathcal{M}_{\tilde{S}}(g(b))$. Since $g(a) \in J$ and J is a $z^{S'}$ -ideal of R' , $s_1 g(b) \in J$, then $s_1 b \in g^{-1}(J)$, where $s_1 \in S$ and $g(s_1) = s_1$, then we are done.

Finally, we will pay attention to the study of the direct image of a z^S -ideal by an homomorphism. As in the previous proposition, we will examine first the S -maximal ideals. But even when $S = \{1\}$ or $S \subseteq U(R)$, it is well-known that the direct image of a maximal ideal is not in general maximal. For this reason, in the following, we will look for an additional condition to achieve our aim.

Let $g: R \rightarrow R'$ be a ring homomorphism and S a m.c.s of R . It is clear that if $S \cap \text{Ker}(f) = \emptyset$, then $f(S)$ is a m.c.s of R' . In the subsequent part, we mean by S a m.c.s of R disjoint with $\text{ker}(g)$.

Definition 4.4: A ring homomorphism $g: R \rightarrow R'$ is called S -strict if it is surjective and if $x \in R \setminus S^* \Rightarrow g(x) \in R' \setminus g(S)$.

Lemma 4.5: If $g: R \rightarrow R'$ is a surjective ring homomorphism. Then g is S -strict $\Leftrightarrow$ For each ideal I of R , if $I \cap S = \emptyset$, then $g(I) \cap g(S) = \emptyset$.

Proof: $\Rightarrow$ Assume that $g(I) \cap g(S) \neq \emptyset$, then there exists $x \in I$ such that $g(x) \in g(S)$. Hence $x \in S^*$, by hypothesis, it follows that $\exists y \in R$ such that $xy \in S$, so $xy \in S \cap I$, as desired.

$\Leftarrow$ Let $x \in R \setminus S^*$ and tack $I = xR$. Assume that $I \cap S \neq \emptyset$ and take $y \in I \cap S$, so $y = xr$ for some $r \in R$, it results that $x \in S^*$, absurd. By assumption, $g(I) \cap g(S) = \emptyset$, and so $g(x) \in R' \setminus g(S)$.

Proposition 4.6: If $g: R \rightarrow R'$ is a S -strict and I a z^S -ideal of R , then $f(I)$ is a $z^{f(S)}$ -ideal of R' .

Proof: Since $I \cap S = \emptyset$, then by the previous lemma, we have $g(I) \cap g(S) = \emptyset$. Now, let $a' \in g(I)$ and $b' \in R'$ such that $\mathcal{M}_{g(S)}(a') \subseteq \mathcal{M}_{g(S)}(b')$. Write $a' = g(a)$ and $b' = g(b)$ for some $a \in I$ and $b \in R$. We will show that $\mathcal{M}_S(a) \subseteq \mathcal{M}_S(b)$. For that, let $M \in \mathcal{M}_S(a)$. There exists $s \in S$ such that if Q is an S -disjoint ideal that containing M , then $sQ \subseteq M$. We have $M \cap S = \emptyset$, then by hypothesis, $g(M) \cap g(S) = \emptyset$. Furthermore, if J is $g(S)$ -disjoint and $g(M) \subseteq J$, then $M \subseteq g^{-1}(J)$. Also, it is easy to prove that $g^{-1}(J) \cap S = \emptyset$, hence $sg^{-1}(J) \subseteq M$, this implies that $g(s)J \subseteq g(M)$. It results that $g(M) \in \text{Max}_{g(S)}(R')$. Moreover, $ua \in M$ for some $u \in S$, then $g(u)g(a) \in g(M)$. According to the above, there exists $t \in S$ such that $g(t)g(b) \in g(M)$, thus $tb \in g^{-1}(g(M))$. But we have $M \subseteq g^{-1}(g(M))$ and $g^{-1}(g(M)) \cap S = \emptyset$, then $sg^{-1}(g(M)) \subseteq M$, thus $stb \in M$, this means that $M \in \mathcal{M}_S(b)$, as required. Since $a \in I$, we have $s_1b \in I$, and $g(s_1)g(b) \in g(I)$. Consequently, $g(I)$ is a $z^{g(S)}$ -ideal of R' .

Lemma 4.7: $\forall x \in R \setminus S^*, \exists N \in \text{Max}_S(R)$ such that $x \in N$.

Proof: Put $I = xR$, so I is S -disjoint. It is well-known that there exists an S -disjoint ideal $P \in \text{Spec}(R)$ such that $I \subseteq P$. Let

$$\mathcal{F} = \{P \in \text{Spec}(R) : I \subseteq P \text{ and } P \cap S = \emptyset\}.$$

$(\mathcal{F}, \subseteq)$ is inductive subset. Hence, $(\mathcal{F}, \subseteq)$ admits a maximal elements denoted by N , that is an S -maximal ideal. Indeed, $N \cap S = \emptyset$ and $I \subseteq N$. Assume that there exists an S -disjoint ideal J of R such that $N \subseteq J$, so there exists an S -disjoint prime ideal P_0 such that $J \subseteq P_0$, this implies that $I \subseteq P_0$, and so $P_0 \in \mathcal{F}$. Since $N \subseteq P_0$, then by the maximality of N in $(\mathcal{F}, \subseteq)$ we have $P_0 = N$, and $J = N$.

Proposition 4.8:

$$x \in \text{Jac}_S(R) \Leftrightarrow \forall s \in S, \forall y \in R, s + xy \in S^*,$$

where $Jac_S(R) = \bigcap_{N \in Max_S(R)} (N : s_N)$.

Proof: ($\Rightarrow$) If $x \in Jac_S(R)$, and suppose that there exists $y \in R$ and $s \in S$ such that $s + xy \notin S^*$. By the previous lemma, $s + xy \in M$ for some $M \in Max_S(R)$. However $x \in (M : s_M)$, this implies that $s_M x \in M$, hence $s_M s + s_M xy \in M$. It follows that $s_M s \in M \cap S$, a contradiction.

($\Leftarrow$) Assume that $x \notin Jac_S(R)$. There exists $N \in Max_S(R)$ such that $x \notin (N : s_N)$, this yields that $s_N x \notin N$, and so $x \notin N$, then $N \subset N + xR$ and $s_N(N + xR)$ not in N , then $S \cap (N + xR) \neq \emptyset$. Pick $s \in S \cap (N + xR)$, then $s = m + xa \in S$ for some $m \in N$ and $a \in R$, thus $m = s - xa \in N \cap S^*$, and so $N \cap S \neq \emptyset$, a contradiction.

Now, by the following remark, we give a class of examples of S -strict ring homomorphism.

Remark 4.9: If $\pi : R \rightarrow R/I$ is the canonical homomorphism, then

$$\pi \text{ is } S\text{-strict} \Leftrightarrow I \subseteq Jac_S(R).$$

References

- [1] A. R. Aliabad, M. Badie, and S. Nazari, "An extension of z -ideals and z° -ideals," *Hacet. J. Math. Stat.*, vol. 49, no. 1, pp. 254-272 (2020).
- [2] D. D. Anderson, T. Arabaci, Ü. Tekir, and S. Koç, "On S -multiplication modules," *Commun. Algebra*, vol. 48, no. 8, pp. 3398-3407 (2020).
- [3] D. D. Anderson and T. Dumitrescu, " S -Noetherian rings," *Comm. Algebra*, vol. 30, no. 9, pp. 4407-4416 (2002).
- [4] M. F. Atiyah and I. G. Macdonald, *Introduction to Commutative Algebra*, vol. 2, Addison-Wesley, Reading (1969).
- [5] A. Benhissi and A. Maatallah, "A question about higher order z -ideals in commutative rings," *Quaest. Math.*, vol. 43, pp. 1155-1157 (2020).
- [6] T. Dube and O. Ighedo, "Higher order z -ideal in commutative rings," *Miscolc Math. Notes*, vol. 17, pp. 171-185 (2016).
- [7] A. Hamed and A. Malek, " S -prime ideals of a commutative ring," *Beitrage Zur Algebra Und Geometrie/ Contributions to Algebra and Geometry*, vol. 61, no. 3, pp. 533-542 (2019).
- [8] C. W. Kohls, "Ideals in rings of continuous functions," *Fund. Math.*, vol. 45, pp. 28-50 (1957).

- [9] G. Mason, “z-ideals and prime ideals,” *J. Algebra*, vol. 26, pp. 280–297 (1973).
- [10] S. Onar, S. Yavuz, B. A. Ersoy, and K. Hila, “Intuitionistic fuzzy 2-absorbing semiprimary ideals of commutative rings,” *J. Discret. Math. Sci. Cryptogr.*, vol. 25, no. 8, pp. 2353–2368 (2022).
- [11] E. S. Sevim, T. Arabaci, Ü. Tekir, and S. Koç, “On S-prime submodules,” *Turk. J. Math.*, vol. 43, no. 2, pp. 1036–1046 (2019).
- [12] E. A. Ugurlu, “Generalizations of r-ideals of commutative rings,” *J. Interdiscip. Math.*, vol. 24, no. 8, pp. 2283–2293 (2021).
- [13] G. Ulucak, Ü. Tekir, and S. Koç, “On S-2-absorbing submodules and vn-regular modules,” *Analele Universitatii “Ovidius” Constanta*, vol. 28, no. 2, pp. 239–257 (2020).
- [14] E. Yıldız, B. A. Ersoy, Ü. Tekir, and S. Koç, “On S-Zariski topology,” *Comm. Algebra*, vol. 49, no. 8, pp. 3387–3397 (2020).

Received March, 2023