

Consequence for the resolution of the complex terms in the case (9,8,3)

Abeer Hussein Ali [†]

Niran Sabah Jasim ^{*}

Department of Mathematics

College of Education for Pure Science (Ibn Al-Haitham)

University of Baghdad

Baghdad 10071

Iraq

and

Ministry of Education

General Directorate of Education Baghdad

Karkh III

Baghdad 10011

Iraq

Abstract

The target of this treatise is to prove that the concatenation of the terms of characteristic zero (cha.zer.) resolution (reso.) is complex in the case (9,8,3) by ply the injective of maps, Capelli identities (Cap.ide.), planner and place polarization (pl.pol.).

Subject Classification (2020): 13D07.

Keywords: Complex terms, Resolution, Characteristic zero, Place polarization.

[†] ORCID-ID: 0009-0000-6998-4747

^{*} ORCID-ID: 000-0001-5340-3020

[†] E-mail: abeer.ali2103m@ihcoedu.uobaghdad.edu.iq

^{*} E-mail: niraan.s.j@ihcoedu.uobaghdad.edu.iq (Corresponding Author)

1. Introduction

Anywhere this treatise $\mathcal{R}$ commutative ring and has 1, $\mathcal{F}$ free $\mathcal{R}$ -module and $\mathcal{D}_i\mathcal{F}$ be the divided power (div.pow.) algebra of degree i , $\Lambda\mathcal{F}$ is the exterior algebra of finitely generated free $\mathcal{R}$ -module $\mathcal{F}$ and for the partition λ/μ , $\mathcal{K}_{\lambda/\mu}\mathcal{F} = \text{Im}(d'_{\lambda/\mu})$ where $d'_{\lambda/\mu}$ is the Weyl map $d'_{\lambda/\mu}: \mathcal{D}\mathcal{F} \rightarrow \Lambda\mathcal{F}$, [1]. The maps $\sum_{k>0} \mathcal{D}_{p+t_1+k} \otimes \mathcal{D}_{q-t_1-k} \otimes \mathcal{D}_r \rightarrow \mathcal{D}_p \otimes \mathcal{D}_q \otimes \mathcal{D}_r$ is k^{th} div.pow. of the pl.pol. from place 1 to place 2 ($\partial_{21}^{(k)}$), and the maps $\sum_{e>0} \mathcal{D}_p \otimes \mathcal{D}_{q+t_2+e} \otimes \mathcal{D}_{r-t_2-e} \rightarrow \mathcal{D}_p \otimes \mathcal{D}_q \otimes \mathcal{D}_r$ explicated as e^{th} div.pow. of the pl.pol. from place 2 to place 3 ($\partial_{32}^{(e)}$), and as in two-rowed case, where the resolution of the Weyl module with the three-rowed $(p, q, r; t_1, t_2)$, [2,3]. Anywhere this treatise we deficiency these Cap.ide.

$$\partial_{32}^{(2)} \partial_{21} = \partial_{21} \partial_{32}^{(2)} + \partial_{32} \partial_{31} \quad \dots(1)$$

$$\partial_{32} \partial_{21}^{(2)} = \partial_{21}^{(2)} \partial_{32} + \partial_{21} \partial_{31} \quad \dots(2)$$

$$\partial_{21}^{(2)} \partial_{32} = \partial_{32} \partial_{21}^{(2)} - \partial_{21} \partial_{31} \quad \dots(3)$$

$$\partial_{32} \partial_{31} = \partial_{31} \partial_{32} \quad \dots(4)$$

The cases (3,3,2), (8,7,3), (6,6,4;0,0) and (8,8)/(1,0) studied respectively in [4-7]. We stratify the same idea in [8] for the case (9,8,3). Also we can employ that for 2-visible submodules and modules, $\mathcal{PSL}(2, \mathfrak{U})$, where $\mathfrak{U} = 23, 29, 31, 37, [9-11] \mathcal{SL}(2, \mathfrak{U})$, $\mathfrak{U} = 23, 29, 31, 37, 125, 3125$, [12-14], $\mathcal{S.U.T.}(2, p)$, for a prime $p \leq 9$ and rad-quasi-prime submodules which are study in [15-16].

2. The treatise and the Results

The cha. ze. reso. in the case (9,8,3) has the formulation:-

$$\begin{aligned} & \mathcal{D}_{11}\mathcal{F} \otimes \mathcal{D}_7\mathcal{F} \otimes \mathcal{D}_2\mathcal{F} \mathcal{D}_{10}\mathcal{F} \otimes \mathcal{D}_7\mathcal{F} \otimes \mathcal{D}_3\mathcal{F} \\ & \mathcal{D}_{11}\mathcal{F} \otimes \mathcal{D}_8\mathcal{F} \otimes \mathcal{D}_1\mathcal{F} \rightarrow \oplus \rightarrow \oplus \rightarrow \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_8\mathcal{F} \otimes \mathcal{D}_3\mathcal{F} \\ & \mathcal{D}_{10}\mathcal{F} \otimes \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_1\mathcal{F} \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_2\mathcal{F} \end{aligned}$$

Behold the press planner:

$$\begin{array}{ccccc} \mathcal{D}_{11}\mathcal{F} \otimes \mathcal{D}_8\mathcal{F} \otimes \mathcal{D}_1\mathcal{F} & \xrightarrow{t_1} & \mathcal{D}_{11}\mathcal{F} \otimes \mathcal{D}_7\mathcal{F} \otimes \mathcal{D}_2\mathcal{F} & \xrightarrow{t_2} & \mathcal{D}_{10}\mathcal{F} \otimes \mathcal{D}_7\mathcal{F} \otimes \mathcal{D}_3\mathcal{F} \\ \mathfrak{f}_1 \downarrow & \mathcal{O} & \downarrow \mathfrak{f}_2 & \mathcal{J} & \downarrow \mathfrak{f}_3 \end{array}$$

$$\mathcal{D}_{10}\mathcal{F} \otimes \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_1\mathcal{F} \xrightarrow{u_1} \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_2\mathcal{F} \xrightarrow{u_2} \mathcal{D}_9\mathcal{F} \otimes \mathcal{D}_8\mathcal{F} \otimes \mathcal{D}_3\mathcal{F}$$

Define the maps $t_1(\mathfrak{k}) = \partial_{32}(\mathfrak{k})$, $\mathfrak{f}_1(\mathfrak{k}) = \partial_{21}(\mathfrak{k})$, and $\mathfrak{f}_2(\mathfrak{k}) = \partial_{21}^{(2)}(\mathfrak{k})$.

We requirement to acquaint the map u_1 and by (1) which manufacture the planner $\mathcal{O}$ commute. Thus $u_1 \circ \mathfrak{f}_1 = \mathfrak{f}_2 \circ d_1$

By the same tactic $\ell_1 \circ \ell_2 = 0$.

References

- [1] D. A. Buchsbaum, "A characteristic-free example of Lascoux resolution and letter place methods for intertwining numbers," *European Journal of Combinatorics*, vol. 25, pp. 1169-1179 (2004).
- [2] L. R. Vermani, *An Elementary Approach to Homological Algebra*, Chapman and Hall/CRC, Monographs and Surveys in Pure and Applied Mathematics (2003).
- [3] R. H. Haytham, "The reduction of resolution of Weyl module from characteristic-free resolution in case $(4,4,3)$," *Ibn Al-Haitham J. for Pure & Appl. Sci.*, vol. 25, no. 3, pp. 341-355 (2012).
- [4] R. H. Haytham and O. A. Alaa, "Complex of Lascoux in Partition $(3,3,2)$," *Ibn Al-Haitham J. for Pure & Appl. Sci.*, vol. 28, no. 1, pp. 171-178 (2015).
- [5] R. H. Haytham and S. J. Niran, "On free resolution of Weyl module and zero characteristic resolution in the case of partition $(8,7,3)$," *Baghdad Science Journal*, vol. 15, no. 4, pp. 455-465 (2018). [Online]. Available: <https://doi.org/10.21123/bsj.2018.15.4.0455>.
- [6] R. H. Haytham and S. J. Niran, "Weyl Module Resolution Res $(6,6,4;0,0)$ in the Case of Characteristic Zero," *Iraqi Journal of Science*, vol. 62, no. 4, pp. 1344-1348 (2021). [Online]. Available: <https://doi.org/10.24996/ij.s.2021.62.4.30>.
- [7] R. H. Haytham and S. J. Niran, "Exactness for complex sequence in the skew shape $(8,8)/(1,0)$," *Journal of Interdisciplinary Mathematics*, vol. 24, no. 6, pp. 1661-1668 (2021). [Online]. Available: <https://doi.org/10.1080/09720502.2021.1892271>.
- [8] S. J. Niran, S. J. Sawsan, and I. A. Ahmed, "Enforcement for the partition $(7,7,4;0,0)$," *Journal of Interdisciplinary Mathematics*, vol. 24, no. 6, pp. 1669-1676 (2021). [Online]. Available: <https://doi.org/10.1080/09720502.2021.1892272>.
- [9] M. S. Fiadh and W. H. Hanoon, "2-Visible Submodules and Fully 2-Visible Modules," *Iraqi Journal for Computer Science and Mathematics*, vol. 1, no. 2, pp. 24-28 (2020). [Online]. Available: <https://doi.org/10.52866/ijcsm.2020.01.01.002>.
- [10] N. A. S. Salem and N. S. Jasim, "Circularity segmentation for the groups $PSL(2,23)$ and $PSL(2,29)$," in 1st Samarra International Conference for Pure and Applied Sciences (SICPS2021), AIP Conf. Proc.,

- vol. 2394, 070032-1-070032-5 (2022). [Online]. Available: <https://doi.org/10.1063/5.0121782>.
- [11] N. S. Noor Alhuda and S. J. Niran, "Periodical split for the groups $PSL(2,31)$ and $PSL(2,37)$," *Journal of Discrete Mathematical Sciences and Cryptography*, vol. 25, no. 2, pp. 605-608 (2022). [Online]. Available: <https://doi.org/10.1080/09720529.2021.1982490>.
- [12] S. A. K. Sherouk and N. S. Jasim, "Calculation for the groups $SL(2,U)$, $U=31$ and 37 ," *Journal of Discrete Mathematical Sciences and Cryptography*, vol. 25, no. 2, pp. 609-613 (2022). [Online]. Available: <https://doi.org/10.1080/09720529.2021.1972614>.
- [13] S. A. Khalaf and N. S. Jasim, "Consequences for the groups $SL(2,U)$, $U=23$ and 29 ," in 1st Samarra International Conference for Pure and Applied Sciences (SICPS2021), AIP Conf. Proc., vol. 2394, 070031-1-070031-7 (2022). [Online]. Available: <https://doi.org/10.1063/5.0121781>.
- [14] M. S. Aldeen Zidan and N. S. Jasim, "Enforcement for the groups $SL(2,125)$ and $SL(2,3125)$," *Journal of Interdisciplinary Mathematics*, vol. 26, no. 7, pp. 1379-1384 (2023). [Online]. Available: <https://doi.org/10.47974/JIM-1554>.
- [15] H. Alameer, L. A. Fiadh, N. S. Jasim, and J. A. Eleiwy, "Applications for the groups $S.U.T.(2,p)$, where p prime greater than 9," *Iraqi Journal for Computer Science and Mathematics*, vol. 5, no. 1, pp. 78-84 (2024). [Online]. Available: <https://doi.org/10.52866/ijcsm.2024.05.01.005>.
- [16] R. N. M. Rana, G. A. Mahmood, S. F. Lemya and A. A. Hadi, "Rad-Quasi-Prime Submodules," *Iraqi Journal for Computer Science and Mathematics*, vol. 5, no. 2, pp. 21-25 (2024). [Online]. Available: <https://doi.org/10.52866/ijcsm.2024.05.02.002>.

Received May, 2024