

A generalization of the extending property for semimodules via weakly supplement subsemimodules

Shimaa M. Dawood *

Saad A. Al-Saadi †

Department of Mathematics

College of Science

Mustansiriyah University

Baghdad

Iraq

Abstract

This work introduces and investigates a generalization of extending semimodules via weak-supplement subsemimodules. An S -semimodule L is called weakly-supplement extending (shortly, wse -semimodule) if every subsemimodule of L is essential in a weakly-supplement subsemimodule of L . We asserted that the class of wse -semimodules properly contains both extending and lifting semimodules. Analogously to extending semimodules, this paper gives many characterizations of wse -semimodules. We showed that L is wse -semimodule if and only if each closed subsemimodule of L is a weakly supplement subsemimodule of L . Moreover, we discussed the relation of wse -semimodule and its relation with other known classes of semimodule, such as $\oplus$ -supplemented, lifting, and uniform semimodules. We inspected whether the wse -semimodule property is satisfied under direct sum, subsemimodule, epimorphic image, and inverse image.

Subject Classification (2020): 16Y60, 16Y80.

Keywords: Semiring, Extending semimodule, Weakly-supplement subsemimodule, Lifting semimodule.

1. Introduction

As algebraic structures, semirings and their applications in various fields have motivated many authors to explore new concepts in semirings and

* ORCID-ID: 0000-0003-4238-6565

† ORCID-ID: 0000-0002-6369-8173

* E-mail: Shamaa_84@uomustansiriya.edu.iq (Corresponding Author)

† E-mail: saalsaadi@uomustansiriya.edu.iq

semimodules. In his book, Golan [1] presented interdisciplinary applications of semirings. The concept of injective semimodules over semirings has been studied extensively (for details, see [2]). Following [3], extending semimodules were introduced as a generalization of injective semimodules. Recently, many papers have appeared providing new characterizations, generalizations, and related work on extending semimodules (cf. [4], [5], [6],[7]). The main focus of this paper is devoted to introducing and studying extending semimodules via weakly-supplement subsemimodules.

For the sake of wholeness, we give some definitions and results that are relevant to our work. Let L be a semimodule over a semiring S . A subsemimodule N of L is called subtractive if $n_1 + n_2 \in N$ and $n_1 \in N$ imply that $n_2 \in N$ [1]. A subsemimodule Y of L is said to be small if for any subsemimodule Y^* of L such that $Y + Y^* = L$ implies $Y^* = L$, denoted by $Y \ll L$ [8]. An epimorphism $f: A \rightarrow B$ (where A and B are S -semimodules) is called small if $\ker(f) \ll A$ [1]. An S - semimodule L is called the direct sum of subsemimodules T_1 , and T_2 of L , if each element $t \in T$ can be uniquely written as $t = t_1 + t_2$ where $t_1 \in T_1$ and $t_2 \in T_2$ [8] and denoted by $T = T_1 \oplus T_2$. Also, T_1 is said to be the summand of T if there is a subsemimodule T_2 of T such that $T = T_1 \oplus T_2$. A semimodule L is called lifting if for every subsemimodule $N \leq L$ of L there exist subsemimodules X and Y such that $L = X \oplus Y$ with $X \leq N$ and $N \cap Y \ll L$ [9]. Let N and K are subsemimodule of a semimodule L . Then N is called a supplement of K in L if $N + K = L$ and $N \cap K \ll L$. A subsemimodule Y is called a supplement subseminodule in L if it is a supplement of some subsemimodule of L . A semimodule L is said to be supplemented if every subsemimodule of L has a supplement [10]. Moreover, a subtractive subsemimodule N of a semimodule L is supplement if and only if it is minimal with the propriety $N + K = L$ [11]. Let X and Y be subsemimodules of a semimodule L such that $X + Y = L$, $X \cap Y \ll Y$ then we say that Y is a weak supplement of X in L [10]. A semimodule L is called weakly supplemented if every subsemimodule of N has a weak supplement in L [10]. It is clear from the definitions that each direct summand is a supplement subsemimodule and every supplement subsemimodule is weakly supplement. A non-zero subsemimodule J of L is called an essential subsemimodule of L , if $J \cap B \neq (0)$ for each nonzero subsemimodule B of L and denoted by $J \leq^e L$ [12]. A semimodule L is called extending if every subsemimodule of L is essential in a direct summand of L [3]. The left annihilator of x in S is defined by $\text{ann}(x) = \{s \in S \mid sx = 0\}$ and the set $Z(L) = \{x \in L \mid \text{ann}(x) \text{ is essential in } S\}$. It is easy to see that $Z(L)$ is a subsemimodule of L and we will call it the singular subsemimodule of L . If $Z(L) = L$ then L is called singular, and if $Z(L) = 0$, then L is called nonsingular [1]. A semimodule L is said to be uniform if any subsemimodule of L is essential [7]. A semimodule L is referred to as strongly extending if closed subsemimodules of L are invariant summands [13]. A semimodule L is said to be semisimple if it is a direct sum of its simple subsemimodules [14]. Let K be a subsemimodule of a semimodule L , then we say L/K is a quotient semimodule of L by K , defined by $L/K = \{[x] \mid x \in L\}$ [10]. A subsemimodule N of L is said

to be fully invariant if $f(N) \subseteq N$ for each endomorphism f on L [14]. A semimodule is said to be duo if each subsemimodule of L is fully invariant [14],[18]. A semimodule C is said to be injective if, for any S -semimodules D and K , any S -monomorphism $\alpha: D \rightarrow K$, and any S -homomorphism $\phi: D \rightarrow C$, there is an S -homomorphism $\beta: K \rightarrow C$ in which $\alpha = \beta \circ \phi$ [15]. If E is an injective S -semimodule and an essential extension of a semimodule C , then E is said to be an injective hull of C , denoted by $E(C)$ [3]. The Jacobson radical of a semimodule L , denoted by $J(L)$, is the intersection of all maximal subsemimodules of L [16]. A semimodule L is referred to as stable-extending whenever a stable subsemimodule of L is contained essentially in a summand of L [16]. A semimodule L is called distributive if, for all subsemimodules A, B , and C of L , the following equality holds $\mathcal{A} \cap (B + C) = (\mathcal{A} \cap B) + (\mathcal{A} \cap C)$ [17],[19].

2. Weakly Supplement Extending Semimodules

Firstly, we introduce the following concept:

Definition 2.1: An S -semimodule L is called *weakly-supplement extending* (abbreviated as *wse* –semimodule) if every subsemimodule of L is essential in a weakly-supplement subsemimodule of L .

The next result provides as a characterization of *wse* –semimodule.

Proposition 2.2: An S -semimodule L is a *wse* –semimodule if and only if every closed subsemimodule of L is weakly-supplement subsemimodule in L .

Proof: Let A be a closed subsemimodule of L . Since L is a *wse* –semimodule, then $A \leq^e T$, where T is a weakly-supplement subsemimodule of L . However, A is a closed subsemimodule of L (that is, A has no proper essential extension in L), hence $A = T$. So A is a weakly-supplement subsemimodule in L . Conversely, let A be a subsemimodule of L and, by [5], A is essential in a closed subsemimodule T of L . By the assumption, T is a weakly-supplement subsemimodule in L , and so L is a *wse* –semimodule.

Remarks and Examples (2.3):

1. Every weakly supplemented semimodule, and hence every supplemented semimodule, is a *wse* –semimodule. The reverse direction is not always true. The $\mathbb{Z}$ as $\mathbb{Z}^+$ -semimodule is not weakly-supplemented (and hence is not supplemented). For example, the subsemimodule $2\mathbb{Z}$ of $\mathbb{Z}_{\mathbb{Z}^+}$ has no weakly-supplement in $\mathbb{Z}_{\mathbb{Z}^+}$. But $\mathbb{Z}$ as $\mathbb{Z}^+$ -semimodule is *wse* –semimodule.
2. Every extending semimodule is a *wse*-semimodule, while the opposite is not generally true. Actually, $\mathbb{Z}_8 \oplus \mathbb{Z}_2$ as an $\mathbb{Z}^+$ -semimodules is a *wse* –semimodule (since it is weakly supplemented) which is not an extending semimodule [5].

3. Every semi-simple semimodule is a *wse* –semimodule. The converse is not true generally. For example, $\mathbb{N}$ as $\mathbb{N}$ -semimodule is *wse* –semimodule but is not semi-simple.
4. Every uniform semimodule is a *wse* –semimodule, while the opposite is not always true. For example, $\mathbb{Z}_{10}$ as $\mathbb{Z}^+$ semimodule is *wse* –semimodule, but it is not uniform.
5. Extending semimodules and lifting semimodules are incomparable; however, every lifting semimodule is a *wse* –semimodule. Moreover, $\mathbb{Z}$ as $\mathbb{Z}^+$ -semimodule is *wse* –semimodule, and it is not lifting.
6. Every $\oplus$ -supplemented semimodule is a *wse* –semimodule, while the converse is not always true. For example, $\mathbb{Z}$ as $\mathbb{Z}^+$ -semimodule is *wse* –semimodule and is not $\oplus$ -supplemented.

The next result provides a sufficient condition for *wse* –semimodules.

Proposition 2.4: An S -semimodule L is a *wse* –semimodule if and only if every subtractive subsemimodule of L is essential in a weakly-supplement subsemimodule of L .

Proof: The necessary part follows directly from the definition of *wse* –semimodule. Conversely, let A be a subsemimodule of L . By [5], there is a closed subtractive subsemimodule K of L such that $A \leq^e K$. By the assumption, there exists a weakly-supplement N of L such that $K \leq^e N$. Hence, $A \leq^e N$. Therefore, L is *wse* –semimodule. $\square$

Proposition 2.5: Let L be a subtractive S -semimodule which has an injective envelope. Then L is *wse* –semimodule if and only if $D \cap L$ is weakly-supplement subsemimodule in L for each summand D of $E(L)$.

Proof: Let D be a summand of the injective hull $E(L)$ of L . Put $E(L) = D + N$, where N is subsemimodule of $E(L)$. Suppose that $D \cap L \leq^e T$, where T is a subsemimodule of L . Let $t \in T$, then $t = x + y$ for some $x \in D$ and $y \in N$. Since D is a summand of the injective hull $E(L)$, by [18, corollary 3.2], D is subtractive. Now, if $t = x + y \in D$ and $x \in D$ then $y \in D$, and this contradicts $D \cap N = 0$. Thus $t \notin D$ and so, $y \neq 0$. Since L is essential in $E(L)$ (by [19, Lemma 2]) then there is $s \in S$ such that $0 \neq sy \in C$. Now $st = sx + sy \in D \cap L \subseteq T$. Again since D is a subtractive and $sx, sx + sy \in D$, then $sy \in D$ and $sy \in T \cap N$. But $N \cap L$ is essential in T , so $(0) = (D \cap L) \cap N \leq^e T \cap N = 0$. Then $sy = 0$, which is a contradiction. Thus $D \cap L$ is a closed subsemimodule of L . By Proposition (2.2), $D \cap L$ is a weakly-supplement of L . Conversely, let N be a subsemimodule of L and Let C be a complement of N in L . By [20, Definition 2.1], C is a subtractive subsemimodule of L . Also by [20, proposition 2.4], then $N \oplus C$ is an essential subsemimodule of L . and by [21, lemma 2.2.3], $E(L) = E(N) \oplus E(C) = E(N \oplus C) = E(L)$. Then, $E(C)$ is a summand of $E(L)$,

and by hypothesis, $E(C) \cap L$ is a weakly-supplement of L . But N is essential in $E(N)$ and L is essential in L , then by [12, Proposition 3.3] $N = N \cap L$ is essential in the weakly-supplement $E(N) \cap L$ in L . Then L is wse -semimodule.

Lemma 2.6: Let $\alpha: X \rightarrow Y$ be an epimorphism image of semimodules X and Y . If N is a closed subsemimodule of Y , then $\alpha^{-1}(N)$ is a closed subsemimodule of X .

Proof: Let $\alpha: X \rightarrow Y$ be an epimorphism, and N be a closed subsemimodule of Y . By the first fundamental theorem of homomorphism for semimodules [22, Theorem 4.2], $X/\ker(\alpha) \cong Y$. Let $K = \alpha^{-1}(N)$, which is a subsemimodule of X , then $N \cong K/\ker(\alpha)$. Suppose that $\ker(\alpha) \leq K \leq_e T \leq X$, then $T/K \cong T/\ker(\alpha)/K/\ker(\alpha)$. By [6, proposition 4.2] T/K is a singular semimodule, and since Y is non-singular, then $K/\ker(\alpha) \leq_e T/\ker(\alpha)$. Thus, $K/\ker(\alpha) = T/\ker(\alpha)$ by [12, proposition 4.2], hence $K = T$, which means that $K = \alpha^{-1}(N)$ is a closed semimodule of X . $\square$

Proposition 2.7: Let X be a wse -semimodule, then every non-singular image of X is a wse -semimodule.

Proof: Let $\alpha: X \rightarrow Y$ be an epimorphism of semimodules with X wse -semimodule and Y a non-singular image of X . Let N be a closed subsemimodule of Y , then by Lemma 2.6, $\alpha^{-1}(N)$ is a closed subsemimodule of X . By proposition (2.2), since X is a wse -semimodule, then $\alpha^{-1}(N)$ is a weakly supplement subsemimodule of X , so there is a subsemimodule H of X such that $X = H + \alpha^{-1}(N)$ and $H \cap \alpha^{-1}(N) \ll X$. then we have $Y = \alpha(H + \alpha^{-1}(N)) = \alpha(H) + \alpha(\alpha^{-1}(N)) = \alpha(H) + N$ and by [15] and since α is epimorphism then $\ker(\alpha) \leq \alpha^{-1}(N)$ we have by [16] $\alpha(H \cap \alpha^{-1}(N)) = \alpha(H) \cap \alpha(\alpha^{-1}(N)) = \alpha(H) \cap N \ll Y$ since $(H \cap \alpha^{-1}(N)) \ll X$. Then N is a weakly-supplement subsemimodule of Y , and so Y is a wse -semimodule of L .

The next result asserts that a wse -semimodule is closed under the isomorphic property.

Proposition 2.8: Let Y_1 and Y_2 are S -semimodule. If Y_1 is a wse -semimodule and $Y_1 \cong Y_2$, then Y_2 is a wse -semimodule.

Proof: Let $\alpha: Y_1 \rightarrow Y_2$ be an isomorphism and W be a subsemimodule of Y_2 . Then $\alpha^{-1}(W)$ is a subsemimodule of Y_1 . Since Y_1 is wse -semimodule of L . Then $\alpha^{-1}(W)$ is essential in a weakly-supplement subsemimodule K of Y_1 . But $\alpha(\alpha^{-1}(W)) = W$ is an essential in $\alpha(K)$. Since K is weakly-supplement subsemimodule in Y_1 then there is H is a subsemimodule of Y_1 such that $Y_1 = K + H$, and $K \cap H \ll Y_1$. Then by [16] $\alpha(Y_1) = \alpha(K) + \alpha(H)$. And so $Y_2 = \alpha(K) + \alpha(H)$, since $K \cap H \ll Y_1$ and $\alpha: Y_1 \rightarrow Y_2$ homomorphism, then $\alpha(K \cap H) \ll Y_2$ [16], since α is isomorphism then α is monomorphism and

$\ker \alpha = 0$, and we have $\ker \alpha \subseteq H$, then by [16], $\alpha(K \cap H) = \alpha(K) \cap \alpha(H) \ll Y_2$, then $\alpha(K)$ is weakly-supplement subsemimodule of Y_2 , hence Y_2 is a $\mathcal{Wse}$ -semimodule.

Proposition 2.9: Let $\alpha: L \rightarrow K$ be a small epimorphism and K be a $\mathcal{Wse}$ -semimodule, such that for every nonzero closed subsemimodule N of L with $\ker \alpha \subseteq N$. Then L is $\mathcal{Wse}$ -semimodule.

Proof: Let K be a $\mathcal{Wse}$ -semimodule, and $\gamma: L \rightarrow K$ be a small epimorphism, and N be a closed subsemimodule of L , since $\ker \gamma \leq N$ by [16] and γ is an epimorphism. Then by the first fundamental theorem of homomorphism for semimodules [7, Theorem 2.4], $\gamma(N) \cong N / \ker \gamma$ is closed in $L / \ker \gamma \cong K$, then $L / \ker \gamma$ is a $\mathcal{Wse}$ -semimodule, so $\gamma(N)$ is a weakly-supplement subsemimodule of K , then N is a weakly-supplement subsemimodule of L . Then L is $\mathcal{Wse}$ -semimodule. $\square$

Proposition 2.10: Let N be a weakly-supplement subsemimodule of $\mathcal{Wse}$ -semimodule L such that the intersection of any two weakly-supplement subsemimodule in L is weakly-supplement in N . Then N is $\mathcal{Wse}$ -semimodule.

Proof: Let N be a weakly-supplement subsemimodule in L and let F be a subsemimodule in N . Since L is a $\mathcal{Wse}$ -semimodule, F is essential in a weakly-supplement subsemimodule H of L . By hypothesis, $N \cap H$ is a weakly-supplement subsemimodule in N since F is essential in H and N is essential in N . Then we have $F \cap N$ is essential in $H \cap N$ and $F \cap N = F$ is essential in $H \cap N$. Then N is $\mathcal{Wse}$ -semimodule. $\square$

Proposition 2.11: Every direct summand of a $\mathcal{Wse}$ -semimodule is a $\mathcal{Wse}$ -semimodule.

Proof: Let K be a direct summand of $\mathcal{Wse}$ -semimodule L , and H be a closed subsemimodule of K . By [5], K is a closed subsemimodule of L . Since L is a $\mathcal{Wse}$ -semimodule and H is a closed subsemimodule of L , H is a weakly-supplement subsemimodule of L . Let N be a subsemimodule of L such that $L = H + N$ and $H \cap N \ll L$. Since every closed subsemimodule is subtractive [5], then by using the modular law for semimodules [20], then $K = K \cap L = K \cap (H + N) = H + (K \cap N)$ and $H \cap (K \cap N) = (H \cap K) \cap N$. Since $(H \cap N) \cap K \subseteq H \cap N$ and $H \cap N \ll L$, so, $H \cap (K \cap N) \ll L$. Since K is a direct summand of L , then by [22], $H \cap (K \cap N) \ll K$ and so H is a weakly-supplement subsemimodule of K . Therefore, K is a $\mathcal{Wse}$ -semimodule. $\square$

Proposition 2.12: If K_1 and K_2 are semimodules such that $K = K_1 \oplus K_2$ is a distributive semimodule. Then $K_1 \oplus K_2$ is a $\mathcal{Wse}$ -semimodule iff K_1 and K_2 are $\mathcal{Wse}$ -semimodules.

Proof: The necessary part is directly by proposition (2.11). Conversely, let D be a closed subsemimodule of $K_1 \oplus K_2$. Since $K_1 \oplus K_2$ is distributive semimodule, then $D = D \cap (K_1 \oplus K_2) = (D \cap K_1) \oplus (D \cap K_2)$. So, $D \cap K_1$ and $D \cap K_2$ are closed subsemimodules of K_1 and K_2 respectively. But K_1 and K_2 are $\mathcal{Wse}$ -semimodule. Thus there is subsemimodules N_1 of K_1 and N_2 of K_2 so $K_1 \oplus K_2 = (N_1 + (D \cap K_1)) \oplus (N_2 + (D \cap K_2))$ such that $(N_1 + N_2) \oplus ((D \cap K_1) \oplus (D \cap K_2)) = (N_1 + N_2) + D$, since $(N_1 \cap D) \ll K_1$ and $(N_2 \cap D) \ll K_2$ are $(N_1 \cap D) + (N_2 \cap D) \ll K_1 \oplus K_2$, Hence D is weakly-supplement subsemimodules of $K_1 \oplus K_2$, therefore $K_1 \oplus K_2$ is $\mathcal{Wse}$ -semimodule. $\square$

Proposition 2.13: Let $L = \bigoplus_{i \in I} K_i$ be a semimodule such that every closed subsemimodule in L is fully invariant. Then L is $\mathcal{Wse}$ -semimodule iff each K_i for each $i \in I$ is $\mathcal{Wse}$ -semimodule.

Proof: The necessary part is directly by proposition (2.11). Conversely, let H be a closed subsemimodule of L and since L is fully invariant then there is a mapping $\alpha: Y \rightarrow Y_i$ define by $\alpha_i(x) = y_i$, where $y_i \in Y_i$ and $x = \sum_{i \in I} y_i$ such that $\alpha_i(H) \subseteq H$ and $\alpha_i(H) \subseteq Y_i$ hence $L = \bigoplus_{i \in I} (H \cap K_i)$. Since $H \cap K_i$ is direct summand then by [5, proposition 2.1.14] $H \cap K_i$ is closed subsemimodules of L . But $H \cap K_i \subseteq K_i \subseteq L$ then $H \cap K_i$ is closed in K_i . Therefore $H \cap K_i$ is a weakly-supplement subsemimodule of K_i . Thus $L = \bigoplus_{i \in I} (H \cap K_i)$ is $\mathcal{Wse}$ -semimodule, Then L is $\mathcal{Wse}$ -semimodule. $\square$

Proposition 2.14: Every weakly-supplement subsemimodule is a direct summand. Then L is $\mathcal{Wse}$ -semimodule if and only if L is extending. $\square$

Lemma 2.15: [5] *A subsemimodule of a semimodule is a complement if and only if it is closed.*

Lemma 2.16: *Let L be an S - semimodule and $N \subseteq L$, then N is closed if and only if $[N:L]$ is a closed ideal in S*

Proof: Let N be a closed subsemimodule of a semimodule L , then there exists a subsemimodule H of L such that N is a relative complement of N if $H \cap N = 0$, where H is the maximal subsemimodule satisfies the condition and we get $[N:L]$ is relative complement of $[H:L]$ in S , Also we suppose that $[N:L] \leq^e A$. Where A is ideal in S and since $[H:L] \cap A \subseteq A$ and $(0) = (A \cap [H:L] \cap [N:L])$ hence $(0) = A \cap [H:L]$. But $[N:L]$ is ideal in S satisfies the relation $[N:L] \cap [H:L] = 0$, hence $[N:L] \subseteq A$, therefore $[N:L] = A$. conversely suppose that $H \subseteq L$ such that $N \leq^e H$ and so $[N:L] \leq^e [H:L]$ and A is ideal in S such that $A \subseteq [H:L]$ and $[N:L] \cap A = 0$, So $N \cap AL = 0$, But $AL \subseteq H$ and $N \leq^e H$ hence $AL = 0$ when $A = 0$, therefore $[N:L] \leq^e [H:L]$ but $[N:L]$ is closed ideal in S , we get $[N:L] = [H:L]$ hence $N = K$ and therefore N be a closed subsemimodule of a semimodule of L . $\square$

Lemma 2.17: Let L be an S -semimodule and $N \subseteq L$ if $N \leq^e L$ then $[N:(x)] \leq^e S, \forall x \in L$.

Proof: Let $x \in L$ and $A \neq 0$ is an ideal in S if $AS_x = 0$ then $\text{ann}(x) \subseteq A$. But, $\text{ann}(x) \subseteq [N:(x)]$. So $[N:(x)] \subseteq A$ then $A \cap [N:(x)] \neq 0$, and if $AS_x \neq 0$ then $N \cap AS_x \neq 0$. Hence $a \in A$ such that $Ax \neq 0$ then $a \in [N:(x)]$, so $A \cap [N:(x)] \neq 0$ and hence $[N:(x)] \leq^e S, \forall x \in L$. $\square$

Proposition 2.18: Every finitely generated faithful multiplication S -semimodule over a commutative $\mathcal{Wse}$ -semiring S is a $\mathcal{Wse}$ -semimodule.

Proof: Let N and L be subsemimodules of S , we set $[N:L] = \{s \in S: sL \subseteq N\}$. It is clear that $[N:L]$ is an ideal of S [21]. Let N be a closed subsemimodule of a semimodule L , Since N is a closed subsemimodule in L , $[H:L]$ is a closed ideal in S by Proposition 2.16. Thus, by hypothesis, $[N:L]$ is weakly-supplement ideal in S . Then there exists an ideal I of S such that $[N:L] + I = S$ and $[N:L] \cap I \ll S$. Now by proposition 2.16, we have $L = SL = ([N:L] + I)L = [N:L]L + IL$. Let $([N:L]L \cap IL) + KL = L$, since L is a multiplication semimodule [23] so $[(N:L) \cap I] + K = L$. But L is a finitely generated faithful multiplication S -semimodule. And $L = SL = ([N:L] \cap I) + K$, then $S = ([N:L] \cap I) + K$ and since $[N:L] \cap I \ll S$, then we have $S = K$ then $L = KL$ is $\mathcal{Wse}$ -semimodule.

Acknowledgments

The authors would like to thank Mustansiriyah University (<http://www.uomustansiriyah.edu.iq>), Baghdad, Iraq, for its moral support in the present work.

References

- [1] J. S. Golan, Theory of Semirings with Applications in Mathematics and Theoretical Computer Science, Pitman Monogr. Surveys Pure Appl. Math., Longman Scientific and Technical, Harlow (1992).
- [2] K. Głazek, A Guide to the Literature on Semirings and Their Applications in Mathematics and Information Sciences: With Complete Bibliography, Dordrecht: Kluwer Academic Publishers (2002).
- [3] S. A. Alhashemi, "Extending Semimodules and Some of Their Generalizations," (Ph.D. thesis), University of Babylon, Babil, Iraq (2022).
- [4] A. H. A. Khaleq and A. M. A. Alhossaini, "Distributive semimodules: Theory, properties, and extensions," *Journal of Kufa for Mathematics and Computer*, vol. 10, no. 2, pp. 90–96 (2023).

- [5] A. J. Naeemah and S. A. Al-Saadi, "Strongly extending semimodules over a semiring," *Journal of Discrete Mathematical Sciences and Cryptography*, vol. 27, no. 5, pp. 1677–1687 (2024).
- [6] K. S. H. Aljebory and A. M. A. Alhossaini, "Almost injective semimodules," *Iraqi Journal of Science*, vol. 64, no. 9, pp. 4634–4643 (2023).
- [7] A. Faris and S. A. Al-Saadi, "Fully stable semirings and related concepts," *Journal of Discrete Mathematical Sciences and Cryptography*, vol. 28, no. 4-A, pp. 1025–1030 (2025).
- [8] N. X. H. Tuyen and H. X. Thang, "On superfluous subsemimodules," *Georgian Mathematical Journal*, vol. 10, no. 4, pp. 763–770 (2003).
- [9] M. Sall, L. Fall, and D. Sow, "On S-lifting semimodules over semirings," in *Scientific Days of Doctoral School of Mathematics and Computer Sciences International Conference*, Cham, Switzerland: Springer Nature, pp. 147–160 (2023).
- [10] M. M. T. Altaee and A. M. A. Alhossaini, "Supplemented and π -projective semimodules," *Iraqi Journal of Science*, vol. 61, no. 6, pp. 1479–1487 (2020).
- [11] M. Sall, M. Barry, and L. Fall, "On small subtractive and projective semimodules over zero-sum semiring," *JP Journal of Algebra, Number Theory and Applications*, vol. 61, no. 2, pp. 117–134 (2023).
- [12] K. Pawar, "A note on essential subsemimodules," *New Trends in Mathematical Sciences*, vol. 1, no. 2, pp. 18–21 (2013).
- [13] A. J. Naeemah and S. A. Al-Saadi, "The class of semimodules with direct summands are stable," *Journal of Discrete Mathematical Sciences and Cryptography*, vol. 28, no. 4-B, pp. 1281–1289 (2025).
- [14] A. M. A. Alhossaini and Z. A. H. Aljebory, "On p-duo semimodules," *Journal of University of Babylon, Pure and Applied Sciences*, vol. 26, no. 4 (2018).
- [15] S. N. Il'in, "On injective envelopes of semimodules over semirings," *J. Algebra Appl.*, vol. 15, no. 7, Art. no. 1650122 (2016).
- [16] S. H. Alsaebari and A. M. Alhossaini, "Nearly injective semimodules," *Journal of University of Babylon for Pure and Applied Sciences*, vol. 27, no. 1, pp. 11–31 (2019).
- [17] A. J. Naeemah and S. A. Al-Saadi, "Stable-extending semimodules over semirings," in *AIP Conference Proceedings*, vol. 3264, no. 1, p. 050110, AIP Publishing LLC (Mar. 2025).

- [18] J. N. Chaudhari and D. R. Bonde, "On direct sum of partitioning sub-semimodules of semimodules over semirings," *Journal of Advanced Research in Pure Mathematics*, vol. 4, no. 1, pp. 81–88 (2012).
- [19] E. D. W. Diop, "On essential subsemimodules and weakly co-Hopfian semimodules," *European Journal of Pure and Applied Mathematics*, vol. 9, no. 3, pp. 250–265 (2016).
- [20] S. E. Atani and F. E. K. Saraei, "On coatomic semimodules over commutative semirings," *Cankaya University Journal of Science and Engineering*, vol. 8, no. 2 (2011).
- [21] M. M. Altaee, π -injective, π -projective semimodules and other related concepts (Ph.D. thesis), University of Babylon, Babil, Iraq (2019).
- [22] R. E. Atani, "Prime subsemimodules of semimodules," *International Journal of Algebra*, vol. 4, no. 26, pp. 1299–1306 (2010).
- [23] R. R. Nazari and S. Ghalandarzadeh, "Multiplication semimodules," *Acta Univ. Sapientiae, Mathematica*, vol. 11, no. 1, pp. 172–185 (2019).

Received November, 2025