

Two-step iterative method for nonlinear equations using Caputo-Fabrizio fractional derivatives

Tania Chib [†]

Surjeet Singh Chauhan (Gonder) ^{*}

Department of Mathematics

University Institute of Sciences (UIS)

Chandigarh University

Mohali 140413

Punjab

India

Abstract

The main objective of this paper is to develop a two-step fractional Newton-Raphson method for solving non-linear equations using the Caputo-Fabrizio fractional derivative. In further detail, we discuss the convergence analysis for the proposed method. In addition to it, we provide an example to compare the method we proposed with the existing approaches.

Subject Classification: 47H10, 47J26.

Keywords: Nonlinear equations, Convergence analysis, Numerical analysis, Caputo-Fabrizio fractional derivative.

1. Introduction

Fractional calculus and nonlinear equations play significant role in arriving of the branches of science and engineering, including applied mathematics, physics, biology, and economics. Instead of classical calculus, fractional derivatives include non-local and memory-dependent properties [1, 2, 3, 4, 5, 6, 7, 8, 9]. Among the various fractional derivatives proposed, the Caputo-Fabrizio derivative has gained significant attention due to its nonsingular kernel [10, 11, 12, 13, 14, 15, 16, 17, 18, 19].

In this paper, we present a two-step fractional Newton-Raphson approach for solving nonlinear equations based on Caputo-Fabrizio fractional derivative. The method proposed here involves an extension of the traditional Newton-Raphson method by utilizing the properties of Caputo-Fabrizio derivative. The prime contribution is the derivation of the two-step fractional

[†] E-mail: chibtania26@gmail.com

^{*} E-mail: surjeetschauhan@yahoo.com (Corresponding Author)

Newton-Raphson method based on the Caputo-Fabrizio derivative and to check its convergence analysis at a local convergence rate of atleast 3α and also a numerical example is provided with its comparison with existing numerical methods.

In Section 2 we state existing results and Section 3 provides the main results that we use to formulate the two-step fractional Newton-Raphson method using the Caputo-Fabrizio derivative that we introduce in section 4. Section 5 provides an example of the application of the proposed method in comparison to previous existing approaches.

2. Preliminaries

We recall some existing results of Caputo fractional derivative and Caputo–Fabrizio fractional derivative in this paper that will be used in the development of our method.

Definition 2.1: (Caputo Fractional Derivative) [14, 15]

$${}^C D_x^\alpha f(x) = \frac{1}{\Gamma(1-\alpha)} \int_0^x (x-\tau)^{-\alpha} f'(\tau) d\tau, \quad (1)$$

in which $0 < \alpha < 1$ is the fractional order, and $\Gamma(s) = \int_0^{+\infty} e^{-t} t^{s-1} dt$ is the Gamma function.

Definition 2.2: (Caputo–Fabrizio Fractional Derivative)[8]. Let $f \in H^1(a, b)$ and $0 < \alpha < 1$. The Caputo–Fabrizio fractional derivative of order α is defined as

$${}^{CF} D_a^\alpha f(t) = \frac{M(\alpha)}{1-\alpha} \int_a^x f'(s) \exp\left(-\frac{\alpha}{1-\alpha}(x-s)\right) ds, \quad (2)$$

where $M(\alpha)$ is a normalization factor having $M(0) = M(1) = 1$.

Lemma [Two-step fractional Newton-Raphson method 1 (TSFNRM1)] [12, 16, 17].

Algorithm: Two-step Caputo fractional Newton-Raphson Method (TSFNRM).

Initially x_0 is sufficiently close to $x_{(i+1)}$ by the following Caputo type two-step scheme:

$$y_i = x_i - \Gamma(\alpha + 1) \frac{f(x_i)}{{}^C D_x^\alpha f(x_i)},$$

$$x_{i+1} = y_i - \Gamma(\alpha + 1) \frac{f(x_i)}{{}^{CF} D_x^\alpha f(x_i)}, \quad i = 0, 1, 2, \dots$$

3. Main Results

Theorem 1: Suppose that $f \in C^{n+1}[a, b]$ and ${}^{CF} D_a^\alpha f(x)$ be the Caputo–Fabrizio fractional derivative of order $0 < \alpha \leq 1$, and let ${}^{CF} D_a^{j\alpha} f(x) \in C[a, b]$ for all $j = 1, 2, \dots, n+1$. Then,

$$f(x) = \sum_{k=0}^n {}^{CF}D_a^{k\alpha} f(a) \cdot \frac{(x-a)^{k\alpha}}{\Gamma(k\alpha+1)} + R_n(x), \quad (3)$$

where $R_n(x) = {}^{CF}D_a^{(n+1)\alpha} f(\xi) \cdot \frac{(x-a)^{(n+1)\alpha}}{\Gamma((n+1)\alpha+1)}$, $\forall \xi$ such that $a \leq \xi \leq b$ is the remainder term.

Proof: Let $n = 0$. Then, equation (3) reduces to the Caputo-Fabrizio fractional mean value theorem:

$$f(x) = f(a) + {}^{CF}D_a^\alpha f(\xi) \frac{(x-a)^\alpha}{\Gamma(\alpha+1)}. \quad (4)$$

Using (2) and integrating both sides from a to x , we have:

$$\int_a^x {}^{CF}D_a^\alpha f(s) ds = \frac{M(\alpha)}{1-\alpha} \int_a^x \int_a^s f'(\tau) \exp\left(-\frac{\alpha(s-\tau)}{1-\alpha}\right) d\tau ds. \quad (5)$$

Here,

$$\int_a^x \int_a^s f'(\tau) \exp\left(-\frac{\alpha(s-\tau)}{1-\alpha}\right) d\tau ds = \int_a^x f'(\tau) \frac{1-\alpha}{\alpha} \left[1 - \exp\left(-\frac{\alpha(x-\tau)}{1-\alpha}\right)\right] d\tau \quad (6)$$

Now (5) implies that

$$\int_a^x {}^{CF}D_a^\alpha f(s) ds = \frac{M(\alpha)}{1-\alpha} \int_a^x f'(\tau) \frac{1-\alpha}{\alpha} \left[1 - \exp\left(-\frac{\alpha(x-\tau)}{1-\alpha}\right)\right] d\tau \quad (7)$$

As

$$f(x) - f(a) = \int_a^x f'(\tau) d\tau \quad (8)$$

Using (8), (7) becomes

$$\int_a^x {}^{CF}D_a^\alpha f(s) ds = \frac{M(\alpha)}{1-\alpha} \left[f(x) - f(a) - \int_a^x f'(\tau) \exp\left(-\frac{\alpha(x-\tau)}{1-\alpha}\right) d\tau \right] \quad (9)$$

On simplifying and by applying mean value theorem, we have,

$$f(x) - f(a) = {}^{CF}D_a^\alpha f(\xi) \frac{1-\alpha}{M(\alpha)} \int_a^x \exp\left(-\frac{\alpha(x-s)}{1-\alpha}\right) ds \quad (10)$$

Upon simplification the integral $\int_a^x \exp\left(-\frac{\alpha(x-s)}{1-\alpha}\right) ds$ we have,

$$\int_a^x \exp\left(-\frac{\alpha(x-s)}{1-\alpha}\right) ds = \frac{1-\alpha}{\alpha} \left[1 - \exp\left(-\frac{\alpha(x-a)}{1-\alpha}\right)\right] \quad (11)$$

For $\alpha \approx 1$, $\exp\left(-\frac{\alpha(x-a)}{1-\alpha}\right) \approx 0$, we have, $f(x) - f(a) \approx {}^{CF}D_a^\alpha f(\xi) \frac{(x-a)^\alpha}{\Gamma(\alpha+1)}$ and now we assume that it holds for $n=m$ and for $n=m+1$ we have to expand $f(x)$ to m terms

$$f(x) = \sum_{k=0}^m {}^{CF}D_a^{k\alpha} f(a) \cdot \frac{(x-a)^{k\alpha}}{\Gamma(k\alpha+1)} + R_m(x) \quad (12)$$

where,

$$R_m(x) = \frac{1}{\Gamma(m\alpha+1)} \int_a^x (x-\tau)^{m\alpha} {}^{CF}D_a^{(m+1)\alpha} f(\tau) d\tau \quad (13)$$

Now, we have to express (13) as an integral. In order to do so let $m=0$ in (12) which implies that $f(x) = f(a) + R_0(x)$ where $R_0(x)$ is obtained by integrating Caputo-Fabrizio derivative,

$$R_0(x) = \frac{1-\alpha}{\alpha} \int_a^x f'(\tau) \left[1 - \exp\left(-\frac{\alpha(x-\tau)}{1-\alpha}\right) \right] d\tau \quad (14)$$

Now, for $\alpha \approx 1$, $\exp\left(-\frac{\alpha(x-\tau)}{1-\alpha}\right) \approx 0$ (14) become

$$R_m(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-\tau)^{\alpha-1} {}^{CF}D_a^\alpha f(\tau) d\tau \quad (15)$$

By assuming that it holds for m and considering for $m+1$ we get that

$$R_m(x) = \frac{1}{\Gamma(m+1)\alpha} \int_a^x (x-\tau)^{(m+1)\alpha-1} {}^{CF}D_a^\alpha f(\tau) d\tau \quad (16)$$

This implies that it is true for $m+1$. Now by applying mean value theorem and after simplifying it we have,

$$R_m(x) = \frac{{}^{CF}D_a^{(n+1)\alpha} f(\xi)}{\Gamma((m+1)\alpha+1)} (x-a)^{(m+1)\alpha} \quad (17)$$

which is our remainder term and (12) and (17) proves our theorem.

Theorem 2: Let $f: D \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function in a domain D that contains the root $\bar{x}$ of $f(x)$. For any positive integers k and α , $\alpha \in (0,1]$, it possesses Caputo-Fabrizio fractional derivatives of order 2α . Also, assume that ${}^{CF}D_x^\alpha f(x)$ is a continuous, nonzero fractional derivative of Caputo-Fabrizio type in a neighbourhood of $\bar{x}$ and the initial guess x_0 is sufficiently close to $\bar{x}$, then

$$x_{k+1} = x_k - M_\alpha \frac{f(x_k)}{{}^{CF}D_x^\alpha f(x_k)}, \quad k = 0, 1, \dots, \quad (18)$$

converges locally at a rate of at least 2α , where M_α is a normalization constant with Caputo-Fabrizio kernel and the error equation is given by

$$e_{k+1} = M(\alpha) \frac{{}^{CF}D_x^{2\alpha} f(\bar{x}) [\Gamma(1+2\alpha) - \Gamma(1+\alpha)^2]}{{}^{CF}D_x^\alpha f(\bar{x}) [\Gamma(1+\alpha)^2 - \Gamma(1+2\alpha)]} e_k^{2\alpha} + O(e_k^{3\alpha}). \quad (19)$$

Proof: By using Caputo-Fabrizio based Taylor series expansion of a function $f(x)$ centered at $a = \bar{x}$

$$f(x_k) = f(\bar{x}) + \frac{{}^{CF}D_x^{2\alpha} f(\bar{x})}{\Gamma(1+\alpha)} e_k^\alpha + \frac{{}^{CF}D_x^{2\alpha} f(\bar{x})}{\Gamma(1+2\alpha)} e_k^{2\alpha} + O(e_k^{3\alpha}).$$

Since $f(\bar{x}) = 0$, then

$$f(x_k) = \frac{{}^{CF}D_x^\alpha f(\bar{x})}{\Gamma(1+\alpha)} e_k^\alpha + \frac{{}^{CF}D_x^{2\alpha} f(\bar{x})}{\Gamma(1+2\alpha)} e_k^{2\alpha} + O(e_k^{3\alpha}) \quad (20)$$

Similarly, by expanding Caputo-Fabrizio derivative at x_k around $\bar{x}$, we have

$${}^{CF}D_x^\alpha f(x_k) = {}^{CF}D_x^\alpha f(\bar{x}) + \frac{{}^{CF}D_x^\alpha f(\bar{x})}{\Gamma(1+\alpha)} e_k^\alpha + O(e_k^{2\alpha}) \quad (21)$$

Subtracting $\bar{x}$ on both sides of (18), we have the following error equation

$$e_{k+1} = e_k - M(\alpha) \frac{f(x_k)}{{}^{CF}D_x^\alpha f(x_k)} \quad (22)$$

Substituting (20) and (21) in (22)

$$e_{k+1} = e_k - M(\alpha) \frac{\frac{{}^{CF}D_{\bar{x}}^{\alpha} f(\bar{x})}{\Gamma(1+\alpha)} e_k^{\alpha} + \frac{{}^{CF}D_{\bar{x}}^{2\alpha} f(\bar{x})}{\Gamma(1+2\alpha)} e_k^{2\alpha} + O(e_k^{3\alpha})}{{}^{CF}D_{\bar{x}}^{\alpha} f(\bar{x}) + \frac{{}^{CF}D_{\bar{x}}^{2\alpha} f(\bar{x})}{\Gamma(1+\alpha)} e_k^{\alpha} + O(e_k^{2\alpha})}$$

To attain a convergence rate of 2α , it is necessary that Caputo-Fabrizio fractional Newton's method leads to the following resulting error equation:

$$e_{k+1} = M(\alpha) \frac{{}^{CF}D_{\bar{x}}^{2\alpha} f(\bar{x}) [\Gamma(1+2\alpha) - \Gamma(1+\alpha)^2]}{{}^{CF}D_{\bar{x}}^{\alpha} f(\bar{x}) [\Gamma(1+\alpha)^2 - \Gamma(1+2\alpha)]} e_k^{2\alpha} + O(e_k^{3\alpha}) \quad (23)$$

This completes the proof.

4. New Fractional Numerical Techniques

In this section, we introduce two-step fractional Newton-Raphson method using Caputo-Fabrizio fractional derivative which is modified version of Newton-Raphson technique.

Algorithm 1: Two-step Caputo-Fabrizio fractional Newton-Raphson method (TSFCFNR).

For an initial approximation x_0 sufficiently close to $x_{(k+1)}$ by the following Caputo-Fabrizio type two-step scheme:

$$y_k = x_k - \Gamma(\alpha + 1) \frac{f(x_k)}{{}^{CF}D_{\bar{x}}^{\alpha} f(x_k)}, \quad (24)$$

$$x_{k+1} = y_k - \Gamma(\alpha + 1) \frac{f(x_k)}{{}^{CF}D_{\bar{x}}^{\alpha} f(x_k)}, \quad k = 0, 1, \dots \quad (25)$$

Now, the following result gives the convergence analysis of Algorithm 1.

Theorem 3: Let us suppose that $f: D \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function in a domain D containing the root $\bar{x}$ of $f(x)$. For any positive integers k and α , $\alpha \in (0, 1]$, suppose that it possesses fractional derivatives of order $k\alpha$ in the Caputo-Fabrizio. Also, assume that ${}^{CF}D_{\bar{x}}^{\alpha} f(x)$ is a continuous, nonzero fractional derivative of Caputo-Fabrizio type at $\bar{x}$. If the initial guess x_0 is sufficiently close to $\bar{x}$, then Algorithm 1 has a local convergence rate of at least 3α , and the error equation is:

$$e_{k+1} = \frac{({}^{CF}D_{\bar{x}}^{2\alpha} f(x_k))^2 (-M(\alpha)\Gamma(\alpha+1) \cdot {}^{CF}D_{\bar{x}}^{\alpha} f(x_k) + \Gamma(2\alpha+1))}{{}^{CF}D_{\bar{x}}^{\alpha} f(x_k)^4 \Gamma(\alpha+1)^2 \Gamma(2\alpha+1)} e_k^{3\alpha} + O(e_k^{4\alpha}) \quad (26)$$

Proof: Let $\{x_k\}_{k \geq 0}$, obtained by (18), be a sequence of iterations produced by $\bar{x}$ of $f(x)$. Expanding $f(x)$ and its Caputo-Fabrizio derivative at x_k , around $\bar{x}$ yields as (20) and (21).

Thus, a Caputo-Fabrizio Newton iteration can be derived and represented in terms of the approximation error $e_k = x_k - \bar{x}$ at the k -th iteration as

$$\frac{f(x_k)}{{}^{CF}D_{\bar{x}}^{\alpha}f(x_k)} = \frac{\frac{{}^{CF}D_{\bar{x}}^{\alpha}f(\bar{x})}{\Gamma(1+\alpha)}e_k^{\alpha} + \frac{{}^{CF}D_{\bar{x}}^{2\alpha}f(\bar{x})}{\Gamma(2\alpha+1)}e_k^{2\alpha} + O(e_k^{3\alpha})}{{}^{CF}D_{\bar{x}}^{\alpha}f(\bar{x}) + \frac{{}^{CF}D_{\bar{x}}^{2\alpha}f(\bar{x})}{\Gamma(\alpha+1)}e_k^{\alpha} + O(e_k^{2\alpha})} \quad (27)$$

So, from (24), we have,

$$y_k = \bar{x} + e_k - \Gamma(\alpha + 1) \frac{f(x_k)}{{}^{CF}D_{\bar{x}}^{\alpha}f(x_k)} = \frac{{}^{CF}D_{\bar{x}}^{2\alpha}f(\bar{x})[\Gamma(2\alpha+1)-\Gamma(\alpha+1)]}{{}^{CF}D_{\bar{x}}^{\alpha}f(\bar{x})[\Gamma(\alpha+1)^2-\Gamma(2\alpha+1)]}e_k^{2\alpha} + O(e_k^{3\alpha}) \quad (28)$$

From (20) and (28), we have,

$$f(y_k) = \frac{{}^{CF}D_{\bar{x}}^{2\alpha}f(\bar{x})[\Gamma(\alpha+1)-\Gamma(2\alpha+1)]}{\Gamma(\alpha+1)^2-\Gamma(2\alpha+1)}e_k^{2\alpha} + O(e_k^{3\alpha}) \quad (29)$$

From (21) and (29), we have,

$$\frac{f(y_k)}{{}^{CF}D_{\bar{x}}^{\alpha}f(x_k)} = \frac{{}^{CF}D_{\bar{x}}^{2\alpha}f(\bar{x})[-\Gamma(\alpha+1)+\Gamma(2\alpha+1)]}{({}^{CF}D_{\bar{x}}^{\alpha}f(\bar{x}))^2\Gamma(\alpha+1)^2\Gamma(2\alpha+1)}e_k^{2\alpha} + O(e_k^{3\alpha}) \quad (30)$$

So, to attain a convergence rate of 3α , it is evident that the two step Caputo-Fabrizio fractional derivative must give $M(\alpha)$ as a normalization factor, and the error equation is:

$$e_{k+1} = y_k - M(\alpha) \frac{f(y_k)}{{}^{CF}D_{\bar{x}}^{\alpha}f(x_k)}, \quad (31)$$

which gives us

$$e_{k+1} = \frac{({}^{CF}D_{\bar{x}}^{2\alpha}f(x_k))^2(-M(\alpha)\Gamma(\alpha+1) \cdot {}^{CF}D_{\bar{x}}^{\alpha}f(x_k)+\Gamma(2\alpha+1))}({}^{CF}D_{\bar{x}}^{\alpha}f(x_k))^4\Gamma(\alpha+1)^2\Gamma(2\alpha+1)}e_k^{3\alpha} + O(e_k^{4\alpha}) \quad (32)$$

This completes the proof.

5. Numerical example of proposed method

In this section, we provided the numerical example of the proposed method. The system that we used for numerical results is Matlab R2014b. Here, Table 1 provides the numerical outcomes of TSFNRM1 and TSFCFNRM for different values of α , x_n is the root root of function, iter is for iteration count, $|x_n - x_{n-1}|$ for absolute error and $|f(x_n)|$ is the functional value.

Table 1

Numerical outcomes of TSFNRM1 and TSFCFNRM for $f(x) = x^3 - 10$.

TSFNRM1					TSFCFNRM		
α	x_n	Iter	$ x_n - x_{n-1} $	$ f(x_n) $	Iter	$ x_n - x_{n-1} $	$ f(x_n) $
0.2	2.154434689989885277	7	1.06×10^{-10}	4.18×10^{-9}	7	3.72×10^{-10}	5.62×10^{-11}
0.3	2.154434690018297478	6	5.26×10^{-10}	6.48×10^{-10}	6	3.92×10^{-11}	5.49×10^{-12}

Contd...

0.4	2.154434690030900574	5	2.43×10^{-10}	5.55×10^{-11}	5	8.39×10^{-11}	1.16×10^{-13}
0.5	2.154434690032031676	4	3.68×10^{-11}	5.54×10^{-11}	4	7.01×10^{-11}	9.76×10^{-13}
0.6	2.154434690032108962	5	4.57×10^{-11}	2.59×10^{-12}	5	5.15×10^{-12}	7.18×10^{-14}
0.7	2.154434690035180481	5	4.32×10^{-11}	6.02×10^{-12}	5	1.40×10^{-11}	1.95×10^{-13}
0.8	2.154434690034441470	5	3.71×10^{-11}	5.24×10^{-13}	5	1.28×10^{-11}	1.78×10^{-13}
0.9	2.154434690037226093	5	3.08×10^{-12}	1.95×10^{-14}	5	1.09×10^{-12}	1.52×10^{-14}

6. Conclusion

This study presents two-step fractional Newton-Raphson method using Caputo-Fabrizio fractional derivative. The convergence properties of the proposed method were established to show that this method have a higher convergence rate as compared to classical Newton- Raphson method and it have a convergence rate of at least 3α and a numerical example is provided to check the accuracy of the proposed method. This result confirms that Caputo-Fabrizio method have potential in solving nonlinear equations. Future research may explore the higher order extensions and multi dimensional system with its applications.

References

- [1] N. Z. Ali, A. G. Khan, M. U. Awan, L. Ciurdariu, and K. Brahim, "Design and applicability of two-step fractional Newton-Raphson method," *Fractal Fract.*, vol. 8, no. 10, Art. no. 582 (2024).
- [2] G. Jumarie, "Modified Riemann-Liouville derivative and fractional Taylor series of nondifferentiable functions: Further results," *Comput. Math. Appl.*, vol. 51, nos. 9-10, pp. 1367-1376 (2006).
- [3] J. H. Mathews and K. D. Fink, *Numerical Methods Using MATLAB*, 4th ed. Upper Saddle River, NJ, USA: Prentice Hall (2004).
- [4] Z. M. Odibat and N. T. Shawagfeh, "Generalized Taylor's formula," *Appl. Math. Comput.*, vol. 186, no. 1, pp. 286-293 (2007), doi: 10.1016/j.amc.2006.07.102.
- [5] M. Vivas-Cortez, N. Z. Ali, A. G. Khan, and M. U. Awan, "Numerical analysis of new hybrid algorithms for solving nonlinear equations," *Axioms*, vol. 12, no. 7, Art. no. 684 (2023).

- [6] A. Torres-Hernandez and F. Brambila-Paz, "Fractional Newton-Raphson method," *arXiv preprint arXiv:1710.07634* (2017).
- [7] A. Akgül, A. Cordero, and J. R. Torregrosa, "A fractional Newton method with 2α th-order of convergence and its stability," *Appl. Math. Lett.*, vol. 98, pp. 344-351 (2019).
- [8] N. Ali, M. Waseem, M. Safdar, A. Akgül, and F. T. Tolasa, "Iterative solutions for nonlinear equations via fractional derivatives: Adaptations and advances," *Appl. Math. Sci. Eng.*, vol. 32, no. 1, Art. no. 2333816 (2024).
- [9] A. Torres-Hernandez and F. Brambila-Paz, "Fractional Newton-Raphson method and some variants for the solution of nonlinear systems," *arXiv preprint arXiv:1908.01453* (2019).
- [10] Z. Cui, "Solutions of some typical nonlinear differential equations with Caputo-Fabrizio fractional derivative," *AIMS Math.*, vol. 7, no. 8, pp. 14139-14153 (2022).
- [11] G. S. Teodoro, J. A. Tenreiro Machado, and E. Capelas de Oliveira, "A review of definitions of fractional derivatives and other operators," *J. Comput. Phys.*, vol. 388, pp. 195-208 (2019), doi: 10.1016/j.jcp.2019.03.008.
- [12] M. D. Ortigueira and J. A. Tenreiro Machado, "What is a fractional derivative?" *J. Comput. Phys.*, vol. 293, pp. 4-13 (2015), doi: 10.1016/j.jcp.2014.07.019.
- [13] H. Yépez-Martínez and J. F. Gómez-Aguilar, "A new modified definition of Caputo-Fabrizio fractional-order derivative and their applications to the Multi Step Homotopy Analysis Method (MHAM)," *J. Comput. Appl. Math.*, vol. 346, pp. 247-260 (2019), doi: 10.1016/j.cam.2018.07.023.
- [14] R. Almeida, "A Caputo fractional derivative of a function with respect to another function," *Commun. Nonlinear Sci. Numer. Simul.*, vol. 44, pp. 460-481 (2017).
- [15] B. S. Katoch and S. S. Chauhan, "Existence of the solution for the Caputo-Fabrizio boundary value problem along with nonlinear integral boundary conditions," in *AIP Conf. Proc.*, vol. 3283, no. 1, Art. no. 040003 (2025).
- [16] Y. Dimitrov, S. Georgiev, and V. Todorov, "Approximation of Caputo fractional derivative and numerical solutions of fractional differential equations," *Fractal Fract.*, vol. 7, no. 10, Art. no. 750 (2023).

- [17] I. Yadav and R. Bhardwaj, "Numerical simulation of nonlinear equations by modified bisection and Regula Falsi method," *Proc. Pakistan Acad. Sci. A Phys. Comput. Sci.*, vol. 62, no. 1, pp. 11-19 (2025).
- [18] S. S. Chauhan, P. Tyagi, and N. Mani, "Rational contraction mapping in F-modular b-metric spaces," *J. Interdiscip. Math.*, vol. 27, no. 8, pp. 1781-1786 (2024), doi: 10.47974/JIM-2041.
- [19] S. Kulkarni, U. Kadam, D. K. Paswan, K. Shipra, B. C. Julme, and S. L. Dhondge, "Numerical stability analysis of nonlinear differential equations in engineering systems," *J. Interdiscip. Math.*, vol. 28, no. 6, pp. 2255-2261 (2025), doi: 10.47974/JIM-2367.

Received February, 2026