

A new kind of topological vector space via proximit structure

Dhuha Abdulameer Kadhim *

*Department of Mathematics
College of Education for Girls
University of Kufa
Iraq*

Boushra Y. Hussein [†]

*Department of Mathematics
College of Education
University of Al-Qadisiyah
Iraq*

Abstract

The concepts of Dh-contraction, proximit semigroup, proximit group, proximit vector space and topological proximit space are introduced. There are a few instances and properties provided. Convergence and Sequential Dh-contraction are both specified, in addition to the terms Dh-contraction and Sequential Dh-contraction are interchangeable.

Subject Classification: Primary 15A60, Secondary 47H09, 46A99.

Keywords: Dh-contraction, Proximit group, Proximit vector space, Proximit space.

1. Introduction

Approach spaces have some applicative backgrounds in all major of mathematics containing probability theory [15], domain theory [16], group theory [13] and vector spaces [14]. It is commonly known that the category met

* Orcid: <https://orcid.org/0000-0002-9152-4162>

[†] Orcid: <https://orcid.org/0000-0003-4909-0126>

* E-mail: dhuha.ebada@uokufa.edu.iq (Corresponding Author)

[†] E-mail: boushra.alshebani@qu.edu.iq

of metric spaces and inexpensive maps lacks infinite products and co-products. In order to address this issue, Robert Lowen [10] created approach spaces in 1989. These spaces are a generalization of metric and topology based on a distance that can be described in a variety of comparable ways, such as in terms of limit, gauge, and distance function between points and sets. Approach spaces [12,17] can be used to define closure operators in topology, extended pseudo-quasi-metrics for defining coarser topologies, and filter limit points, respectively. R. Lowen and K. Robeys [11] established some basic definitions and theorems in approach space which is a topological and metric space extension. More information about approach space can be found in [9]. M. Sion and W. Haute [18] investigated an alternative notion of approach spaces based on approach cores. [19] presented the relationship between approach space and metric space, which is similar to the relationship between topological and ordered space and he presented a sober metric approach space. A new kind of topological vector space which is called approach vector space made by [2]. Furthermore, Abbas and Hussein worked the completion normed approach space in [1]. Additional information can be found in [3,4,5,6,7]. This research looks into the regularity of functionals between sets. In section 2, we introduce some concepts of proximit space and we look at some of the findings and discuss the concept of Dh-contraction. Section 3 introduces the concept of proximit semigroup and proximit group. We also provide basic features and examples. Finally, topological proximit vector space, Dh-contraction, and proximit vector space are demonstrated.

2. Preliminaries

Definition 2.1: [12] A collection of functions $t_\varepsilon: 2^{\mathbb{X}} \rightarrow 2^{\mathbb{X}}$, $\varepsilon \in R^+$ is called a tower on $\mathbb{X}$ if the following conditions are satisfied :

- (1) $\forall \hat{\mathcal{A}} \in 2^{\mathbb{X}}, \forall \varepsilon \in R^+: \hat{\mathcal{A}} \subset t_\varepsilon(\hat{\mathcal{A}})$,
- (2) $\forall \varepsilon \in R^+: t_\varepsilon(\emptyset) = \emptyset$
- (3) $\forall \hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^{\mathbb{X}}, \forall \varepsilon \in R^+: t_\varepsilon(\hat{\mathcal{A}} \cup \hat{\mathcal{B}}) = t_\varepsilon(\hat{\mathcal{A}}) \cup t_\varepsilon(\hat{\mathcal{B}})$,
- (4) $\forall \hat{\mathcal{A}} \in 2^{\mathbb{X}}, \forall \varepsilon, \gamma \in R^+: t_\varepsilon(t_\gamma(\hat{\mathcal{A}})) \subset t_{\varepsilon+\gamma}(\hat{\mathcal{A}})$,
- (5) $\forall \hat{\mathcal{A}} \in 2^{\mathbb{X}}, \forall \varepsilon \in R^+: t_\varepsilon(\hat{\mathcal{A}}) = \bigcap_{\varepsilon < \gamma} t_\gamma(\hat{\mathcal{A}})$,

From (3) and (5), Then $\forall \hat{\mathcal{A}} \subset \hat{\mathcal{B}} \subset \mathbb{X}, \forall \alpha \leq \beta \in R^+: t_\alpha(\hat{\mathcal{A}}) \subset t_\beta(\hat{\mathcal{B}})$.

Definition 2.2: [12] An operator $cl: 2^{\mathbb{X}} \rightarrow 2^{\mathbb{X}}$ that determines the qualities of a pretopology on a set $\mathbb{X}$ for all $\mathcal{A}, \mathcal{B} \in 2^{\mathbb{X}}$ is determined by an operator that determines the properties of an operator on a set.

1. $\hat{\mathcal{A}} \subset cl(\hat{\mathcal{A}})$
2. $cl(\emptyset) = \emptyset$,
3. $cl(\hat{\mathcal{A}} \cup \hat{\mathcal{B}}) = cl(\hat{\mathcal{A}}) \cup cl(\hat{\mathcal{B}})$.

This operator is said to be a pretopological closure operator. We call a set $\mathbb{X}$ with a pretopology a pretopological space.

Definition 2.3: [8] A functional $\rho: 2^{\mathbb{X}} \times 2^{\mathbb{X}} \rightarrow [0, \infty]$ is said to be an Dh-functional on $\mathbb{X}$ ($\mathbb{X}$ is a set) if and only if the following conditions are satisfies:

- (G1) $\forall \hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^{\mathbb{X}}: \rho(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = \rho(\hat{\mathcal{B}}, \hat{\mathcal{A}}),$
- (G2) $\forall \hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^{\mathbb{X}}, \hat{\mathcal{A}} = \emptyset \text{ or } \hat{\mathcal{B}} = \emptyset \Rightarrow \rho(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = \infty,$
- (G3) $\forall \hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^{\mathbb{X}}, \rho(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = 0 \Rightarrow \hat{\mathcal{A}} \cap \hat{\mathcal{B}} \neq \emptyset,$
- (G4) $\forall \hat{\mathcal{A}}, \hat{\mathcal{B}}, \hat{\mathcal{C}} \in 2^{\mathbb{X}}: \rho(\hat{\mathcal{A}}, \hat{\mathcal{B}} \cup \hat{\mathcal{C}}) = \min\{\rho(\hat{\mathcal{A}}, \hat{\mathcal{B}}), \rho(\hat{\mathcal{A}}, \hat{\mathcal{C}})\},$
- (G5) $\forall \hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^{\mathbb{X}}, \forall \varepsilon, \eta \in [0, \infty]: \rho(\hat{\mathcal{A}}, \hat{\mathcal{B}}) \leq \rho(\hat{\mathcal{A}}^\varepsilon, \hat{\mathcal{B}}^\eta) + \varepsilon + \eta.$

For every $\hat{\mathcal{A}} \in 2^{\mathbb{X}}, \varepsilon \in [0, \infty]$, we write $t_\varepsilon(\hat{\mathcal{A}}) := \{\mathfrak{x} \in \mathbb{X} \mid \rho(\{\mathfrak{x}\}, \hat{\mathcal{A}}) \leq \varepsilon\}$. Therefore the triple $(\mathbb{X}, \rho, t_\varepsilon)$ is called proximit space.

Example 2.4: [8] Let $\mathbb{X} = [0, \infty]$, define $\rho: 2^{[0, \infty]} \times 2^{[0, \infty]} \rightarrow [0, \infty]$ by

$$\rho(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = \begin{cases} 0 & \hat{\mathcal{A}}, \hat{\mathcal{B}} \text{ unbounded} \\ \infty & \hat{\mathcal{A}}, \hat{\mathcal{B}} \text{ bounded} \\ \inf_{n \in \hat{\mathcal{A}}} \inf_{m \in \hat{\mathcal{B}}} |n - m| & n, m < \infty \end{cases}$$

Then $(\mathbb{X}, \rho, t_\varepsilon)$ is proximit space.

Definition 2.5: A function $f: (\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon) \rightarrow (\mathbb{Y}, \rho_{\mathbb{Y}}, t_\varepsilon)$ such that $(\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon)$ and $(\mathbb{Y}, \rho_{\mathbb{Y}}, t_\varepsilon)$ are proximit space is called Dh-contraction if

$$f(t_{\varepsilon_{\mathbb{X}}}(\hat{\mathcal{A}})) \subseteq t_{\varepsilon_{\mathbb{Y}}}(f(\hat{\mathcal{A}})) \quad \forall \hat{\mathcal{A}} \subseteq \mathbb{X}, \forall \varepsilon \in R^+.$$

Proposition 2.6: Suppose that $(\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon)$ is proximit spaces. Then identity map $I_{\mathbb{X}}: (\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon) \rightarrow (\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon) \quad \forall \hat{\mathcal{A}} \subseteq \mathbb{X}, \forall \varepsilon \in R^+$ is Dh-contraction.

Proof: Let $\hat{\mathcal{A}} \subseteq \mathbb{X}, \varepsilon \in R^+$ such that $I_{\mathbb{X}}(\hat{\mathcal{A}}) = \hat{\mathcal{A}}$. Let $\mathfrak{x} \in I_{\mathbb{X}}(t_\varepsilon(\hat{\mathcal{A}}))$, then $\mathfrak{x} \in t_\varepsilon(\hat{\mathcal{A}})$. Hence $\mathfrak{x} \in t_\varepsilon(I_{\mathbb{X}}(\hat{\mathcal{A}}))$, So $I_{\mathbb{X}}$ is Dh-contraction.

Proposition 2.7: Suppose that $(\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon)$, $(\mathbb{Y}, \rho_{\mathbb{Y}}, t_\varepsilon)$ and $(\mathbb{Z}, \rho_{\mathbb{Z}}, t_\varepsilon)$ are proximit spaces. If the function $h: (\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon) \rightarrow (\mathbb{Y}, \rho_{\mathbb{Y}}, t_\varepsilon)$ and $j: (\mathbb{Y}, \rho_{\mathbb{Y}}, t_\varepsilon) \rightarrow (\mathbb{Z}, \rho_{\mathbb{Z}}, t_\varepsilon)$ are Dh-contraction, then $j \circ h$ is Dh-contraction.

Proof: Let $\hat{\mathcal{A}} \subseteq \mathbb{X}, \varepsilon \in R^+$ such that $j \circ h: (\mathbb{X}, \rho_{\mathbb{X}}, t_\varepsilon) \rightarrow (\mathbb{Z}, \rho_{\mathbb{Z}}, t_\varepsilon)$ be a function. Let $\mathfrak{x} \in (j \circ h)(t_{\varepsilon_{\mathbb{X}}}(\hat{\mathcal{A}}))$, then $\mathfrak{x} \in (j(h(t_{\varepsilon_{\mathbb{X}}}(\hat{\mathcal{A}}))))$. But h is Dh-contraction, it follows that $\mathfrak{x} \in j(t_{\varepsilon_{\mathbb{Y}}}(h(\hat{\mathcal{A}})))$. Also j is Dh-contraction, hence $\mathfrak{x} \in t_{\varepsilon_{\mathbb{Z}}}(j(h(\hat{\mathcal{A}})))$. Therefore $\mathfrak{x} \in t_{\varepsilon_{\mathbb{Z}}}(j \circ h(\hat{\mathcal{A}}))$. Then $(j \circ h)(t_{\varepsilon_{\mathbb{X}}}(\hat{\mathcal{A}})) \subseteq t_{\varepsilon_{\mathbb{Z}}}(j \circ h(\hat{\mathcal{A}}))$, so that $j \circ h$ is Dh-contraction.

Proposition 2.8: If $(\mathbb{X}, \rho_{\mathbb{X}}, t_{\varepsilon}), (\mathbb{Y}, \rho_{\mathbb{Y}}, t_{\varepsilon})$ is a proximit space and $h: (\mathbb{X}, \rho_{\mathbb{X}}, t_{\varepsilon}) \rightarrow (\mathbb{Y}, \rho_{\mathbb{Y}}, t_{\varepsilon})$ is Dh-contraction, then the restriction $h|_S$ is Dh-contraction $\forall S \subseteq \mathbb{X}$.

Proof: Assume that $h: (\mathbb{X}, \rho_{\mathbb{X}}, t_{\varepsilon}) \rightarrow (\mathbb{Y}, \rho_{\mathbb{Y}}, t_{\varepsilon})$ is Dh-contraction. We define restriction $j: S \rightarrow (\mathbb{Y}, \rho_{\mathbb{Y}}, t_{\varepsilon})$ as $j(v) = h(v) \forall v \in S$, assume $\mathfrak{x} \in j(t_{\varepsilon_{\mathbb{X}}}(\hat{\mathcal{A}}))$. Then $\mathfrak{x} \in h(t_{\varepsilon_{\mathbb{X}}}(\hat{\mathcal{A}}))$, but h is Dh-contraction. Hence $\mathfrak{x} \in t_{\varepsilon_{\mathbb{Y}}}(h(\hat{\mathcal{A}}))$, so $\mathfrak{x} \in t_{\varepsilon_{\mathbb{Y}}}(j(\hat{\mathcal{A}}))$. Thus $j(t_{\varepsilon_{\mathbb{X}}}(\hat{\mathcal{A}})) \subseteq t_{\varepsilon_{\mathbb{Y}}}(j(\hat{\mathcal{A}}))$, hence $h|_S$ is Dh-contraction.

3. Proximit Semi Group and Proximit Group

The terms "proximit semigroup," "proximit group," and "proximit subgroup" are defined in this section. Furthermore, instances and theorems are investigated.

Definition 3.1: A quadruple $(\mathbb{X}, \rho, t_{\varepsilon}, +)$ is said to be proximit semi group if and only if

1. $(\mathbb{X}, \rho, t_{\varepsilon})$ is proximit space.
2. $(\mathbb{X}, +)$ is semi group.
3. $+: \mathbb{X} \otimes \mathbb{X} \rightarrow \mathbb{X}$ such that $(\mathfrak{x}, \mathfrak{y}) \mapsto \mathfrak{x} + \mathfrak{y}$ is Dh-contraction.

Definition 3.2: A quadruple $(\mathbb{X}, \rho, t_{\varepsilon}, +)$ is said to be proximit group if and only if

1. $(\mathbb{X}, \rho, t_{\varepsilon})$ is proximit space.
2. $(\mathbb{X}, +)$ is group.
3. $+: \mathbb{X} \otimes \mathbb{X} \rightarrow \mathbb{X}$ such that $(\mathfrak{x}, \mathfrak{y}) \mapsto \mathfrak{x} + \mathfrak{y}$ is Dh-contraction.
4. $-: \mathbb{X} \rightarrow \mathbb{X}$ such that $\mathfrak{x} \mapsto -\mathfrak{x}$ is Dh-contraction.

Example 3.3: If R is the set of real numbers and $+$ is an addition binary operation with Dh-functional ρ , then $(R^n, \rho, t_{\varepsilon}, +)$ is proximit group.

Solve:

1. $(R^n, \rho, t_{\varepsilon})$ is proximit space.
2. $(R^n, +)$ is group.
3. We prove $+: R^n \otimes R^n \rightarrow R^n$ such that Dh-contraction is defined by $(\mathfrak{x}, \mathfrak{h}) \mapsto \mathfrak{x} + \mathfrak{h}$ for all $\mathfrak{x} = (\mathfrak{x}_1, \mathfrak{x}_2, \dots, \mathfrak{x}_n), \mathfrak{h} = (\mathfrak{h}_1, \mathfrak{h}_2, \dots, \mathfrak{h}_n)$.

Let $\hat{\mathcal{A}}, \hat{\mathcal{B}} \subseteq R^n, \varepsilon \in R^+$. Assume that $\mathfrak{x} \in f(t_{\varepsilon}(\hat{\mathcal{A}}), t_{\varepsilon}(\hat{\mathcal{B}}))$, then $\mathfrak{x} \in [t_{\varepsilon}(\hat{\mathcal{A}}) + t_{\varepsilon}(\hat{\mathcal{B}})]$. So that $\mathfrak{x} \in \min\{t_{\varepsilon}(\hat{\mathcal{A}}), t_{\varepsilon}(\hat{\mathcal{B}})\} = \min\{\rho(\{\mathfrak{x}\}, \hat{\mathcal{A}}) \leq \varepsilon, \rho(\{\mathfrak{x}\}, \hat{\mathcal{B}}) \leq \varepsilon\}$ it follows that $\min\{\rho(\{\mathfrak{x}\}, \hat{\mathcal{A}}), \rho(\{\mathfrak{x}\}, \hat{\mathcal{B}})\} \leq \varepsilon$ and $\rho(\{\mathfrak{x}\}, \hat{\mathcal{A}} + \hat{\mathcal{B}}) \leq \varepsilon$ therefore $\mathfrak{x} \in t_{\varepsilon}(\hat{\mathcal{A}} + \hat{\mathcal{B}})$. So $\mathfrak{x} \in t_{\varepsilon}(f(\hat{\mathcal{A}}), f(\hat{\mathcal{B}}))$

4. We show that $-:R^n \rightarrow R^n$ such that $\mathbf{x} \mapsto -\mathbf{x}$ is Dh-contraction. Let $\mathbf{x} \in f(t_\varepsilon(\hat{\mathcal{A}}))$, then $\mathbf{x} \in (-t_\varepsilon(\hat{\mathcal{A}}))$. So that $\rho(\{\mathbf{x}\}, -\hat{\mathcal{A}}) \leq \varepsilon$ implies that $\mathbf{x} \in t_\varepsilon(-\hat{\mathcal{A}})$. Thus $\mathbf{x} \in t_\varepsilon(f(\hat{\mathcal{A}}))$, so that $(R^n, \rho, t_\varepsilon, +)$ is proximit group.

Example 3.4: The set $C[a, b]$ of all continuous functions on the interval $[a, b]$. Then $(C[a, b], \rho, t_\varepsilon, +, \cdot)$ is proximit group.

$$\rho(\hat{\mathcal{A}}, G) = \begin{cases} \infty & \hat{\mathcal{A}} = \emptyset \text{ or } G = \emptyset \\ \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in G} |h(\mathbf{x}) - j(\mathbf{x})| & \hat{\mathcal{A}} \neq \emptyset \text{ and } G \neq \emptyset \end{cases}$$

Solve: Let $\hat{\mathcal{A}}, G, H \in 2^{C[a, b]}$

(G1) if $\hat{\mathcal{A}} \neq \emptyset, G \neq \emptyset$

$$\rho(\hat{\mathcal{A}}, G) = \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in G} |h(\mathbf{x}) - j(\mathbf{x})| = \sup_{j \in G} \sup_{h \in \hat{\mathcal{A}}} |j(\mathbf{x}) - h(\mathbf{x})| = \rho(\hat{\mathcal{A}}, G)$$

(G2) If $\hat{\mathcal{A}} = \emptyset, \rho(\hat{\mathcal{A}}, G) = \sup_{h \in \emptyset} \sup_{j \in G} |\emptyset - j(\mathbf{x})| = \infty$

$$\text{If } G = \emptyset, \rho(\hat{\mathcal{A}}, G) = \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in \emptyset} |h(\mathbf{x}) - \emptyset| = \infty$$

(G3) If $\rho(\hat{\mathcal{A}}, G) = 0$, then $0 = \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in G} |h(\mathbf{x}) - j(\mathbf{x})|$. It follows that $h(\mathbf{x}) = j(\mathbf{x})$, so $\hat{\mathcal{A}} \cap G \neq \emptyset$.

$$\begin{aligned} \text{(G4) Let } \rho(\hat{\mathcal{A}}, G \cup H) &= \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in G \cup H} |h(\mathbf{x}) - j(\mathbf{x})| \\ &= \sup_{h \in \hat{\mathcal{A}}} \left(\min \left\{ \sup_{j \in G} |h(\mathbf{x}) - j(\mathbf{x})|, \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in H} |h(\mathbf{x}) - j(\mathbf{x})| \right\} \right) \\ &= \min \left\{ \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in G} |h(\mathbf{x}) - j(\mathbf{x})|, \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in H} |h(\mathbf{x}) - j(\mathbf{x})| \right\} \\ &= \min\{\rho(\hat{\mathcal{A}}, G), \rho(\hat{\mathcal{A}}, H)\} \end{aligned}$$

(G5) Let $\hat{\mathcal{A}}, G \in 2^{C[a, b]}, \varepsilon, \eta \in [0, \infty]$

$$\begin{aligned} \rho(\hat{\mathcal{A}}, G) &= \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in G} |h(\mathbf{x}) - j(\mathbf{x})| \leq \sup_{h \in \hat{\mathcal{A}}} \sup_{j \in G} |h(\mathbf{x}) - j(\mathbf{x})| + \varepsilon + \eta \\ &= \rho(\hat{\mathcal{A}}^\varepsilon, G^\eta) + \varepsilon + \eta \end{aligned}$$

2. $(C[a, b], +)$ is group.

3. We prove $+: C[a, b] \oplus C[a, b] \rightarrow C[a, b]$ such that $(h, j) \mapsto h + j$ is Dh-contraction. Let $\mathbf{x} \in f(t_\varepsilon(\hat{\mathcal{A}}), t_\varepsilon(G))$, then $\mathbf{x} \in t_\varepsilon(\hat{\mathcal{A}}) + t_\varepsilon(G)$. It follows $\min\{\rho(\{\mathbf{x}\}, \hat{\mathcal{A}}), \rho(\{\mathbf{x}\}, G)\} \leq \varepsilon$,

$$\rho(\{\mathbf{x}\}, \hat{\mathcal{A}} + G) \leq \varepsilon, \text{ hence } \mathbf{x} \in t_\varepsilon(f(\hat{\mathcal{A}} + G))$$

4. Let $\mathbf{x} \in f(t_\varepsilon(\hat{\mathcal{A}}))$, then $\mathbf{x} \in (-t_\varepsilon(\hat{\mathcal{A}}))$. It follows that $\rho(\{\mathbf{x}\}, -\hat{\mathcal{A}}) \leq \varepsilon$, so that $\mathbf{x} \in (t_\varepsilon(f(\hat{\mathcal{A}})))$. Then $(C[a, b], \rho, t_\varepsilon, +)$ is proximit group.

Definition 3.5: If $(\mathbb{X}, \rho, t_\varepsilon, +)$ is proximit group and $\mathbb{Y} \subseteq \mathbb{X}$, then $\mathbb{Y}$ is said to be proximit subgroup if the following conditions are satisfied:

1. $(\mathbb{Y}, \rho, t_\varepsilon)$ is proximit space
2. $(\mathbb{Y}, +)$ is subgroup
3. $j: \mathbb{Y} \times \mathbb{Y} \rightarrow \mathbb{Y}$ such that $j(\mathfrak{x}, \mathfrak{y}) = \mathfrak{x} * \mathfrak{y}^{-1}$ is Dh-contraction.

Example 3.6: Let Z be the set of integers and if ρ is an Dh-functional defined by

$$\rho(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = \begin{cases} \infty & \hat{\mathcal{A}} = \emptyset \text{ or } \hat{\mathcal{B}} = \emptyset \\ \inf_{\mathfrak{x} \in \hat{\mathcal{A}}} \inf_{\mathfrak{y} \in \hat{\mathcal{B}}} |\mathfrak{x} - \mathfrak{y}| & \hat{\mathcal{A}} \neq \emptyset \text{ and } \hat{\mathcal{B}} \neq \emptyset \end{cases}$$

Then $(Z, \rho, t_\varepsilon, +)$ is a proximit subgroup of Q .

4. Proximit Vector Space

We present the idea of proximit vector space and show the new results.

Definition 4.1: A quintuple $(\mathbb{X}, \rho, t_\varepsilon, +, \cdot)$ is called proximit vector space such that $\mathbb{X}$ is a non-empty set with two binary operations (addition and scalar multiplication) and triple $(\mathbb{X}, \rho, t_\varepsilon)$ is Proximit space if satisfy the following conditions $\forall \mathfrak{g}, \mathfrak{g}^* \in \mathbb{X}, \vartheta, \zeta \in \text{field } E$.

1. $(\mathbb{X}, \rho, t_\varepsilon, +)$ is proximit group.
2. $\zeta \cdot \mathfrak{g} \in V$
3. $\zeta \cdot (\mathfrak{g} + \mathfrak{g}^*) = \zeta \mathfrak{g} + \zeta \mathfrak{g}^*$
4. $(\mathfrak{g} + \mathfrak{g}^*) \zeta = \mathfrak{g} \cdot \zeta + \mathfrak{g}^* \cdot \zeta$
5. $(\vartheta \cdot \zeta) \cdot \mathfrak{g} = \vartheta(\zeta \mathfrak{g})$
6. $1 \cdot \mathfrak{g} = \mathfrak{g}$

Example 4.2: Let $C[a, b]$ be set of all continuous functions on the interval $[a, b]$. Then $(C[a, b], \rho, t_\varepsilon, +, \cdot)$ is proximit vector space.

Proof:

1. $(C[a, b], \rho, t_\varepsilon, +)$ is proximit group by Example (3.4).
2. $\alpha \cdot f(\mathfrak{x}) \in C[a, b]$.
3. Since $C[a, b]$ is linear space, then $\alpha(f(\mathfrak{x}) + g(\mathfrak{x})) = \alpha f(\mathfrak{x}) + \alpha g(\mathfrak{x})$
4. $(f(\mathfrak{x}) + g(\mathfrak{x}))\alpha = f(\mathfrak{x}) \cdot \alpha + g(\mathfrak{x}) \cdot \alpha$
5. $(\vartheta \cdot \zeta) \cdot f(\mathfrak{x}) = \vartheta(\zeta f(\mathfrak{x}))$
6. $1 \cdot f(\mathfrak{x}) = f(\mathfrak{x})$.

Then $(C[a, b], \rho, t_\varepsilon, +, \cdot)$ is proximit vector space.

Definition 4.3: A subset U of proximit vector space $(\mathbb{X}, \rho, t_\varepsilon, +, \cdot)$ over a field E is said to be proximit subspace if the following conditions are holds:

1. U is subspace of vector space
2. $(U, \rho_U, t_\varepsilon)$ is proximit space.

Remark 4.4: We denoted topological space by $(\mathbb{X}, T)$ where T is a family of open sets such as associated closure operator, denoted by cl_T .

Definition 4.5: A topological space $(\mathbb{X}, T)$ that associated with a proximit space which its defined as a function $\rho_T: 2^{\mathbb{X}} \times 2^{\mathbb{X}} \rightarrow [0, \infty]$ such that

$$(\hat{\mathcal{A}}, \hat{\mathcal{B}}) \mapsto \begin{cases} 0, \hat{\mathcal{A}} \in cl_T(\hat{\mathcal{B}}) \\ \infty, \hat{\mathcal{A}} \notin cl_T(\hat{\mathcal{B}}) \end{cases}$$

Proximit space of kind $(\mathbb{X}, T, \rho_T)$ for T of $\mathbb{X}$ is said to be a topological proximit space. Also, Dh-functional of type ρ_T is said to be a topological Dh-functional.

Proposition 4.6: Suppose that $(Y, \rho, t_\varepsilon, +, \cdot)$ is proximit vector space on a field E . A topological proximit space induced by topology T_Y , denoted by (Y, T_Y) satisfy the following:

1. The map $+$: $Y \times Y \rightarrow Y$ such that $(x, y) \mapsto x + y$ is Dh-contraction
2. The map $\cdot$: $E \times Y \rightarrow Y$ such that $(\alpha, x) \mapsto \alpha x$ is Dh-contraction.

Proposition 4.7: Let (Y, T) be topological space, then a function $\rho_T: 2^Y \times 2^Y \rightarrow [0, \infty]$ defined by

$$\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = \begin{cases} 0, \hat{\mathcal{A}} \subseteq cl_T(\hat{\mathcal{B}}) \\ \infty, \hat{\mathcal{A}} \not\subseteq cl_T(\hat{\mathcal{B}}) \end{cases}$$

is Dh-functional.

Proof:

- (1) Let $\hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^Y$, if $\hat{\mathcal{A}} \subseteq cl_T(\hat{\mathcal{B}})$. Then $\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = 0 = \rho_T(\hat{\mathcal{B}}, \hat{\mathcal{A}})$.
if $\hat{\mathcal{A}} \not\subseteq cl_T(\hat{\mathcal{B}})$. Then $\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = \infty = \rho_T(\hat{\mathcal{B}}, \hat{\mathcal{A}})$.
- (2) Let $\hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^Y$, if $\hat{\mathcal{A}} \subseteq cl_T(\hat{\mathcal{B}})$ and if $\hat{\mathcal{B}} = \emptyset$. Thus $cl_T(\emptyset) = \emptyset$, so $\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = \infty$.
- (3) Let $\hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^Y$, if $\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}}) = 0$. Then $\hat{\mathcal{A}} \subseteq cl_T(\hat{\mathcal{B}})$, but $\hat{\mathcal{A}} \subseteq cl_T(\hat{\mathcal{A}})$ and $\hat{\mathcal{B}} \subseteq cl_T(\hat{\mathcal{B}})$. Thus $\hat{\mathcal{A}} \cap \hat{\mathcal{B}} \neq \emptyset$.
- (4) Let $\hat{\mathcal{A}}, \hat{\mathcal{B}}, \hat{\mathcal{C}} \in 2^Y$, Thus $cl_T(\hat{\mathcal{A}} \cup \hat{\mathcal{B}}) = c_l(\hat{\mathcal{A}}) \cup cl_T(\hat{\mathcal{B}})$, so
 $\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}} \cup \hat{\mathcal{C}}) = \min(\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}}), \rho_T(\hat{\mathcal{A}}, \hat{\mathcal{C}}))$.
- (5) Let $\hat{\mathcal{A}}, \hat{\mathcal{B}} \in 2^Y, \varepsilon, \gamma \in [0, \infty]$, Thus $\hat{\mathcal{A}}^\varepsilon = cl_T(\hat{\mathcal{A}}), \hat{\mathcal{B}}^\gamma = cl_T(\hat{\mathcal{B}}), \hat{\mathcal{A}}^\varepsilon = Y$ and $\hat{\mathcal{B}}^\gamma = Y$. So that $\rho_T(\hat{\mathcal{A}}, \hat{\mathcal{B}}) \leq \rho_T(\hat{\mathcal{A}}^\varepsilon, \hat{\mathcal{B}}^\gamma) + \varepsilon + \gamma$.

Theorem 4.8: If $(Y, \rho_Y, t_{\varepsilon_Y})$ is proximit vector space, $\mathcal{M}$ is closed proximit subspace of $(Y, \rho_Y, t_{\varepsilon_Y})$. Then $(Y/\mathcal{M}, \rho'_{Y/\mathcal{M}}, t_{\varepsilon_{Y/\mathcal{M}}})$ is proximit vector space such that $\rho'_{Y/\mathcal{M}} : 2^{Y/\mathcal{M}} \times 2^{Y/\mathcal{M}} \rightarrow [0, \infty]$ is defined by $\rho'_{Y/\mathcal{M}}(\mathcal{A}, \widehat{\mathcal{B}}) = \rho'_{Y/\mathcal{M}}(\mathcal{P} + \mathcal{A}, \mathcal{Q} + \widehat{\mathcal{B}}) = \rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{B}})$

Proof: Let $\widehat{\mathcal{A}}, \widehat{\mathcal{B}}, \widehat{\mathcal{C}} \subseteq Y$

1. $\rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{B}} + \mathcal{M}) = \rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{B}}) = \rho_Y(\widehat{\mathcal{B}}, \widehat{\mathcal{A}}) = \rho'_{Y/\mathcal{M}}(\widehat{\mathcal{B}} + \mathcal{M}, \widehat{\mathcal{A}} + \mathcal{M})$
2. If $\widehat{\mathcal{A}} = \emptyset$ or $\widehat{\mathcal{B}} = \emptyset$, then $\rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{B}} + \mathcal{M}) = \rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{B}}) = \infty$
3. If $\rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{B}} + \mathcal{M}) = 0$, then $\rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{B}}) = 0$. So that $\widehat{\mathcal{A}} \cap \widehat{\mathcal{B}} \neq \emptyset$.
4. $\rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, (\widehat{\mathcal{B}} + \mathcal{M}) \cup (\widehat{\mathcal{C}} + \mathcal{M})) = \rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, (\widehat{\mathcal{B}} \cup \widehat{\mathcal{C}}) + \mathcal{M})$,
then $\rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{B}} \cup \widehat{\mathcal{C}}) = \min\{\rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{B}}), \rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{C}})\}$
 $= \min\{\rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{B}} + \mathcal{M}), \rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{C}} + \mathcal{M})\}$
5. Let $\widehat{\mathcal{A}}, \widehat{\mathcal{B}} \in 2^{Y/\mathcal{M}}, \varepsilon, \gamma \in [0, \infty]$
 $\rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{B}} + \mathcal{M}) = \rho_Y(\widehat{\mathcal{A}}, \widehat{\mathcal{B}}) \leq \rho_Y(\widehat{\mathcal{A}}, \widehat{\widehat{\mathcal{B}}}) + \varepsilon + \gamma$
 $\leq \rho'_{Y/\mathcal{M}}(\widehat{\mathcal{A}}^\varepsilon + \mathcal{M}, \widehat{\mathcal{B}}^\gamma + \mathcal{M}) + \varepsilon + \gamma.$

Then $(Y/\mathcal{M}, \rho'_{Y/\mathcal{M}}, t_\varepsilon)$ is proximit space.

2. $(Y/\mathcal{M}, +)$ is group.
3. $+: Y/\mathcal{M} \times Y/\mathcal{M} \rightarrow Y/\mathcal{M}$ such that $(x + \mathcal{M}, y + \mathcal{M}) \mapsto x + y + \mathcal{M}$
Let $\widehat{\mathcal{A}}, \widehat{\mathcal{B}} \subseteq Y/\mathcal{M}, \varepsilon \in R^+$. Suppose that $x \in f(t_\varepsilon(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{B}} + \mathcal{M}))$, then $x \in t_\varepsilon(\widehat{\mathcal{A}} + \widehat{\mathcal{B}} + \mathcal{M})$. It follows $\rho'_{Y/\mathcal{M}}(\{x\}, \widehat{\mathcal{A}} + \widehat{\mathcal{B}} + \mathcal{M}) \leq \varepsilon$. Thus $x \in t_\varepsilon(f(\widehat{\mathcal{A}} + \mathcal{M}, \widehat{\mathcal{B}} + \mathcal{M}))$
4. Let $\widehat{\mathcal{A}} \subseteq Y/\mathcal{M}, \varepsilon \in R^+$. Suppose that $x \in f(t_{Y/\mathcal{M}}(\widehat{\mathcal{A}}))$, then $x \in (-t_{Y/\mathcal{M}}(\widehat{\mathcal{A}} + \mathcal{M}))$. Hence $\rho'_{Y/\mathcal{M}}(\{x\}, -(\widehat{\mathcal{A}} + \mathcal{M})) \leq \varepsilon_{Y/\mathcal{M}}$, so $x \in (t_\varepsilon(f(\widehat{\mathcal{A}})))$, then $(Y/\mathcal{M}, \rho'_{Y/\mathcal{M}}, t_\varepsilon, +)$ is proximit group.

The remaining condition is direct.

Definition 4.9: If (Y, ρ, t_ε) is proximit vector space and $\langle \mathfrak{X}_n \rangle$ is a sequence in (Y, ρ, t_ε) , then $\langle \mathfrak{X}_n \rangle$ is said to be convergent if $\exists \mathfrak{X} \in Y$ such that $\lim_{n \rightarrow \infty} \inf \rho(\{\mathfrak{X}_n\}, \widehat{\mathcal{B}}) = 0$ and $\lim_{n \rightarrow \infty} \sup \rho(\{\mathfrak{X}_n\}, \widehat{\mathcal{B}}) = 0$

Definition 4.10: If $(Y, \rho_Y, t_\varepsilon)$ and $(\mathcal{W}, \rho_{\mathcal{W}}, t_\varepsilon)$ are proximit spaces. We say that the function $\varphi: Y \rightarrow \mathcal{W}$ is sequentially Dh-contraction if $\lim_{n \rightarrow \infty} \varphi(t_{\varepsilon_Y}(\{\mathfrak{X}_n\})) = 0$ whenever $\lim_{n \rightarrow \infty} t_{\varepsilon_{\mathcal{W}}}(\varphi(\{\mathfrak{X}_n\})) = 0$.

Theorem 4.11: If $(Y, \rho_Y, t_\varepsilon)$ and $(\mathcal{W}, \rho_{\mathcal{W}}, t_\varepsilon)$ are proximit spaces. Then a function $\varphi: Y \rightarrow \mathcal{W}$ is Dh-contraction if and only if its sequentially Dh-contraction.

Proof: Let φ be an Dh-contraction, let $\{\mathfrak{X}_n\}$ be convergent sequence. Then $\lim_{n \rightarrow \infty} \inf \rho(\{\mathfrak{X}_n\}, \widehat{\mathcal{B}}) = 0$ and $\lim_{n \rightarrow \infty} \sup \rho(\{\mathfrak{X}_n\}, \widehat{\mathcal{B}}) = 0$, but $\varphi(t_{\varepsilon_Y}(\{\mathfrak{X}_n\})) \subseteq t_{\varepsilon_{\mathcal{W}}}(\varphi(\{\mathfrak{X}_n\}))$. Then $\lim_{n \rightarrow \infty} t_{\varepsilon_{\mathcal{W}}}(\varphi(\{\mathfrak{X}_n\})) = 0$ implies that $\lim_{n \rightarrow \infty} \varphi(t_{\varepsilon_Y}(\{\mathfrak{X}_n\})) = 0$. Thus φ is sequentially Dh-contraction.

Conversely, assume that φ is sequentially Dh-contraction. If φ is not Dh-contraction, then $\varphi(t_{\varepsilon_Y}(\{\mathfrak{X}_n\})) \not\subseteq t_{\varepsilon_{\mathcal{W}}}(\varphi(\{\mathfrak{X}_n\}))$ such that there exists $\mathfrak{f} \in \varphi(t_{\varepsilon_Y}(\{\mathfrak{X}_n\}))$, $\mathfrak{f} \notin t_{\varepsilon_{\mathcal{W}}}(\varphi(\{\mathfrak{X}_n\}))$. Since φ is sequentially Dh-contraction, then $\lim_{n \rightarrow \infty} \varphi(t_{\varepsilon_Y}(\{\mathfrak{X}_n\})) = 0$ whenever $\lim_{n \rightarrow \infty} t_{\varepsilon_{\mathcal{W}}}(\varphi(\{\mathfrak{X}_n\})) = 0$ which is contradiction.

5. Conclusion

The regularity of functionals between sets is the subject of this study. The terms proximit semigroup and proximit group, as well as other proximit space notions, are introduced here. The notion of Dh-contraction is discussed as we look at some of the results. Furthermore, we include fundamental details and illustrations. Topological proximit vector space, Dh-contraction, and proximit vector space are studied.

References

- [1] R.K. Abbas, B.Y Hussein, New Results of Completion Normed Approach Space, *Iraqi Journal of Science*, 63(5) (2022), <https://doi.org/10.24996/ij.s.2022.63.5.25>.
- [2] R.K. Abbas, B.Y Hussein, A new kind of topological vector space: Topological approach vector space, *AIP Conference Proceedings* 2386, 060008 (2022). <https://doi.org/10.1063/5.0066971>.
- [3] F. Afzal, Shah Rukh, D.Afzal, M.R.Farahani, M.Cancan, S. Ediz. A study of non-commutative Von-Neumann regular rings. *Journal of Information and Optimization Sciences*, 42(7), 1425-1436 (2021). <https://doi.org/10.1080/02522667.2021.1896650>.

- [4] Z. Ahmad, M.A. Chaudhary, A.Q. Baig, M.A. Zahid. On metric dimension of $P(n,2)*K_1$ graph. *Journal of Discrete Mathematical Sciences and Cryptography*. 24(2), 629-645 (2021). <https://doi.org/10.1080/09720529.2021.1907017>.
- [5] Z. Ahmad, M.A. Chaudhary, A.Q. Baig, M.A. Zahid. Fault-tolerant metric dimension of $P(n,2)*K_1$ graph. *Journal of Discrete Mathematical Sciences and Cryptography*. 24(2), 647-656 (2021). <https://doi.org/10.1080/09720529.2021.1899209>.
- [6] D.G. Bhimani, Contraction of functions in modulation spaces, *Integral Transforms and Special Functions*, 26 (2015). <https://doi.org/10.1080/10652469.2015.1036055>.
- [7] S. Ediz, İ. Çiftçi, M. Cancan, M.R. Farahani, On k-total distance degrees and k-total Wiener polarity index. *Journal of Information and Optimization Sciences*. 42(7), 1469-1477 (2021). <https://doi.org/10.1080/02522667.2021.1896652>.
- [8] N. Fabiano, V. Parvaneh, D. Mirković, L. Paunović, S. Radenović, On W-contractions of Jungck-Ćirić-Wardowski-type in metric spaces, *Cogent Math., Statistics*, 7 (2020). <https://doi.org/10.1080/25742558.2020.1792699>.
- [9] A.P. Farajzadeh, J. Zafarani, Equilibrium problems and variational inequalities in topological vector spaces, *Optimization*, 59, (2010). <https://doi.org/10.1080/02331930801951090>.
- [10] G. Jäger, Extensions of contractions and uniform contractions on dense subspaces, *Quaestiones Mathematicae*, 37 (2014). <https://doi.org/10.2989/16073606.2013.779984>.
- [11] G. Jäger, Quantale-valued generalizations of approach spaces and quantale-valued topological spaces, *Quaestiones Mathematicae*, 42 (2019). <https://doi.org/10.2989/16073606.2018.1516250>.
- [12] Dh. A. Kadhim, A New Type of Metric space Via Proximity Structure, In press.
- [13] E. Lowen, R. Lowen, A quasitopos containing CONV and MET as full subcategories, *Internat. J. Math. and Math. Sci.* 11, 417-433 (1988). <https://doi.org/10.1155/S016117128800050X>.
- [14] R. Lowen, Approach spaces: a common supercategory of TOP and MET, *Math. Nachr.* 141, 183-226 (1989). <https://doi.org/10.1002/mana.19891410120>.

- [15] R. Lowen, K. Robey, Completion of product of metric spaces, *Quart. J. Math. Oxford*, 43, 319-338 (1991). <https://doi.org/10.1093/qmath/43.3.319>.
- [16] R. Lowen, Approach spaces: the missing link in the topologies-uniformities-metrics, Oxford University Press, (1997).
- [17] R. Lowen, B. Windels, Approach Groups, *Journal of Math.* 30 (2000).
- [18] R. Lowen, S. Verwulgen, Approach vector spaces, *Houston J. Math* 30, 1127-1142 (2004).
- [19] R. Lowen, B. Berckmoes and J. Van Casteren, Approach theory meets probability theory, *Topology and its Applications* 158, 836-852 (2011).
- [20] R. Lowen, E. Colebunders, S. De Wachter, Intrinsic approach spaces on domains, *Topology and its Applications* 158, 2343-2355 (2011).
- [21] R. Lowen, Index Analysis: Approach Theory at Work, Springer, (2015).
- [22] M. Nadeem, S. Ahmad, M.K. Siddiqui, M.A. Ali, M.R.Farahani, A.J.M. Khalaf. On some applications related with algebraic structures through different well known graphs. *Journal of Discrete Mathematical Sciences and Cryptography*. 24(2), 451-471 (2021). <https://doi.org/10.1080/09720529.2021.1885806>.
- [23] A. Alsinai, H.M.U. Rehman, Y. Manzoor, M. Cancan, Z. Taş, M.R. Farahani, Sharp upper bounds on forgotten and SK indices of cactus graph, *Journal of Discrete Mathematical Sciences and Cryptography*. 1-22 (2022). <https://doi.org/10.1080/09720529.2022.2027605>.
- [24] M.S. Sardar, M. Alaeiyan, M.R.Farahani, M.Cancan, S. Ediz. Resistance distance in some classes of rooted product graphs obtained by Laplacian generalized inverse method. *Journal of Information and Optimization Sciences* 42(7), 1447-1467 (2021). <https://doi.org/10.1080/02522667.2021.1899210>.
- [25] M. Sioen, W. Van Den Haute, An Alternative Description of Approach Spaces Via Approach Cores, *Appl. Categor. Struct.* 24, 545-558 (2016).
- [26] W. Li, D. Zhang, Sober metric approach spaces, *Topology and its Applications* 233, 67-88 (2018). <https://doi.org/10.1016/j.topol.2017.10.019>.