

Double ZMA-transform for solving linear telegraph equation

Zainab Mohammed Alwan

Department of Mathematics

College of Science

Mustansiriyah University

Baghdad

Iraq

Abstract

Recently, a new technique to solve initial-boundary value problems (IBVP) for general linear telegraph equation (GLTE), namely Double ZMA- transform (DZMAT) introduced. This transform is an efficient and attractive technique; moreover accurate to find the exact solution by using the procedure for the transform of (LTE).

Subject Classification: 15A04, 20M20.

Keywords: Transform, Double ZMAT, General LTE, LTE, ZMAT, PDEs.

1. Introduction

Most events are modeled using partial differential equations (PDEs) in mathematical physic [1], and have confirmed and calculated the connection between the double transform, as well as their applications to differential equations by [2-4].

From [5] have established the relation between the double transform and discussed their applications. Quite recently, the general linear telegraph and partial integral-differential equations have been settled by [6].

In [1],[7-8],[19-20], a lot of problems have been explained and discussed by integral transforms. Also, [9-16] applied many problems by differential transform method. In this task, discuss some majorities new transformation to find solution of PDEs and understand procedure by discusses applications of (IBVP) for (GLTE) and confirmation the Convergence Theorem of (DZMAT).

E-mail: lionwight_2009@uomustansiriya.edu.iq

2. Definition and derivations of Double ZMA- transform

In [17],[18] ZMAT realized as:

$$\Lambda\{\wp(t)\} = \left\{ \wp(t): \exists M, \Omega_1, \Omega_2 > 0, |\wp(t)| < Me^{-\frac{t}{\Omega v}} \right\} \quad (2.1)$$

$$\mathbb{Z}_{MA}(v; s) = \Lambda\{\wp(t)\} = \frac{1}{sv} \int_0^\infty \wp(t) e^{-\frac{t}{sv}} dt \quad (2.2)$$

To find (DZMAT) for PDEs as:

Let $\wp(x, t)$ where $x, t \in \mathbb{R}^+$ be a capacity which can be represented as convergent series:

$$\mathbb{Z}_{MA}(u, v; s) = \Lambda_2\{\wp(x, t)\} = \frac{1}{s^2 uv} \int_0^\infty \int_0^\infty \wp(x, t) e^{-\left(\frac{x}{su} + \frac{t}{sv}\right)} dx dt, x, t > 0 \quad (2.3)$$

Where u, v are complex values, by (DZMAT) can be obtained the telegraph and wave equation. In the following find the 1st and 2nd- order derivatives by (DZMAT) as:

$$\Lambda_2 \left\{ \frac{\partial \wp(x, t)}{\partial x} \right\} = \frac{1}{su} \mathbb{Z}_{MA}(u, v; s) - \frac{1}{su} \mathbb{Z}_{MA}(0, v; s) \quad (2.4)$$

$$\Lambda_2 \left\{ \frac{\partial^2 \wp(x, t)}{\partial x^2} \right\} = \frac{1}{s^2 u^2} \mathbb{Z}_{MA}(u, v; s) - \frac{1}{s^2 u^2} \mathbb{Z}_{MA}(0, v; s) - \frac{1}{su} \frac{\partial \mathbb{Z}_{MA}(0, v; s)}{\partial x} \quad (2.5)$$

$$\Lambda_2 \left\{ \frac{\partial \wp(x, t)}{\partial t} \right\} = \frac{1}{sv} \mathbb{Z}_{MA}(u, v; s) - \frac{1}{sv} \mathbb{Z}_{MA}(u, 0; s) \quad (2.6)$$

$$\Lambda_2 \left\{ \frac{\partial^2 \wp(x, t)}{\partial t^2} \right\} = \frac{1}{s^2 v^2} \mathbb{Z}_{MA}(u, v; s) - \frac{1}{s^2 v^2} \mathbb{Z}_{MA}(u, 0; s) - \frac{1}{sv} \frac{\partial \mathbb{Z}_{MA}(u, 0; s)}{\partial t} \quad (2.7)$$

$$\begin{aligned} \Lambda_2 \left\{ \frac{\partial^2 \wp(x, t)}{\partial x \partial t} \right\} &= \frac{1}{s^2 uv} \wp(0, 0) - \frac{1}{sv} \mathbb{Z}_{MA}(u, 0; s) + \frac{1}{s^2 uv} \mathbb{Z}_{MA}(u, v; s) \\ &\quad - \frac{1}{s^2 uv} \mathbb{Z}_{MA}(0, v; s) \end{aligned} \quad (2.8)$$

3. Methodology

3.1 The Convergence Theorem of (DZMAT)

This action proves the convergence of (DZMAT) informed (2.3).

Theorem 3.1: If two variables' function $\wp(x, t)$ is continuous in the 1st- quadrant of the xt - plane. Where (DZMAT) realized in (2.3) converges at $s = s_0, u = u_0$ and $v = v_0$. When $s > s_0, u > u_0$ and $v > v_0$ then the integral converges, assuming that $\frac{1}{su} = \frac{1}{s_0 u_0} > 0$ and $\frac{1}{sv} = \frac{1}{s_0 v_0} > 0$.

Proof: Rewrite the integral (2.3) as:

$$\begin{aligned} \Lambda_2\{\wp(x, t)\} &= \frac{1}{s^2 uv} \int_0^\infty \int_0^\infty \wp(x, t) e^{-\left(\frac{x}{su} + \frac{t}{sv}\right)} dx dt, x, t > 0 \\ &= \frac{1}{su} \int_0^\infty e^{-\frac{x}{su}} \left[\frac{1}{sv} \int_0^\infty e^{-\frac{t}{sv}} \wp(x, t) \right] dt dx \\ &= \frac{1}{su} \int_0^\infty e^{-\frac{x}{su}} \Psi(x, t) dx \end{aligned} \quad (3.1)$$

where

$$\Psi(x, t) = \frac{1}{sv} \int_0^\infty e^{-\frac{t}{sv}} \wp(x, t) dt \quad (3.2)$$

Firstly, prove that the integral (3.2) for $s > s_0$ and $v > v_0$, where $\frac{1}{sv} = \frac{1}{s_0 v_0} > 0$, suppose that integral (3.1) converges at

$$s > s_0 \text{ and } v > v_0 \quad \mathbb{Q}(x, t) = \frac{1}{s_0 v_0} \int_0^\infty e^{-\frac{\kappa}{s_0 v_0}} \wp(x, \kappa) d\kappa, \quad 0 < t < \infty \quad (3.3)$$

$\lim_{t \rightarrow \infty} \mathbb{Q}(x, t)$ exists and $\mathbb{Q}(x, t) = 0$ since (3.2) converges at $s > s_0$ and $v > v_0$. Using the fundamental theorem of calculus, find:

$$\mathbb{Q}_t(x, t) = \frac{1}{s_0 v_0} e^{-\frac{t}{s_0 v_0}} \wp(x, \kappa) \quad (3.4)$$

Select α and β , $0 < \alpha < \beta$, then from (3.2) obtain:

$$\begin{aligned} \frac{1}{sv} \int_0^\infty e^{-\frac{t}{sv}} \wp(x, t) dt &= \frac{1}{sv} \int_\alpha^\beta e^{-\frac{t}{sv} e^{\frac{t}{s_0 v_0}}} \mathbb{Q}_t(x, t) dt \\ &= \frac{1}{sv} \int_\alpha^\beta e^{-\left(\frac{1}{sv} - \frac{1}{s_0 v_0}\right)t} \mathbb{Q}_t(x, t) dt \end{aligned} \quad (3.5)$$

by the integration by parts for (3.5) get:

$$\begin{aligned} \frac{1}{sv} \left[e^{-\left(\frac{1}{sv} - \frac{1}{s_0 v_0}\right)\alpha} \mathbb{Q}(x, \alpha) - e^{-\left(\frac{1}{sv} - \frac{1}{s_0 v_0}\right)\beta} \mathbb{Q}(x, \beta) \right] \\ + \left(\frac{1}{sv} - \frac{1}{s_0 v_0} \right) \int_0^\beta e^{-\left(\frac{1}{sv} - \frac{1}{s_0 v_0}\right)t} \mathbb{Q}(x, t) dt \end{aligned} \quad (3.6)$$

when $\alpha \rightarrow 0$ from $\left(\frac{1}{sv} = \frac{1}{s_0 v_0} > 0\right)$ the 1st-term on (3.6) converges to zero, lastly, find:

$$\frac{1}{sv} \int_0^\infty e^{-\frac{t}{sv}} \wp(x, t) dt = \left(\frac{1}{sv} - \frac{1}{s_0 v_0} \right) \int_0^\infty e^{-\left(\frac{1}{sv} - \frac{1}{s_0 v_0}\right)t} \mathbb{Q}(x, t) dt \quad (3.7)$$

It is clear that the integral on the right converges, then the integral (3.2) converges for $s > s_0$ and $v > v_0$ where $\frac{s}{s_0} = \frac{v}{v_0} > 0$.

In the same way, it proves the convergence of the integral $\frac{1}{su} \int_0^\infty e^{-\frac{x}{su}} \Psi(x, t) dx$ $s > s_0, u > u_0$ with $\frac{s}{s_0} = \frac{u}{u_0} > 0$. Thus, the integral on the right-hand side of (3.1) converges for $s > s_0, u > u_0$ and $v > v_0$ with assumption $\frac{s}{s_0} = \frac{u}{u_0} > 0$ and $\frac{s}{s_0} = \frac{v}{v_0} > 0$. Therefore, the (DZMAT) transform defined by (2.3) converges, and the proof of the theorem is then complete.

4. Procedure

In this portion, applied (DZMAT) to solve (GLTE) as:

- Make the (DZMAT) for PDEs.
- Find the single ZMAT of conditions.
- Substitute the 2nd step in the 1st step and solve the resulting equation. I, lastly applied the double inverse of ZMA-transform to obtain the general solution.

5. Applications

This work established some applications to understand the procedure of (DZMAT):

Application 5.1: Look at the (LTE).

$$U_{xx} = U_{tt} + U_t + U \quad (5.1)$$

$$\text{with the frontier condition } U(0, t) = e^{-t}, U_x(0, t) = e^{-t} \quad (5.2)$$

$$\text{And the incipient conditions } U(x, 0) = e^x, U_t(x, 0) = -e^x \quad (5.3)$$

$$\text{The careful solution is } U(x, t) = e^{x-t} \quad (5.4)$$

Solution: By (DZMAT) of (5.1), find:

$$\begin{aligned} & \frac{1}{s^2 u^2} \mathbb{Z}_{MA}(u, v; s) - \frac{1}{s^2 u^2} \mathbb{Z}_{MA}(0, v; s) - \frac{1}{su} \frac{\partial \mathbb{Z}_{MA}(0, v; s)}{\partial x} \\ &= \frac{1}{s^2 v^2} \mathbb{Z}_{MA}(u, v; s) - \frac{1}{sv} \frac{\partial \mathbb{Z}_{MA}(u, 0; s)}{\partial t} - \frac{1}{s^2 v^2} \mathbb{Z}_{MA}(u, 0; s) \\ &+ \frac{1}{sv} \mathbb{Z}_{MA}(u, v; s) - \frac{1}{sv} \mathbb{Z}_{MA}(u, 0; s) + \mathbb{Z}_{MA}(u, v; s) \end{aligned} \quad (5.5)$$

By (ZMAT) of (5.2) and (5.3) obtain:

$$\mathbb{Z}_{MA}(0, v; s) = \frac{1}{1+sv} \quad (5.6)$$

$$\frac{\partial \mathbb{Z}_{MA}(0, v; s)}{\partial x} = \frac{1}{1+sv} \quad (5.7)$$

$$\mathbb{Z}_{MA}(u, 0; s) = \frac{1}{1-su} \quad (5.8)$$

$$\frac{\partial \mathbb{Z}_{MA}(u, 0; s)}{\partial t} = -\frac{1}{1-su} \quad (5.9)$$

Now, substituting (5.6)-(5.9) in (5.5)

$$\left[\frac{1}{s^2 u^2} - \frac{1}{s^2 v^2} - \frac{1}{sv} - 1 \right] \mathbb{Z}_{MA}(u, v; s) = \frac{1}{1+sv} \left[\frac{1}{s^2 u^2} + \frac{1}{su} \right] - \frac{1}{s^2 v^2} \left[\frac{1}{1-su} \right] \quad (5.10)$$

$$\Rightarrow \mathbb{Z}_{MA}(u, v; s) = \frac{1}{(1-su)(1+sv)} \quad (5.11)$$

Now, take the double inverse of ZMA- transform for (5.11) and get:

$$U(x, t) = e^x e^{-t} = e^{x-t} \quad (5.12)$$

Application 5.2: Compose the (LTE).

$$U_{xx} = U_{tt} + U_t - U \quad (5.13)$$

$$\text{with the frontier conditions } U(0, t) = e^{-2t}, U_x(0, t) = e^{-2t} \quad (5.14)$$

$$\text{And the incipient conditions } U(x, 0) = e^x, U_t(x, 0) = -2e^x \quad (5.15)$$

$$\text{The careful solution is } U(x, t) = e^{x-2t} \quad (5.16)$$

Solution: By (DZMAT) of (5.13) obtain:

$$\begin{aligned}
& \frac{1}{s^2 u^2} \mathbb{Z}_{\text{MA}}(u, v; s) - \frac{1}{s^2 u^2} \mathbb{Z}_{\text{MA}}(0, v; s) - \frac{1}{su} \frac{\partial \mathbb{Z}_{\text{MA}}(0, v; s)}{\partial x} \\
&= \frac{1}{s^2 v^2} \mathbb{Z}_{\text{MA}}(u, v; s) - \frac{1}{sv} \frac{\partial \mathbb{Z}_{\text{MA}}(u, 0; s)}{\partial t} - \frac{1}{s^2 v^2} \mathbb{Z}_{\text{MA}}(u, 0; s) \\
&+ \frac{1}{sv} \mathbb{Z}_{\text{MA}}(u, v; s) - \frac{1}{sv} \mathbb{Z}_{\text{MA}}(u, 0; s) - \mathbb{Z}_{\text{MA}}(u, v; s)
\end{aligned} \tag{5.17}$$

By (ZMAT) of (5.14) and (5.15), get:

$$\mathbb{Z}_{\text{MA}}(0, v; s) = \frac{1}{1+2sv} \tag{5.18}$$

$$\frac{\partial \mathbb{Z}_{\text{MA}}(0, v; s)}{\partial x} = \frac{1}{1+2sv} \tag{5.19}$$

$$\mathbb{Z}_{\text{MA}}(u, 0; s) = \frac{1}{1-su} \tag{5.20}$$

$$\frac{\partial \mathbb{Z}_{\text{MA}}(u, 0; s)}{\partial t} = -\frac{2}{1-su} \tag{5.21}$$

Now, substituting (5.18)-(5.20) in (5.17)

$$\left[\frac{s^2 v^2 - s^2 u^2 - (s^2 u^2)sv + (s^2 u^2)(s^2 v^2)}{(s^2 u^2)(s^2 v^2)} \right] \mathbb{Z}_{\text{MA}}(u, v; s) = \frac{1+su}{s^2 u^2 (1+2sv)} + \frac{-1+sv}{s^2 v^2 (1-su)} \tag{5.22}$$

$$\Rightarrow \mathbb{Z}_{\text{MA}}(u, v; s) = \frac{1}{(1-su)(1+2sv)} \tag{5.23}$$

Know, by the double inverse of ZMA- transform for (5.23), find:

$$U(x, t) = e^x e^{-2t} = e^{x-2t} \tag{5.24}$$

Application 5.3: Take the (LTE).

$$U_{tt} + aU_t + bU = c^2 U_{xx} \tag{5.25}$$

$$\text{with the frontier conditions } U(0, t) = \ell_1(t), U_z(0, t) = k_1(t) \tag{5.26}$$

$$\text{And the incipient conditions } U(x, 0) = \ell_2(t), U_t(x, 0) = k_2(t) \tag{5.27}$$

Solution: Occupy (DZMAT) of (5.25) find:

$$\begin{aligned}
& \frac{1}{s^2 v^2} \mathbb{Z}_{\text{MA}}(u, v; s) - \frac{1}{s^2 v^2} \mathbb{Z}_{\text{MA}}(u, 0; s) - \frac{1}{sv} \frac{\partial \mathbb{Z}_{\text{MA}}(u, 0; s)}{\partial t} \\
&+ \frac{a}{sv} \mathbb{Z}_{\text{MA}}(u, v; s) - \frac{a}{sv} \mathbb{Z}_{\text{MA}}(u, 0; s) + \frac{b}{sv} \mathbb{Z}_{\text{MA}}(u, v; s) \\
&= \frac{c^2}{s^2 u^2} \mathbb{Z}_{\text{MA}}(u, v; s) - \frac{c^2}{s^2 u^2} \mathbb{Z}_{\text{MA}}(0, v; s) - \frac{c^2}{sv} \frac{\partial \mathbb{Z}_{\text{MA}}(0, v; s)}{\partial x}
\end{aligned} \tag{5.28}$$

By (ZMAT) of (5.26) and (5.27) resulted:

$$\mathbb{Z}_{\text{MA}}(0, v; s) = L_1(v) \tag{5.29}$$

$$\frac{\partial \mathbb{Z}_{\text{MA}}(0, v; s)}{\partial x} = K_1(v) \tag{5.30}$$

$$\mathbb{Z}_{\text{MA}}(u, 0; s) = L_2(u) \tag{5.31}$$

$$\frac{\partial \mathbb{Z}_{\text{MA}}(u, 0; s)}{\partial t} = K_2(u) \tag{5.32}$$

Know, exchange (5.29)-(5.32) in (5.28) as:

$$\left[\frac{1}{s^2 v^2} + \frac{a}{sv} + \frac{b}{sv} - \frac{c^2}{s^2 u^2} \right] \mathbb{Z}_{MA}(uv; s) = \frac{1}{s^2 v^2} L_2(u) + \frac{1}{sv} K_2(u) + \frac{a}{sv} L_2(v) - \frac{c^2}{s^2 u^2} L_1(v) - \frac{c^2}{sv} K_1(v) \quad (5.33)$$

$$\mathbb{Z}_{MA}(u, v; s) = \frac{\frac{1}{s^2 v^2} L_2(u) + \frac{1}{sv} K_2(u) + \frac{a}{sv} L_2(v) - \frac{c^2}{s^2 u^2} L_1(v) - \frac{c^2}{sv} K_1(v)}{\left[\frac{1}{s^2 v^2} + \frac{a}{sv} + \frac{b}{sv} - \frac{c^2}{s^2 u^2} \right]} = \lambda(u, v) \quad (5.34)$$

by the double inverse of ZMA- transform for (5.23) obtain (GLTE) for (5.25):

$$U(x, t) = \Lambda_2^{-1}[\lambda(u, v)] = \mathbb{Z}_{MA}(x, t; s) \quad (5.35)$$

where

$$\Lambda\{e^{ax+bt}\} = \frac{1}{(1-asu)(1-asv)} \quad (5.36)$$

6. Conclusion

The primary goal of this research was to define a novel method called (DZMAT). After establishing that this transformation note is useful and accurate for locating the solution to PDEs, we then reviewed and proved the primary and theorem for this technique, and finally demonstrated the application of (IBVP) for (GLTE) using (DZMAT).

References

- [1] Debnath, L., & Bhatta, D. : Integral transforms and their applications. CRC press (2014).
- [2] Estrin, T. A., & Higgins, T. J. : The solution of boundary value problems by multiple Laplace transformations. *Journal of the Franklin Institute*, 252(2), 153-167 (1951).
- [3] Eltayeb, H., & Kilicman, A. : On double Sumudu transform and double Laplace transform. *Malaysian Journal of Mathematical Sciences*, 4(1), 17-30 (2010).
- [4] Kilicman, A., Eltayeb, K. Some remarks on the sumudu and Laplace transforms and applications to differential equations, *ISRN Applied Mathematics* (2012), doi:10.5402/2012/591517.
- [5] Singh, Y., Mandia, H. K. Relationship between double Laplace transform and double Mellin transform in terms of generalized hypergeometric function with applications, *International Journal of Scientific and Engineering Research* 3(5), 1- 5(2012).

- [6] Eltayeb, H., Kilicman, A. A note on double Laplace transform and telegraphic equations, *Abstract and Applied Analysis* Article ID 932578 (2013), 6 pages <http://dx.doi.org/10.1155/2013/932578>.
- [7] J. Zhang, A Sumudu based algorithm m for solving differential equations, *Comp. Sci. J. Moldova* 15(3), pp – 303-313 (2007).
- [8] Eltayeb, H., & Kilicman, A. : A note on the Sumudu transforms and differential equations. *Applied Mathematical Sciences*, 4(22), 1089-1098 (2010).
- [9] Hassan, I. A. H. : Differential transformation technique for solving higher-order initial value problems. *Applied Mathematics and Computation*, 154(2), 299-311 (2004).
- [10] Bongsoo Jang: Solving linear and nonlinear initial value problems by the projected differential transform method. Ulsan National Institute of Science and Technology (UNIST), Banyeon-ri-100, Ulsan 689-798 Korea. *Compu. Phys. Communication* (2009).
- [11] Chen, C. O. K., & Ho, S. H. : Solving partial differential equations by two-dimensional differential transform method. *Applied Mathematics and Computation*, 106(2-3), 171-179 (1999).
- [12] Ayaz, F. : Solutions of the system of differential equations by differential transform method. *Applied Mathematics and Computation*, 147(2), 547-567 (2004).
- [13] Kanglgil, F., & Yaz, F. A. : Solitary wave Solution for kdv and M kdv equations by differential transform method, chaos solutions and fractions (2008) do1: 10.1016/j. Chaos, 2(009).
- [14] Hashim, I., Noorani, M. S. M., Ahmad, R., Bakar, S. A., Ismail, E. S., & Zakaria, A. M. : Accuracy of the Adomian decomposition method applied to the Lorenz system. *Chaos, Solitons & Fractals*, 28(5), 1149-1158 (2006).
- [15] Zhou, J. K. : Differential transformation and its applications for electrical circuits (1986).
- [16] Zainab Mohammed Alwan, (ZMA)-transform method. *Journal of Interdisciplinary Mathematics* (2021), DOI: 10.1080/09720502.2021.1963520.

- [17] Alwan, Z. M. : Solution Non-Linear Partial Differential Equations By ZMA Decomposition Method. WSEAS Transactions on Mathematics, 20, 712-716 (2021).
- [18] Zahurul, A., Sazzad, B. S., Use of integration to calculate the magnetic field in a space by using mathematical software for the effective production of nano-particles, *Journal of Interdisciplinary Mathematics*, (2022) DOI:10.1080/09720502.2022.2133228.
- [19] Ridha, Z. D., & Mohommed, F. A. : Solving linear ordinary differential equations by using Shehu transform with variable coefficients. *Journal of Interdisciplinary Mathematics*, 24(8), 2391-2400 (2021).
- [20] Eiber, M. S., & Kadhem, H. H. : On Soft Pre-Compact Maps. *Wasit Journal of Computer and Mathematics Science*, 1(2) (2022).