

Pr-small R-submodules of modules and Pr-radicals

Khawlah Ahmmed *

Nuhad S. Al. Mothafar †

Department of Mathematics

College of Science

University of Baghdad

Baghdad

Iraq

Abstract

The goal of this discussion is to study the twiggged of pure-small (pr-small) sub-module of a module W as recirculation of a small sub-module, and we give some basic idiosyncrasy and instances of this kind of sub-module. Also, we give the acquaint of pure radical of a module W (pr-radical) with peculiarities.

Subject Classification: 16P70, 16S90.

Keywords: Pure small, Pure radicals.

Introduction

Throughout this paper, W will be left R -module, in [1] A proper sub-module K of a module W is Named small ($K \ll W$) if $K + H = W$ for any sub-module H of W namely $H = W$. also in [2], A module W is named hallow module if the whole part sub-module of W is a small sub-module of W . In [3] A sub-module P of a module W is named a pure sub-module of W if $P \cap IW = IP$, for whole ideal I of R . Abdul Jaleel and Yaseen in [4] evoked the twiggged of large and small sub-modules as a recirculation of the class of teeny sub-module. in this paper, we evoked the twiggged of pure small sub-modules, which is another recirculation of the class of small sub-module, where a sub-module K of an module W is named pure-small ($K \ll_{pr} W$), if $K + H = W$ then, $0 \neq H$ pure sub-module of W . This paper consists of two parts; in part 1, we give some of the important peculiarities of this module. In part 2, we give a definition of Pr-Radical and some remarks on Pr-radical modules, that is $Rad_{pr}(W) = \sum \{N : N \ll_{pr} W\}$, which is named pure-radical of W , and we present some peculiarities of it.

* E-mail: khawlahahmmed@gmail.com (Corresponding Author)

† E-mail: nuhad.salim@sc.uobaghdad.edu.iq

1. Pr-Small R-sub modules

In this section, we introduced the twiggged of a pure small R-sub module and we gave few of its peculiarities.

Definition 1.1: If W is a module proper sub- module K of W is named pure-small (pr-small) sub-module of W means $(K \ll_{pr} W)$ if $H + K = W$, then $0 \neq H$ is pure sub- module of W ; in [3] H is a pure sub- module of W if $H \cap IW = IH$ for each ideal I of R .

Noticing and Instances 1.2:

1. It is nakedly that the whole small sub- modules pr-small sub- mod-ule. Thus, in Z_4 as Z -module $\{0,2\} \ll_{pr} Z_4$ since $\{0,2\} + Z_4 = Z_4$ and Z_4 is non-zero pure sub- mod-ule, in Z_4 .
2. The covers of the point above is false in general as the subsequent show, consider $W = Z_6$, $\{0,3\}$ is pr-small in Z_6 but not small.
3. $2Z + 3Z = Z$ as Z -module, $3Z$ is not pure in Z , so $2Z$ not pr-small in Z .
4. If W is a semisimple mod-ule, then every part sub-module of W is pr-small.

We know that in [5] an R -module W is called a F -regular if each sub-module of W is pure.

In [6],[7], A module W is named a simple if $W \neq \langle 0 \rangle$, and it has no sub-modules except $\langle 0 \rangle$ and W . A module is semi simple if W is a total of simple sub-modules or every sub-module of W is directly summoned of W . [8] A sub-module H of a module W is named an e-small sub-module in W ($H \ll_e W$), if $H + K = W$, when $K \leq_e W$ implies $K = W$.

Remark 1.3: E --small sub- mod-ule, need not pr-small as the subsequent instance in Z as Z -module $2Z + 3Z = Z$, $2Z$ is e-small but not pr-small since $3Z$ is not pure.

Remark 1.4: If W is F -regular module, then every pr-small sub- module of W is a small sub- mod-ule.

In the subsequent proposition, we give properties of pr-small R-sub modules:

Proposition 1.5: If W be a module and K, H are sub-modules of W so $K \leq H \leq W$ if $H \ll_{pr} W$, then $K \ll_{pr} W$.

Proof: Suppose U be R-sub module of W so $K + U = W$ since $K \leq H$, thence $H + U = W$ we defined $H \ll_{pr} W$ implies U is a pure R-sub module of W hence $K \ll_{pr} W$.

Remark 1.6: If W is a module and $K \leq H \leq W$, if $K \ll_{pr} H$, then it is not true that $K \ll_{pr} W$, as the subsequent instance. In Z as a Z -module let $K = 4Z$, $H = 2Z$ so $4Z \leq 2Z \leq Z$, we have $4Z \ll_{pr} 2Z$ since we have only $4Z + 2Z = 2Z$ and $2Z$ is a pure R -sub module of $2Z$ in Z , but $4Z$ is Lise Pr-small in Z because $4Z + 3Z = Z$ and $3Z$ is not pure in Z .

Corollary 1.7: If W is R -module such that $H_1 \ll_{pr} W$, then $H_1 \cap H_2 \ll_{pr} W$.

Proof: $H_1 \cap H_2 \leq H_1$, since $H_1 \ll_{pr} W$ implies $H_1 \cap H_2 \ll_{pr} W$ used proposition (1.5). Similarly if $H_2 \ll_{pr} W$ we have $H_1 \cap H_2 \ll_{pr} W$.

Remark 1.8: The covers of corollary (1.7) is Lise true as the subsequent instance: In Z as Z -module $K_1 = 2Z_{12}$, $K_2 = 3Z_{12}$ so $K_1 \cap K_2 = 2Z_{12} \cap 3Z_{12} = 6Z_{12} \ll_{pr} Z_{12}$ but K_1, K_2 are Lise pr-small in Z_{12} .

Corollary 1.9: If W is an R -module and H_1, H_2 be sub-modules of W , if $H_1 + H_2 \ll_{pr} W$, then $H_1 \ll_{pr} W$ and $H_2 \ll_{pr} W$.

Proof: Since $H_1 \leq H_1 + H_2 \leq W$ and $H_1 + H_2 \ll_{pr} W$ implies $H_1 \ll_{pr} W$ used proposition (1.5). In the same way, we have $H_2 \leq H_1 + H_2 \leq W$ and $H_1 + H_2 \ll_{pr} W$ implies $H_2 \ll_{pr} W$.

Remark 1.10: The opposite of corollary (1.9) is Lise true as the subsequent instance, in Z_{12} as Z -module $2Z_{12} + 3Z_{12} = Z_{12}$ is pr-small in Z_{12} but $2Z_{12}$ is not pure in Z_{12} .

Proposition 1.11: If W is an R -module and H, L are sub-modules of W so $H \leq L \leq W$ if $H \ll_{pr} L$ and L pure of W then $H \ll_{pr} W$.

Proof: Assume P be sub-module of W such that $P + H = W$, then $(P + H) \cap L = W \cap L = L$. assist modular law $H + (P \cap L) = L$ since $H \ll_{pr} L$, then $P \cap L$ is pure in L , but L is pure in W namely $P \cap L$ pure in W used (1.1) so $H \ll_{pr} W$.

Remark 1.12: If $f: W \rightarrow K$ is an isomorphism where W and K are an R -module such that $H \ll_{pr} W$, then $f(H) \ll_{pr} K$.

Proof: Assume L is a sub-module of K so $f(H) + L = K$ since f is an isomorphism, then $f^{-1}(f(H) + L) = W$, $H + f^{-1}(L) = W$ since $H \ll_{pr} W$, then $f^{-1}(L)$ is pure in W , then $f(f^{-1}(L))$ is pure in W , but $f(f^{-1}(L)) = L$ implies $f(H) \ll_{pr} W$.

Proposition 1.13: If W is an R -module and H, L are sub-modules of W so $H \leq L \leq W$ if $\frac{L}{H} \ll_{pr} \frac{W}{H}$, then $L \ll_{pr} W$.

Proof: Assume K be a sub-module of w so $L + K = W$, since there $\frac{L}{H} \ll_{pr} \frac{W}{H}$, there exist $\frac{K}{H}$ is pure in $\frac{W}{H}$ since H R -sub module of W such that $l + H = w$, $\frac{L}{H} + \frac{K}{H} = \frac{W}{H}$ but $\frac{K}{H}$ is pure of $\frac{W}{H}$, then used [8] K is pure of W there fore $L \ll_{pr} W$.

Proposition 1.14: If W is an R -module and H, L are sub-modules of W so $H \leq L \leq W$ if $L \ll_{pr} W$ then $\frac{L}{H} \ll_{pr} \frac{W}{H}$.

Proof: Suppose $\frac{K}{H}$ be an R -sub-module of $\frac{W}{H}$, $H \leq K$ so $\frac{L}{H} + \frac{K}{H} = \frac{W}{H}$, then $\frac{L+K}{H} = \frac{W}{H}$ implies $L + K = W$, since $L \ll_{pr} W$ so K is pure in W Also $H \leq K$, then $\frac{L}{H}$ pure a R -sub-module in $\frac{W}{H}$ used [9], then $\frac{L}{H} \ll_{pr} \frac{W}{H}$.

[10] An R -module W have the pure crossing idiosyncrasy (briefly PIP) if the crossing of any two pure sub-modules is pure.

Proposition 1.15: If W is an R -module with (PIP) and K_1, K_2 be sub-modules of W , then $K_1 \ll_{pr} W$ and $K_2 \ll_{pr} W$ iff $K_1 + K_2 \ll_{pr} W$.

Proof: $\Rightarrow$) $K_1 \ll_{pr} W$ and $K_2 \ll_{pr} W$ then $K_1 + K_2 \ll_{pr} W$. Suppose H be an R -sub-module of W such that $(K_1 + K_2) + H = W$ Since $K_1 \ll_{pr} W$ and $K_2 \ll_{pr} W$ then $K_2 + H$ pure in W and $H_1 + K$ pure in W , Since W has the (PIP) idiosyncrasy $(K_1 + H) \cap (K_2 + H)$ is pure in W , But $(K_1 + H) \cap (K_2 + H) = H$, namely H is a pure R -sub module of W , then $K_1 + K_2 \ll_{pr} W$.

If $K_1 + K_2 \ll_{pr} W$, then $K_1 \ll_{pr} W$ and $K_2 \ll_{pr} W$, Since $K_1 \leq K_1 \oplus K_2 \leq W$ and $K_1 \oplus K_2 \ll_{pr} W$, then used Remark (1.4); we have $k_1 \ll_{pr} w$, likeness we prove that $K_2 \ll_{pr} W$.

2. Pure Radicals (Pr-Radicals)

In [9] the Jacobson radical of a module W name $\text{Rad}(W)$ is the total of all small R -sub-modules of W .

Definition 2.1: If W is a module, the total of all pr-small sub-module of W name pure-radical (pr-radical) of W signify $\text{Rad}_{pr}(W) = \sum \{H : H \ll_{pr} W\}$.

Remark 2.2: $\text{Rad}_{pr}(W) \leq W$.

Proof: Let $H_i \ll_{pr} W$, $i \in I$ then $\bigoplus H_i \in I \ll_{pr} W$ by Remark (1.4) implies $\text{Rad}_{pr}(W) \leq W$.

Instance and Remark 2.3:

1. $\text{Rad}_{\text{pr}} Z_4 = \{\bar{0}, \bar{2}\}$.
2. $\text{Rad}_{\text{pr}} Z_6 = \{\bar{0}, \bar{3}\} + \{\bar{0}, \bar{2}, \bar{4}\} = Z_6$.
3. Every radical of R-mod-ule W contains in pr-radical of W i. e. $\text{Rad}(W) \leq \text{Rad}_{\text{pr}}(W)$.

Proof: Suppose $H \in \text{Rad}(W)$ implies $H \in \text{Rad}_{\text{pr}} W$ since every small is pr-small, then $\text{Rad}(W) \leq \text{Rad}_{\text{pr}} W$.

For instance: $\text{Rad}_{\text{pr}}(Z_6) = Z_6$, $\text{Rad } Z_6 = \{\bar{0}\}$ clearly, $\text{Rad}(Z_6) \leq \text{Rad}_{\text{pr}}(Z_6)$.

Theorem 2.4: Suppose W, H be R-modules and $\varphi: W \rightarrow H$ be an isomorphism, then $\varphi(\text{Rad}_{\text{pr}}(W)) \leq (\text{Rad}_{\text{pr}}(H))$.

Proof: Suppose $A \ll_{\text{pr}} W$, then by proposition (1.10), $\varphi(A) \ll_{\text{pr}} H$ hence $\varphi(A) \leq \text{Rad}_{\text{pr}}(H) \forall A \ll_{\text{pr}} W$, then $\varphi(\text{Rad}_{\text{pr}}(W)) \leq (\text{Rad}_{\text{pr}}(H))$.

Corollary 2.5: If W is an R-module and K be a R-sub- module of W , then $\text{Rad}_{\text{pr}}(K) \leq \text{Rad}_{\text{pr}}(W)$.

Proof: Suppose H be an R-sub module of $\text{Rad}_{\text{pr}}(K)$ such that $H \ll_{\text{pr}} K$ since $K \ll_{\text{pr}} W$ use proposition (1.5), we get $H \ll_{\text{pr}} W$ hence K be a pr-small R-sub module of W so K R-sub- module of $\text{Rad}_{\text{pr}}(W)$, then $\text{Rad}_{\text{pr}}(K) \leq \text{Rad}_{\text{pr}}(W)$.

Proposition 2.6: If $\varphi: W \rightarrow K$ is an isomorphism where W and K be R-modules, then $\varphi^{-1}(\text{Rad}_{\text{pr}}(K)) \leq \text{Rad}_{\text{pr}}(W)$.

Proof: Suppose $H \ll_{\text{pr}} K$, then used proposition (1.12), $\varphi^{-1}(H) \ll_{\text{pr}} W$ hence $\varphi^{-1}(H) \subseteq \text{Rad}_{\text{pr}}(W) \forall H \ll_{\text{pr}} K$ thus $\varphi^{-1}(\text{Rad}_{\text{pr}}(K)) \subseteq \text{Rad}_{\text{pr}}(W)$.

3. Conclusion

In this paper, we generalized small submodules as pure-small submodules and radical as pure-radical modules, and we explained:

1. If W be an R-module and K, H are R-sub- modules of W so $K \leq H \leq W$ if $H \ll_{\text{pr}} W$, then $K \ll_{\text{pr}} W$.
2. If W is an F-regular mod-ule, then every pr-small R-sub- module of W is a small R-sub- mod-ule.
3. If W is a semisimple module, then each part R-sub- module of W is pr-small.
4. If W is an R-module and K be an R-sub- module of W , then $\text{Rad}_{\text{pr}}(K) \leq \text{Rad}_{\text{pr}}(W) \cap K$.

References

- [1] Wisam A. Ali & Nuhad S. Al. Mothafar Small prime and quasi-small prime modules, *Journal of Discrete Mathematical Sciences and Cryptography*, 24:7, 1967-1971 (2021).
- [2] Abduljaleel, A. A., & Yaseen, S. M., On Large-Small submodule and Large-Hollow module. In *Journal of Physics: Conference Series*, Vol. 1818, No. 1, p. 012214, (2021, March). IOP Publishing.
- [3] Artin, E., Algebraic numbers and algebraic functions, Vol. 358 (2005). *American Mathematical Soc.*
- [4] Fieldhous. D. J, " Pure Theories, math." Ann. 184, pp.1-18 (1969).
- [5] Omar M. Al-Ragab & Nuhad S. Al-Mothafar Radical semiprime submodules, *Journal of Discrete Mathematical Sciences and Cryptography*, 24:7, 1895-1899 (2021).
- [6] Kasch F. Modules and Rings," *Academic Press*, London (1982).
- [7] Israa Hassan Muslem, Layla Salman Mahmood, Some types of Retractable and Compressible Modules, thesis Pure Science, University of Baghdad (2016).
- [8] Al-Bahraany B.H," Modules with the pure intersection property", Ph.D. Thesis, college of Science, University of Baghdad (2000).
- [9] Wisbauer R, Foundation of module and ring theorem" Gordon and Breach, Philadelphia (1991).
- [10] Eiber, M. S., & Kadhem, H. H., On Soft Pre-Compact Maps. *Wasit Journal of Computer and Mathematics Science*, 1(2) (2022).
- [11] Omar M Al-Ragab & Nuhad S. Al-Mothafar. Radical semiprime submodules, *Journal of Discrete Mathematical Sciences and Cryptography*, 24:7, 1895-1899 (2021), DOI: 10.1080/09720529.2021.1957180.