

Pre measure defined on O -ring and some related concepts

Hind F. Abbas *

Directorate General of Education Salahaddin

Ministry of Education

Salahaddin

Iraq

Ibrahim S. Ahmed [†]

Department of Physics

College of Education-Tuzkhurmatu

Tikrit University

Salahaddin

Iraq

Abstract

The aim of this work is deals with to study pre measure defined on O -ring and its completeness as well as some properties of this concept have been investigated. We introduce the notion of extension for pre measure defined on O -ring and we discuss uniqueness of this notion.

Subject Classification: 00A05, 00A06.

Keywords: O -ring, Measure, Null-additive, O -field.

1. Introduction

Measure theory and integration are the most serious sides of modern analysis with topology and functional analysis. Measure is basis in probability theory. Far reach generalizations of measure are very available in quantum physics. Wang and Klir in 2009 [1] discussed many subjects in generalized measure theory where $\mathfrak{F}$ is class of subset of universal set, then $p: \mathfrak{F} \rightarrow [0, \infty]$ is called measure if $\exists \mathcal{C} \in \mathfrak{F}$ s.t $p(\mathcal{C}) < \infty$ and if $p(\bigcup_n \mathcal{C}_n) = \sum_n p(\mathcal{C}_n)$ for any

* E-mail: hind.f.abbas35386@st.tu.edu.iq (Corresponding Author)

[†] E-mail: ibrahim1992@tu.edu.iq

disjoint sets $\mathcal{C}_1, \mathcal{C}_2, \dots \in \mathfrak{F}$. The notion of outer measure on a σ -field was introduced by Ahmed [2] in 2021 which is generalization for the measure. Many authors was studied generalization of measure see, Wang, Yu and Li [3], Pap [4], Jun, Mesiar, and Pap [5] and Mohammed, Asaad, and Ahmed [6]. Ash [7] and Mohammed, Asaad, and Ahmed [8] have been studied the σ -field as family $\mathfrak{F}$ s.t $\mathcal{N} \in \mathfrak{F}$ and if $\mathcal{C}_1, \mathcal{C}_2, \dots \in \mathfrak{F}$, then $\bigcup_{j=1}^{\infty} \mathcal{C}_j \in \mathfrak{F}$ and if $\mathcal{C} \in \mathfrak{F}$, then $\mathcal{C}^c \in \mathfrak{F}$. The O -ring was studied by Halmos [9] which is weaker than of σ -field where a class $\mathfrak{F}$ is O -ring iff it is closed under the difference between any two sets and countable union [10]. The goal of this research is to discuss some concepts about a field real analysis and in particular we discuss the idea of pre measure defined on O -ring which is generalization of measure.

2. The results

Definition 2.1: If $\mathfrak{F}$ is O -ring then $p: \mathfrak{F} \rightarrow [0, \infty]$ is said to be pre measure defined on O -ring if the following hold:

1. $p(\Phi) = 0$.
2. if $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_n$ is disjoint sets in $\mathfrak{F}$, then $p(\bigcup_{j=1}^n \mathcal{C}_j) = \sum_{j=1}^n p(\mathcal{C}_j)$.
3. if $\mathcal{C}_1, \mathcal{C}_2, \dots \in \mathfrak{F}$, then $p(\bigcup_{j=1}^{\infty} \mathcal{C}_j) \leq \sum_{j=1}^{\infty} p(\mathcal{C}_j)$.

Example 2.2: Let $\mathcal{N} = \{1, 2, 3\}$ and let $\mathfrak{F} = \{\Phi, \{1\}, \{2\}, \{1, 2\}, \{2, 3, 4\}, \{1, 3, 4\}, \mathcal{N}\}$. Then $\mathfrak{F}$ is O -ring. Define $p: \mathfrak{F}_1 \rightarrow [0, \infty]$ by

$$p(\mathcal{C}) = \begin{cases} 0 & \text{if } \mathcal{C} = \Phi \\ \frac{1}{2} & \text{if } \mathcal{C} = \{1\} \text{ or } \{2\} \\ 1 & \text{o. w.} \end{cases}$$

It is easy to see that all conditions in above definition are hold, so we conclude that p is a pre measure defined on $\mathfrak{F}$.

Definition 2.3: we shall denote to pre measure space over O -ring $\mathfrak{F}$ by $(\mathcal{N}, \mathfrak{F}, p)$ where $\mathcal{N}$ is universal set and p is a pre measure defined on $\mathfrak{F}$.

Theorem 2.4: If $\mathfrak{F}$ is O -ring, then every measure defined on $\mathfrak{F}$ is pre measure defined on $\mathfrak{F}$.

Proof: Assume p is measure defined on O -ring $\mathfrak{F}$ and $\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_k$ are disjoint sets in $\mathfrak{F}$, then $p(\Phi) = 0$ and $p(\bigcup_{n=1}^k \mathcal{C}_n) = \sum_{n=1}^k p(\mathcal{C}_n)$. Let $\mathcal{C}_1, \mathcal{C}_2, \dots \in \mathfrak{F}$. Since p is measure defined on $\mathfrak{F}$, so p is a countably sub-additive and thus $p(\bigcup_{n=1}^{\infty} \mathcal{C}_n) \leq \sum_{n=1}^{\infty} p(\mathcal{C}_n)$. Therefore, p is pre measure defined on O -ring $\mathfrak{F}$.

Definition 2.5: Assume p is pre measure defined on O -ring $\mathfrak{F}$, the set $\mathcal{C}$ in $\mathfrak{F}$ is called p -null set, if $p(\mathcal{C}) = 0$.

Definition 2.6: A pre measure p on O -ring $\mathfrak{F}$ is said to be complete if each of p - null set $\mathcal{C}$ in $\mathfrak{F}$, we have $\mathcal{V} \in \mathfrak{F} \quad \forall \mathcal{V} \subset \mathcal{C}$.

Theorem 2.7: [1]. Any σ -field implies O -ring.

Definition 2.8: Let $(\mathcal{N}, \mathfrak{F}_1, p_1)$ & $(\mathcal{N}, \mathfrak{F}_2, p_2)$ are pre measure spaces over O -ring $\mathfrak{F}_1$ & $\mathfrak{F}_2$. Then we say that $(\mathcal{N}, \mathfrak{F}_2, p_2)$ is an extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$, if $\mathfrak{F}_1 \subseteq \mathfrak{F}_2$ and $p_1(\mathcal{C}) = p_2(\mathcal{C})$ for all $\mathcal{C} \in \mathfrak{F}_1$.

Example 2.9: Let $\mathcal{N} = \{1, 2, 3\}$ and let $\mathfrak{F}_1 = \{\Phi, \mathcal{N}\}$, $\mathfrak{F}_2 = \{\Phi, \{1\}, \{2, 3\}, \mathcal{N}\}$. Then we note that $\mathfrak{F}_1, \mathfrak{F}_2$ is O -ring. Define $p_1: \mathfrak{F}_1 \rightarrow [0, \infty]$ by

$$p_1(\mathcal{C}) = \begin{cases} 0 & \text{if } \mathcal{C} = \Phi \\ 1 & \text{if } \mathcal{C} = \mathcal{N} \end{cases}$$

And $p_2: \mathfrak{F}_2 \rightarrow [0, \infty]$ by:

$$p_2(\mathcal{V}) = \begin{cases} 0 & \text{if } \mathcal{V} = \Phi \\ \frac{1}{2} & \text{if } \mathcal{V} = \{1\} \text{ or } \mathcal{V} = \{2, 3\} \\ 1 & \text{if } \mathcal{V} = \mathcal{N} \end{cases}$$

Then each of p_1, p_2 is pre measure on O -ring $\mathfrak{F}_1, \mathfrak{F}_2$ respectively. Now, for any $\mathcal{C} \in \mathfrak{F}_1$, then $\mathcal{C} = \Phi$ or $\mathcal{C} = \mathcal{N}$, hence $\mathcal{C} \in \mathfrak{F}_2$, thus $\mathfrak{F}_1 \subseteq \mathfrak{F}_2$. it is clear that $p_1(\mathcal{C}) = p_2(\mathcal{C})$ for all $\mathcal{C} \in \mathfrak{F}_1$. So $(\mathcal{N}, \mathfrak{F}_2, p_2)$ is an extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$.

Definition 2.10: Let $\mathfrak{F}$ be O -ring. Then $p: \mathfrak{F} \rightarrow [0, \infty]$ is said to be null additive iff $\mathcal{C}, \mathcal{V}$ disjoint in $\mathfrak{F}$, $p(\mathcal{C}) = 0$, then $p(\mathcal{C} \cup \mathcal{V}) = p(\mathcal{V})$.

Definition 2.11: Let $\mathfrak{F}$ be O -ring. Then $p: \mathfrak{F} \rightarrow [0, \infty]$ is said to be weakly null additive iff $\mathcal{C}, \mathcal{V}$ disjoint in $\mathfrak{F}$, $p(\mathcal{C}) = p(\mathcal{V}) = 0$, then $p(\mathcal{C} \cup \mathcal{V}) = 0$.

Theorem 2.12: Let $\mathfrak{F}$ be a O -ring. Then every pre measure defined on $\mathfrak{F}$ is null additive on $\mathfrak{F}$.

Proof: Let p be a pre measure defined on O -ring $\mathfrak{F}$ and $\mathcal{C}, \mathcal{V}$ disjoint sets in $\mathfrak{F}$ s.t $p(\mathcal{C}) = 0$. Since p is a pre measure, then p is a finitely additive. So, $p(\mathcal{C} \cup \mathcal{V}) = p(\mathcal{C}) + p(\mathcal{V}) = p(\mathcal{V})$ since $p(\mathcal{C}) = 0$. Therefore, p is a null additive on $\mathfrak{F}$.

Definition 2.13: Let $\mathfrak{F}$ be a O -ring & p be a pre measure on $\mathfrak{F}$. Define m^* as $p^*(\mathcal{C}) = \inf\{\sum_{i=1}^{\infty} p(\mathcal{C}_i): \mathcal{C} \subset \bigcup_{i=1}^{\infty} \mathcal{C}_i \text{ \& } \mathcal{C}_i \in \mathfrak{F}\}$.

Proposition 2.14: Let $\mathfrak{F}$ be O -ring and p be a pre measure on $\mathfrak{F}$. Then p^* is an outer measure.

Proof: The result Follows.

Proposition 2.15: Let $\mathfrak{F}$ be O -ring and p be a pre measure on $\mathfrak{F}$. If p is σ -finite, then p^* is its unique extension to $\mathfrak{F}$ and therefore to $\sigma(\mathfrak{F})$.

Proof: Direct.

Theorem 2.16: Suppose $t \in \mathbb{R}$ and p is a pre measure on σ -ring $\mathfrak{F}$ and $\mathcal{C} \subset \mathbb{R}$. Then $p^*(t + \mathcal{C}) = p^*(\mathcal{C})$.

Proof: Suppose $I_1, I_2, \dots$ are family of sequence for open intervals and its union includes $\mathcal{C}$. then $t + I_1, t + I_2, \dots$ are collection of sequence for open intervals and its union includes $t + \mathcal{C}$, so $p^*(t + \mathcal{C}) \leq \sum_{k=1}^{\infty} p(t + I_k) = \sum_{k=1}^{\infty} p(I_k)$.

We taking infimum for the last term over every $I_1, I_2, \dots$ such that and its union includes $\mathcal{C}$, then we get $p^*(t + \mathcal{C}) \leq p^*(\mathcal{C})$. To get the inequality in the other side, since $\mathcal{C} = -t + (t + \mathcal{C})$. then stratifying the inequality by a former passage with $\mathcal{C}$ changed by $t + \mathcal{C}$ & t commuted by $-t$, we get $p^*(\mathcal{C}) = p^*(-t + (t + \mathcal{C})) \leq p^*(t + \mathcal{C})$. So, $p^*(t + \mathcal{C}) = p^*(\mathcal{C})$.

Theorem 2.17: Let $\mathfrak{F}$ be a O -ring. Then every pre measure defined on $\mathfrak{F}$ is weakly null additive defined on $\mathfrak{F}$.

Proof: Let p be a pre measure defined on O -ring $\mathfrak{F}$ and $\mathcal{C}, \mathcal{V}$ disjoint sets in $\mathfrak{F}$ s.t $p(\mathcal{C}) = p(\mathcal{V}) = 0$. Since p is pre measure, so p is a finitely additive, thus $p(\mathcal{C} \cup \mathcal{V}) = p(\mathcal{C}) + p(\mathcal{V}) = 0$. Hence, p is weakly null additive defined on $\mathfrak{F}$.

Theorem 2.18: Let $(\mathcal{N}, \mathfrak{F}_1, p_1)$ be a pre measure space over O -ring, then there exists the unique smallest complete extension $(\mathcal{N}, \mathfrak{F}_2, p_2)$ for $(\mathcal{N}, \mathfrak{F}_1, p_1)$.

Proof: Let $\mathfrak{F}_2 = \{\mathcal{C} : \mathcal{C} = W \cup M, \text{ where } W \in \mathfrak{F}_1 \text{ and } M \subseteq \text{of some } H \in \mathfrak{F}_1 \text{ with } p_1(H)=0\}$. To claim $\mathfrak{F}_2$ is O -ring, it clear $\mathfrak{F}_2$ is closed under complementation, so we get it is closed under the difference between any two sets which is belong to $\mathfrak{F}_2$. Let $\mathcal{C}_1, \mathcal{C}_2, \dots \in \mathfrak{F}_2$. Then $\mathcal{C}_i = W_i \cup M_i \forall i$ where $W_i \in \mathfrak{F}_1$ and $M_i \subseteq \text{of some } H_i \in \mathfrak{F}_1 \text{ with } p_1(H_i)=0 \forall i$ implies that $\sum_{i=1}^{\infty} p_1(H_i) = 0$. So, $\bigcup_{i=1}^{\infty} \mathcal{C}_i = \bigcup_{i=1}^{\infty} (W_i \cup M_i) = (\bigcup_{i=1}^{\infty} W_i) \cup (\bigcup_{i=1}^{\infty} M_i)$. but $W_i \in \mathfrak{F}_1 \forall i$ & $\mathfrak{F}_1$ is O -ring, so $\bigcup_{i=1}^{\infty} W_i \in \mathfrak{F}_1$. Now, $M_i \subseteq H_i$, thus $\bigcup_{i=1}^{\infty} M_i \subseteq \bigcup_{i=1}^{\infty} H_i$. Because $H_i \in \mathfrak{F}_1 \forall i$ & $\mathfrak{F}_1$ is a O -ring, so $\bigcup_{i=1}^{\infty} H_i \in \mathfrak{F}_1$. Since p_1 is pre measure defined on O -ring $\mathfrak{F}_1$, then $0 \leq p_1(\bigcup_{i=1}^{\infty} H_i) \leq \sum_{i=1}^{\infty} p_1(H_i) = 0$ which is implies that $p_1(\bigcup_{i=1}^{\infty} H_i) = 0$, hence $\bigcup_{i=1}^{\infty} \mathcal{C}_i \in \mathfrak{F}_2$. Therefore, $\mathfrak{F}_2$ is a O -ring. We now prove that p_2 is a pre measure defined on O -ring $\mathfrak{F}_2$. Let $\mathcal{C} \in \mathfrak{F}_2$, then $\mathcal{C} = W \cup M$, where $W \in \mathfrak{F}_1$ and $M \subseteq \text{of some } H \in \mathfrak{F}_1 \text{ and } p_1(H)=0$ and we define $p_2(\mathcal{C}) = p_1(W)$, since $\Phi \in \mathfrak{F}_2$, then $\Phi = \Phi \cup \Phi$ where $\Phi \in \mathfrak{F}_1$ and $\Phi \subseteq \Phi$, but p_1 is an pre measure defined on O -ring $\mathfrak{F}_1$ implies that $p_1(\Phi) = 0$, hence $p_2(\Phi) = p_1(\Phi) = 0$. Let $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_n$ are disjoint sets in $\mathfrak{F}_2$. Then $\mathcal{C}_i = W_i \cup M_i \forall i$ where W_i are disjoint sets in $\mathfrak{F}_1$ and $M_i \subseteq \text{of some } H_i \in \mathfrak{F}_1 \text{ with } p_1(H_i)=0$ for all $i=1, 2, \dots, n$. Now, $\bigcup_{i=1}^n \mathcal{C}_i = (\bigcup_{i=1}^n W_i) \cup (\bigcup_{i=1}^n M_i) \cup \bigcup_{i=1}^n \mathcal{C}_i \in \mathfrak{F}_2$ & $\bigcup_{i=1}^n W_i \in \mathfrak{F}_1$ & $\bigcup_{i=1}^n M_i \subseteq \bigcup_{i=1}^n H_i$ & $\bigcup_{i=1}^n H_i \in \mathfrak{F}_1$ and $p_1(\bigcup_{i=1}^n H_i) = 0$. Then by

definition of p_2 we have $p_2(\bigcup_{i=1}^n C_i) = p_1(\bigcup_{i=1}^n W_i)$. Since $W_1, W_2, \dots, W_n$ are disjoint sets in $\mathfrak{F}_1$ and p_1 is a pre measure defined on O -ring $\mathfrak{F}_1$, then we get $p_2(\bigcup_{i=1}^n C_i) = p_1(\bigcup_{i=1}^n W_i) = \sum_{i=1}^n p_1(W_i) = \sum_{i=1}^n p_2(C_i)$. Let $C_1, C_2, \dots \in \mathfrak{F}_2$. so as above $\bigcup_{i=1}^\infty C_i = (\bigcup_{i=1}^\infty W_i) \cup (\bigcup_{i=1}^\infty M_i)$ and $\bigcup_{i=1}^\infty C_i \in \mathfrak{F}_2$ and $\bigcup_{i=1}^\infty W_i \in \mathfrak{F}_1$ and $\bigcup_{i=1}^\infty M_i \subseteq \bigcup_{i=1}^\infty H_i$ and $\bigcup_{i=1}^\infty H_i \in \mathfrak{F}_1$ and $p_1(\bigcup_{i=1}^\infty H_i) = 0$. Then by definition of p_2 we have $p_2(\bigcup_{i=1}^\infty C_i) = p_1(\bigcup_{i=1}^\infty W_i)$. Since $W_1, W_2, \dots \in \mathfrak{F}_1$ and p_1 is a pre measure defined on O -ring $\mathfrak{F}_1$, then we get $p_2(\bigcup_{i=1}^\infty C_i) = p_1(\bigcup_{i=1}^\infty W_i) \leq \sum_{i=1}^\infty p_1(W_i) = \sum_{i=1}^\infty p_2(C_i)$. So p_2 is a pre measure defined on O -ring $\mathfrak{F}_2$.

To claim p_2 is a complete pre measure defined on O -ring $\mathfrak{F}_2$, let $C \in \mathfrak{F}_2$ such that $p_2(C) = 0$ and let $S \subseteq C$. Then $C = W \cup M$, where $W \in \mathfrak{F}_1$ and $M \subseteq$ of some $H \in \mathfrak{F}_1$ and $p_1(H) = 0$. From definition of p_2 , we get $p_2(C) = p_1(W) = 0$. Put $S = K \cup T$, when $K = \Phi$ & $T \subseteq Q$ & $Q = W \cup H$, so $K = \Phi \in \mathfrak{F}_1$. Since $\mathfrak{F}_1$ is a O -ring and $W, H \in \mathfrak{F}_1$, then $W \cup H \in \mathfrak{F}_1$, hence $Q \in \mathfrak{F}_1$ and $p_1(Q) = p_1(W \cup H) \leq p_1(W) + p_1(H) = 0$, thus $p_1(Q) = 0$ which implies that $S \in \mathfrak{F}_2$ and p_2 is complete. To claim $(\mathcal{N}, \mathfrak{F}_2, p_2)$ is an extension for $(\mathcal{N}, \mathfrak{F}_1, p_1)$. Assume $W \in \mathfrak{F}_1$. So, $W = W \cup \Phi$ & $\Phi \subseteq \Phi \in \mathfrak{F}_1$ & $p_1(\Phi) = 0$, hence $W \in \mathfrak{F}_2$, therefore $\mathfrak{F}_1 \subseteq \mathfrak{F}_2$. Now, from definition of p_2 , we get $p_2(W) = p_1(W)$ for any $W \in \mathfrak{F}_1$. Therefore $(\mathcal{N}, \mathfrak{F}_2, p_2)$ is an extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$.

Let $(\mathcal{N}, \mathfrak{F}_3, p_3)$ be another complete extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$ and we prove that $(\mathcal{N}, \mathfrak{F}_3, p_3)$ is an extension of $(\mathcal{N}, \mathfrak{F}_2, p_2)$. Let $C \in \mathfrak{F}_2$. Then $C = W \cup M$ for $W \in \mathfrak{F}_1$ and $M \subseteq$ of some $H \in \mathfrak{F}_1$ and $p_1(H) = 0$. Since $(\mathcal{N}, \mathfrak{F}_3, p_3)$ is an extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$, then $\mathfrak{F}_1 \subseteq \mathfrak{F}_3$, hence $W, H \in \mathfrak{F}_3$ and $p_3(H) = p_1(H) = 0$. Now, $H \in \mathfrak{F}_3$ and $p_3(H) = 0$ and $M \subseteq$ of some H and p_3 is complete, then $C \in \mathfrak{F}_3$. Since $W, M \in \mathfrak{F}_3$ and $\mathfrak{F}_3$ is a O -ring, then $W \cup M \in \mathfrak{F}_3$, but $C = W \cup M$, hence $C \in \mathfrak{F}_3$ and $\mathfrak{F}_2 \subseteq \mathfrak{F}_3$. Now, since p_3 is pre measure defined on O -ring $\mathfrak{F}_3$ and $W \subseteq W \cup M = C$, then $p_3(W) \leq p_3(C)$. Since $M \subseteq H$, then $0 \leq p_3(M) \leq p_3(H) = 0$, hence $p_3(C) = 0$. But $p_3(U) = p_3(W \cup M) \leq p_3(W) + p_3(N) = p_3(W)$. So, $p_3(C) = p_3(W)$, but p_3 is complete and $W \in \mathfrak{F}_1$ implies that $p_3(W) = p_1(W)$ and hence $p_3(C) = p_1(W)$. Now from definition of p_2 we get, $p_1(A) = p_2(C)$. Therefore $p_3(C) = p_2(C)$. Which implies, $(\mathcal{N}, \mathfrak{F}_3, p_3)$ is an extension of $(\mathcal{N}, \mathfrak{F}_2, p_2)$. Hence $(\mathcal{N}, \mathfrak{F}_2, p_2)$ is smallest complete extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$. Now, assume $(\mathcal{N}, \mathfrak{F}_*, p_*)$ & $(\mathcal{N}, \mathfrak{F}_\#, p_\#)$ are two smallest complete extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$, now $(\mathcal{N}, \mathfrak{F}_*, p_*)$ is a smallest complete extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$ and $(\mathcal{N}, \mathfrak{F}_\#, p_\#)$ is a complete extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$, then $(\mathcal{N}, \mathfrak{F}_\#, p_\#)$ is a complete extension of $(\mathcal{N}, \mathfrak{F}_*, p_*)$ and by same way we have, $(\mathcal{N}, \mathfrak{F}_*, p_*)$ is a complete extension of $(\mathcal{N}, \mathfrak{F}_\#, p_\#)$. Hence $(\mathcal{N}, \mathfrak{F}_*, p_*) = (\mathcal{N}, \mathfrak{F}_\#, p_\#)$. So, there is a unique smallest complete extension of $(\mathcal{N}, \mathfrak{F}_1, p_1)$.

Conclusion

The main idea of our paper is to allocate to real valued function defined on the generalization for σ -field which is O -ring and it's our domain. Our aims in

this article are to show that, the concept of pre measure defined on O -ring is a generalization of the measure defined on O -ring while each of null additive and weakly null additive defined on O -ring are generalization of the pre measure defined on O -ring as well as any pre measure defined on a O -ring having unique complete extension.

References

- [1] Z. Wang and G. Klir, *Generalized Measure Theory*, 1st ed. New York: Springer Science and Business Media, LLC (2009).
- [2] I. S. Ahmed, S. H. Asaad, and H. H. Ebrahim, "Some new properties of an outer measure on a σ -field," *J. Interdiscip. Math.*, vol. 24, no. 4, pp. 947–952 (2021), doi: 10.1080/09720502.2021.1884386.
- [3] P. Wang, M. Yu, and J. Li, "Monotone measures defined by Pan-integral," *Adv. Pure Math.*, vol. 8, no. 6, pp. 535–547 (2018), doi: 10.4236/apm.2018.86031.
- [4] E. Pap, "Extension of null-additive set functions on algebra of subsets," *Novi Sad J. Math.*, vol. 31, no. 2, pp. 9–13 (2001).
- [5] L. Jun, R. Mesiar, and E. Pap, "Atoms of weakly null-additive monotone measures and integrals," *Inf. Sci. (Ny)*, vol. 257, pp. 183–192 (2014), doi: 10.1016/j.ins.2013.09.013.
- [6] A. S. Mohammed, S. H. Asaad, and I. S. Ahmed, "New properties of strongly regular left almost semi-group," *J. Discret. Math. Sci. Cryptogr.*, vol. 25, no. 2, pp. 615–622 (2022), doi: 10.1080/09720529.2021.1982491.
- [7] R. B. Ash, *Real Analysis and Probability*, 1st ed. London: Academic Press, Inc. (1972).
- [8] M. H. Alobaidi, I. S. Ahmed, and S. H. Asaad, "Study of the generalized P-contractions on Banach spaces and uniqueness for stability fixed points," *Int. J. Nonlinear Anal. Appl.*, vol. 12, no. 2, pp. 1903–1908 (2021), doi: 10.22075/IJNAA.2021.5324.
- [9] P. R. Halmos, *Measure Theory*, 1st ed. New York: Springer-Verlag (1974).
- [10] M. M. Abbood, H. H. Ebrahim, A. Al-Fayadh, and A. J. Obaid, "Measure defined on Γ -algebra and some of their generalizations," *J. Interdiscip. Math.*, vol. 26, no. 5, pp. 821–827 (2023), doi: 10.47974/JIM-1502.

Received May, 2024