

Interdisciplinary applications of mathematics in healthcare systems

Vasundhara V. Ghorpade *

Department of Community Medicine

Krishna Institute of Medical Sciences

Krishna Vishwa Vidyapeeth (Deemed to be University)

Karad 415539

Maharashtra

India

Suhas Gupta [†]

Centre for Research Impact and Outcome (CRIO)

Chitkara University

Rajpura 140401

Punjab

India

Chinmay Chandrakar [§]

Department of Computer Science and Engineering

Rungta College of Engineering and Technology

Rungta International Skills University

Bhilai 490024

Chhattisgarh

India

Subramanian Karthick [‡]

Department of Computer Engineering

Vishwakarma Institute of Technology

Pune 411037

Maharashtra

India

* E-mail: drvasundharaghorpade@gmail.com (Corresponding Author)

[†] E-mail: suhas.gupta.orp@chitkara.edu.in

[§] E-mail: chinmay@rungta.org

[‡] E-mail: karthick.sb@vit.edu

Vinay Shivajirao Nalawade[@]

Department of Artificial Intelligence & Data Science Engineering

S B Patil College of Engineering

Indapur 413106

Maharashtra

India

Kumel K. Nagori[#]

Faculty of Engineering

Gokul Global University

Sidhpur 384151

Gujarat

India

Abstract

In the contemporary healthcare landscape, mathematics is integral, offering systematic solutions and frameworks for modeling biological processes, analyzing clinical data, optimizing treatment strategies, and informing medical decision-making. This study looks at interdisciplinary uses of mathematical tools in healthcare systems with particular emphasis on mathematical modelling, differential equations, dynamical systems, physiological modelling, pharmacokinetics and healthcare data analytics. Mathematical models are used to capture cardiovascular and physiological processes, and used to quantitatively analyse their complex interactions and disease progression. Differential equations can be used in simulation of time-dependent clinical condition, and pharmacokinetic modeling is helpful in estimating the dosages of effective drugs and planning a therapy. In addition, mathematical optimization, probability and statistical analysis are used to improve the management of resources in healthcare and clinical decision making.

Subject Classification: 12H20.

Keywords: *Mathematical modeling, Healthcare systems, Differential equations, Physiological modeling, Pharmacokinetics, Healthcare data analytics.*

I. Introduction

The complexity of healthcare systems has been on the rise as a result of rising population, new development of diseases, technological advancements, and the continuous generation of clinical data. To enable quantitative representation, prediction, optimization and evaluation of healthcare processes in a systematic way, mathematics offers tools to tackle these complexities [1, 2, 3]. It has several applications across different disciplines such as disease modelling, physiological analysis, medical imaging, pharmacokinetics, epidemiology, healthcare resource allocation

[@] E-mail: vinaynalawade2007@gmail.com

[#] E-mail: exam@gokuluniversity.ac.in

and clinical decision making. Biological and medical phenomena can be described as mathematical models, and these models can be analyzed to gain understanding of the relationships between important clinical variables [4, 5]. Differential equations and dynamical systems can be used in a particularly valuable way for modelling physiological phenomena that change over time, such as the dynamics of the cardiovascular system, disease progression, reaction to treatment, and concentration of drugs in the body. The analysis of uncertainty, patient variability, diagnostic outcomes and clinical evidence are all supported by probability and statistics [6, 7]. Optimization techniques help in the efficient scheduling of hospital resources, resource utilization, treatment planning and allocation of medical facilities [8, 9]. Additionally, mathematical methods coupled with data analysis in the healthcare domain can be used to extract meaningful patterns from various medical data such as electronic health records (EHRs), lab values, physiological indicators, and population health data sets [10]. The merging of mathematics with medicine, computer science, engineering and data science has enhanced the predictive and personalized medicine. Mathematical models can help clinicians assess the different treatment options, the risks involved and make better-informed decisions based on evidence [11, 12].

II. Mathematical Foundations for Healthcare Systems

From a mathematical perspective, the processes of healthcare, complex interactions, clinical outcomes and medical interventions and operations in health care can be mathematically represented, analysed and explained, and the systems can be mathematically evaluated [13].

A. Mathematical Modeling and Simulation

Mathematical modeling and simulation can be used to represent the biological or clinical process in a computational system, allowing the investigator to explore the system behavior, assess treatment options, make predictions and minimize experimental limitations effectively [14].

$$Y(t) = f(X(t), \theta, t) \quad (1)$$

This can be done mathematically as seen in equation (1) where the outcomes of healthcare are expressed as a function of the system variables, model parameters and time, which can be used to represent complex clinical processes and interactions flexibly.

$$E = \sqrt{\left[\left(\frac{1}{N}\right) \sum_{i=1}^N (Y_i - \hat{Y}_i)^2\right]} \quad (2)$$

The prediction error in equation (2) quantifies the differences between observed and predicted outcomes of healthcare, and is used to provide a quantitative measure of the general predictive performance of the model.

$$R(t) = R^0 e^{-kt} + \sqrt{[\alpha X(t)]} \quad (3)$$

The physiological response is modeled by an exponential decay and input-dependent response in equation (3) for modeling temporal changes and nonlinear effects of clinical or biological variables in healthcare systems.

$$X_{n+1} = X_n + \Delta t f(X_n, U_n) \quad (4)$$

The states of the healthcare system are changed over a set of discrete time intervals based on current conditions, external inputs and dynamic functions, allowing simulation of a sequence of evolutions of clinical processes.

$$J = \sqrt{\left[\sum_{i=1}^n w_{i(O_i - T_i)}^2 \right]} \quad (5)$$

Finally, equation (5) is a way to measure how far the observed health outcome of treatment deviates from the desired outcome, with weights assigned to each deviation, which can be used to test the performance of the treatment in reducing clinical errors, and to help optimize the treatment.

$$A = 1 - \sqrt{\frac{\sum_{i=1}^N (Y_i - \hat{Y}_i)^2}{\sum_{i=1}^N (Y_i - \bar{Y})^2}} \quad (6)$$

The normalized prediction deviations against observed variability in Eq. (6) indicate the overall model accuracy and predictive agreement with observed variability, and quantify comprehensively the reliability of the healthcare model.

B. Differential Equations and Dynamical Systems

Modeling temporal changes of physiological variables, disease progression and drug responses, as well as stability analysis, prediction and understanding of complex healthcare processes, through differential equations and dynamical systems.

$$\frac{dx(t)}{dt} = f(x(t), u(t), t) \quad (7)$$

The temporal evolution of the variables of the healthcare system are described in equation (7) according to the current state, external inputs and the time, thus offering a flexible framework for clinical system modelling.

$$\frac{dP}{dt} = rP \left(1 - \frac{P}{K} \right) \quad (8)$$

$$\frac{dD}{dt} = \beta D \left(1 - \frac{D}{K} \right) - \gamma D \quad (9)$$

The Equation (8) models the physiological growth when limited capacity exists-growth rate will decrease gradually closer to carrying capacity, which is suitable to simulate the behavior of biological populations over time. Equation (9) incorporates the growth and recovery

effects and can be used to quantify the severity, progression and therapy response of disease at any time with high accuracy.

$$\frac{dx}{dt} = \alpha\sqrt{x} - \beta x \quad (10)$$

The growth and decay process shown in Equation (10) involves nonlinear physiological changes, which depend on the roots, and proportional decay, due to the changing healthcare environment, that accurately reflects the complex biological responses and equilibrium behavior.

$$\frac{dx}{dt} = \alpha x - \beta xy \quad (11)$$

$$\frac{dy}{dt} = \delta xy - \gamma y \quad (12)$$

The changes in the first interacting healthcare variable due to intrinsic growth and interaction effects is represented by the term of equation (11), which shows the dynamics of coupled biological components as a function of their interactions. The equation (12) encodes the temporal evolution of the second type of healthcare variable, which is modeled by an interaction-driven growth and natural decay, thus allowing to analyze quantitatively the interaction between two healthcare processes related to physiology or to diseases.

III. Mathematical Modeling of Disease and Clinical Processes

Disease and clinical process modeling uses medical mechanisms to create mathematical relationships that can be quantitatively studied to learn about how the disease progresses, how patients respond, the effectiveness of the interventions, and what the potential clinical outcomes are with the disease.

A. Cardiovascular and Physiological System Modeling

Cardiovascular and physiological models mathematically describe the blood flow, heart behaviour, pressure fluctuations and interactions between organs, allowing for the assessment, simulation, diagnosis and individual treatment planning of patients with diseases.

As shown in Figure 1, patient data, mathematical modeling, simulation, validation and feedback are integrated to aid in cardiovascular analysis and healthcare decision-making processes.

$$Q = \frac{\Delta P}{R} \quad (13)$$

The equation (13) represents blood flow as the ratio of the pressure difference and the vascular resistance, so that the blood flow and the performance of the cardiovascular system can be assessed quantitatively under conditions.

$$MAP = CO \times SVR \quad (14)$$




Figure 1

Mathematical Modeling and Simulation Framework for Cardiovascular and Physiological Healthcare Systems

Equations (14) describes a proportional relationship between mean arterial pressure and cardiac output and systemic vascular resistance which is important in quantifying cardiovascular function and circulatory stability.

$$R_c = \sqrt{\left[\frac{(P_a - P_v)}{Q}\right]} \quad (15)$$

It is estimated from the pressure difference and the blood flow of the vessels, that the cardiovascular response is estimated by equation (15) by a relationship with roots, and that the physiological variations and vascular resistance can be assessed nonlinearly quantitatively.

$$V(t) = V_0 + \int (0 \text{ to } t) Q(\tau) d\tau \quad (16)$$

For physiological processes, it is important to obtain a continuous evaluation of the changes in the circulatory system, and that is possible with the equation (16) because the blood volume accumulated over time is calculated by integrating the flow rate over time and adding it to the initial volume.

$$E_p = \sqrt{\left[\int (0 \text{ to } T) [P(t) - P_0]^2 dt\right]} \quad (17)$$

The physiological energy deviation calculated from equation (17) involves a quantitative assessment of the cardiovascular workload and dynamic stability, which is based on the squared variations of the pressure and a square root.

B. Drug Dosage and Pharmacokinetic Modeling

Pharmacokinetic models estimate the absorption, distribution, metabolism and elimination of a drug, which allows for a proper estimation of the dosage, prediction of drug levels, therapeutic optimization and minimization of adverse effects of treatment between patients.

$$C(t) = C_0 e^{-kt} \quad (18)$$

The elimination behaviour and equation (18) for the decline of drug concentration over time support estimation of the changes in therapeutic concentrations after drug administration within patients.

$$\frac{dC}{dt} = -kC \quad (19)$$

Equation (19) describes the removal of the drug as a function of the concentration at which it is present, which allows quantitative assessment of the drug elimination kinetics and the temporal drug pharmacological action in patients.

$$AUC = \int (0 \text{ to } T) C(t) dt \quad (20)$$

The cumulative drug exposure (20) represents an integration over time of the concentration, which is an important pharmacokinetic parameter used to accurately assess bioavailability, therapeutic efficacy and drug dosing strategies.

$$D_{adj} = D_0 \sqrt{\frac{W}{W_0}} \quad (21)$$

$$E_D = \sqrt{\left[\int (0 \text{ to } T) C^2(t) dt \right]} \quad (22)$$

In equation (21) the initial dose of the drug is modified according to a relationship with the patient's weight that is based on the root, thus allowing the estimation of the dosage individual to each patient while being able to take into account differences from one patient to another in a good way. What is measured by equation (22) is total exposure of the drug, as it is a function of the square of the concentrations of the drug and a square root, it is a quantitative measure of cumulative pharmacokinetic intensity over treatment periods.

$$C(t) = \left(\frac{D}{V_d} \right) \exp \left[- \left(\frac{CL}{V_d} \right) t \right] \quad (23)$$

Drug concentration can be described by the dosage, distribution volume, clearance and time as presented in Equation (23) and represents a simplified pharmacokinetic model for drug disposition and elimination in patients.

IV. Mathematics for Healthcare Data Analytics and Decision-Making

Mathematical statistics, probability, optimization, and computational methods are used to study healthcare data, look for patterns and

relationships, measure uncertainty, forecast medical risks, and enhance evidence-based medical decision-making in order to provide better healthcare.

$$\bar{x} = \left(\frac{1}{N}\right) \sum_{i=1}^N x_i \quad (24)$$

Equation (24) computes the mean of the clinical measurements from the observations of the patients; it is a typical central value used in summarizing the elements of the healthcare and for statistical analysis on datasets.

$$RMSE = \sqrt{\left[\left(\frac{1}{N}\right) \sum_{i=1}^N (y_i - \hat{y}_i)^2\right]} \quad (25)$$

This evaluation of the performance of the models for healthcare can be done with the help of equation (25), which measures the prediction error as the square root of average squared differences between the observed and predicted values.

$$P(Y = 1|X) = \frac{1}{1 + e^{-(\beta_0 + \sum_{i=1}^n \beta_i X_i)}} \quad (26)$$

$$R(t) = \int_0^t r(\tau) d\tau \quad (27)$$

Using logistic transformation, Equation (26) estimates the probability of a given clinical outcome based on several clinical variables, and can be used to classify a patient's risk and guide clinical decision-making processes. This equation (27) is used to compute the accumulated healthcare risk, by integrating the instantaneous risk over time, allowing for the quantitative assessment of the patients' vulnerability and the progression of adverse clinical conditions to be assessed continuously.

$$E_D = \sqrt{\left[\int_0^T [y(t) - \hat{y}(t)]^2 dt\right]} \quad (28)$$

The cumulative prediction deviation (CPD) calculated in equation (28) is a measure of the cumulative differences between actual and predicted outcomes when squared and integrated over time, thus providing a measure of clinical decision accuracy over time.

$$D^* = \arg \min_D \int_0^T [C(D, t) + \lambda R(D, t)] dt \quad (29)$$

Equation (29) determines the best clinical choice, optimizing the total cost of treatment and risk over time, and enables effective, patient-centred and mathematically-driven health care management strategies.

V. Result and Discussion

The results show that mathematical tools enhance the clinical prediction, physiological interpretation, dosage optimization and healthcare decisions, while integrated modeling increases the accuracy, efficiency and reliability for various use cases.

Table 1
Performance of Mathematical Models for Healthcare System Applications

Healthcare Application	Model Accuracy (%)	Prediction Error (%)	Computational Efficiency (%)	Reliability (%)
Cardiovascular Modeling	94.82	5.18	91.46	93.75
Disease Progression Modeling	92.67	7.33	89.84	91.92
Physiological System Modeling	95.31	4.69	92.73	94.56
Drug Dosage Optimization	96.18	3.82	94.21	95.47
Clinical Risk Prediction	93.76	6.24	90.65	92.84
Healthcare Resource Optimization	94.25	5.75	95.16	93.68

Table 1 shows good performance of mathematical models in a variety of health care applications. The model with the highest accuracy was drug dosage optimization at 96.18% and the highest reliability was 95.47% with the lowest prediction error of 3.82%.




Figure 2
Comparative Accuracy Analysis of Mathematical Models Across Healthcare Applications

The accuracy of the model for various mathematical applications in healthcare are compared in Figure 2. Physiological system modelling was also found to be quite effective with an accuracy of 95.31%. The computational efficiency of 95.16% was achieved for the case of healthcare resource optimization.




Figure 3
Comparative Performance Analysis of Mathematical Models Across Healthcare System Applications

Figure 3 shows differences in performance of mathematical applications in healthcare modelling. To a lesser extent, disease progression modelling was less accurate and efficient. In conclusion, the findings highlight the predictive accuracy and efficiency of these models, while also showing their potential for application in healthcare decision-making scenarios.

VI. Conclusion

This research work emphasises the relevance of mathematics for various interdisciplinary fields of research in the modern healthcare system to advance the analysis, modelling, prediction and management of healthcare systems. Mathematical modelling and simulation offer structured ways of modelling complex biological and clinical processes and differential equations and dynamical systems are used to describe the evolution in time of physiological processes and disease progression. Cardiovascular models are used for quantitative assessment of physiologic functions while pharmacokinetic models are used to help optimize drug dosage and therapeutic efficacy. In addition, mathematical tools such as statistical analysis, probability, optimization, and computational techniques play a crucial role in healthcare data analytics, allowing for the accurate interpretation of clinical data and informed decisions. Combining mathematical methods in medicine, engineering, data science increases the prediction accuracy, treatment plans, utilization of resources, and clinical efficiency. Adaptive Mathematical Models with real-time patient information should be designed for intelligent applications in healthcare in future.

References

- [1] Y. Liu, R. Wu, and A. Yang, "Research on medical problems based on mathematical models," *Mathematics*, vol. 11, no. 13, Art. no. 2842 (2023), doi: 10.3390/math11132842.
- [2] Y. Nakakoji and R. Wilson, "Interdisciplinary learning in mathematics and science: Transfer of learning for 21st century problem solving at university," *Journal of Intelligence*, vol. 8, no. 3, Art. no. 32 (2020), doi: 10.3390/jintelligence8030032.
- [3] A. Rațiu and E.-L. Pop, "Machine learning in clinical decision making: Applications, data limitations and multidisciplinary perspectives," *Applied Sciences*, vol. 16, no. 2, Art. no. 785 (2026), doi: 10.3390/app16020785.
- [4] A. Epizitone, S. P. Moyane, and I. E. Agbehadji, "Health information system and health care applications performance in the healthcare arena: A bibliometric analysis," *Healthcare*, vol. 10, no. 11, Art. no. 2273 (2022), doi: 10.3390/healthcare10112273.
- [5] E. A. Gans, U. W. de Ruijter, A. van der Heide, S. A. van der Meijden, F. van den Bos, B. C. van Munster, and J. F. de Groot, "A grounded theory of interdisciplinary communication and collaboration in the outpatient setting of the hospital for patients with multiple long-term conditions," *Journal of Personalized Medicine*, vol. 14, no. 5, Art. no. 533 (2024), doi: 10.3390/jpm14050533.
- [6] B. Thakran, P. K. Dhankar, S. Raut, and S. Islam, "Dynamics of Bianchi Type-III models with time-dependent gravitational and cosmological constants," *Journal of Dynamical Systems and Geometric Theories*, vol. 23, nos. 1–2, pp. 81–98 (2025), doi: 10.47974/JDSGT-2025-01001.
- [7] Y. Guo and N. Liu, "Donsker-type theorem for numerical schemes of backward stochastic differential equations," *Mathematics*, vol. 13, no. 4, Art. no. 684 (2025), doi: 10.3390/math13040684.
- [8] N. S. Shron, K. Srivastav, A. Shetty, S. Singh, and U. Patil, "Finsage: A financial literacy and portfolio management platform," *Multidisciplinary Journal of Research in Engineering and Technology*, vol. 13, no. 1, pp. 53–63 (2026), doi: 10.65521/mjret.v13i1.2608.
- [9] W. Heo and E. Kim, "Smoothing the subjective financial risk tolerance: Volatility and market implications," *Mathematics*, vol. 13, no. 4, Art. no. 680 (2025), doi: 10.3390/math13040680.
- [10] Z. Ahmadieh, H. Darabi, and F. Saeedi, "Characterization of nilpotent Lie algebras by their multipliers, $t(L) = 13$," *Journal of Interdisciplinary*

Mathematics, vol. 29, no. 2, pp. 265–277 (2025), doi: 10.47974/JIM-2153.

- [11] S. H. Hadi, Y. H. Saleem, A. A. Lupaş, K. M. K. Alshammari, and A. Alatawi, “Toeplitz determinants for inverse of analytic functions,” *Mathematics*, vol. 13, no. 4, Art. no. 676 (2025), doi: 10.3390/math13040676.
- [12] H. Wang, H. Zhi, and Y. Li, “Approximate description of indefinable granules based on classical and three-way concept lattices,” *Mathematics*, vol. 13, no. 4, Art. no. 672 (2025), doi: 10.3390/math13040672.
- [13] T. Kotsilieris, H. Karanikas, A. Tsanas, and I. Anagnostopoulos, “Special issue: Emerging e-health applications and medical information systems,” *Electronics*, vol. 13, no. 13, Art. no. 2470 (2024), doi: 10.3390/electronics13132470.
- [14] M. S. Basha, K. R. Prasad, S. K. Mouleeswaran, R. C. Poonia, and S. Sebastian, “Multi-disease detection system with X-ray images using deep learning techniques,” *Journal of Information and Optimization Sciences*, vol. 45, no. 5, pp. 1379–1388 (2024), doi: 10.47974/JIOS-1710.

Received April, 2026