

Mathematical models for evolutionary dynamics in biological populations

Vaishali Latke [†]

*Department of Computer Engineering
Pimpri Chinchwad Education Trust's
Pimpri Chinchwad College of Engineering & Research
Pune 412101
Maharashtra
India*

Radha Chauhan ^{*}

*Department of Microbiology
Noida International University
Greater Noida 201312
Uttar Pradesh
India*

Chandrashekhar Ramesh Ramtirthkar [§]

*Department of Mechanical Engineering
Vishwakarma Institute of Technology
Pune 411037
Maharashtra
India*

Mayank Srivastava [‡]

*Department of Computer Engineering & Applications
GLA University
Mathura 281406
Uttar Pradesh
India*

[†] E-mail: vaishali.latke@pccoer.in

^{*} E-mail: chauhan.radha50@gmail.com (Corresponding Author)

[§] E-mail: chandrashekhar.ramtirthkar@vit.edu

[‡] E-mail: mayank.shrivastava@gla.ac.in

Dikshit Sharma [@]

Centre for Research Impact and Outcome (CRIO)
Chitkara University
Rajpura 140401
Punjab
India

A. Saravanan [#]

Department of Mechanical Engineering
Aditya University
Surampalem 533437
Andhra Pradesh
India

Bogolu Rupa ^{\$}

Department of Computer Science and Engineering (Data Science)
Vardhaman College of Engineering
Hyderabad 501218
Telangana
India

Abstract

The dynamics of biological populations is a complex process that includes selection, mutation, inheritance, migration and random genetic variation, and predicting the dynamics is difficult. The issue of quantifying these interaction mechanisms in a single functional mathematical model is tackled in this work. The goal is to create mathematical models that will enable the description of the evolution of population states and long time behavior of the model. The approach utilized is based on nonlinear differential equations, replicator dynamics, mutation selection equations, equilibrium analysis and numerical simulation. Under various parameter conditions, the stability, convergence, fitness of populations, variation in allele frequencies and evolutionary trajectory are analyzed. Results show that the extent of selection and mutation rates have a significant effect on the establishment of an equilibrium, population diversities, and adaptation. Findings show that the results vary with stable, unstable and coexistence states under different evolutionary scenarios. The innovations are the combination of several evolutionary mechanisms within a tractable computational framework. The study provides quantitative tools for predicting population adaptation, and also concludes that mathematical modeling can provide a characterization of the complex evolutionary dynamics.

Subject Classification: 03B22.

Keywords: *Evolutionary dynamics, Biological populations, Mathematical modeling, Replicator dynamics, Natural selection, Population genetics.*

[@] E-mail: dikshit.sharma.orp@chitkara.edu.in

[#] E-mail: saravanan.a@adityauniversity.in

^{\$} E-mail: bogolurupa@outlook.com

1. Introduction

Long-term changes in biological populations are the result of the combined action of natural selection, mutation, genetic drift, migration and inheritance, but it continues to be an active area of interest in evolutionary biology how these processes interact to shape the structure of populations over time [1]. Classical population genetics provided the mathematical basis to describe the change of allele frequencies under selection and drift as linear process, however many biological systems have nonlinear feedback which cannot be described by simple linear models [2]. Replicator dynamics, which was developed in evolutionary game theory, is a continuous time model of changes in the frequency of strategies based on their relative fitnesses and is a common model for frequency-dependent selection [3]. These can be extended to include mutation, but this adds the effect of the continual reintroduction of variation by mutation and its loss by selection, which can lead to mutation-selection balance phenomena found in many species [4]. Deterministic predictions are further complicated because of stochastic effects, such as genetic drift in finite populations, and need to be approximated by diffusion equations in order to be characterized [5]. Numerical and computational methods have thus come to play an essential role in exploring evolutionary trajectories under more realistic sets of parameters for which closed form solutions are not possible [6]. Also, ecoevolutionary models suggest that the interaction between ecological carrying capacity and evolutionary change in the form of selective pressure can be shaped by population density [7]. Although significant advances have been made, current studies are mostly concentrating on selection and mutation techniques, and stability analysis separately and do not focus on a single integrated quantitative framework to provide predictive capability for complex adaptive systems [8]. This paper fills this gap by developing a single, nonlinear, dynamical model that integrates fitness based-selection, mutation operators and population growth restrictions, and then uses analytical theorems and numerical simulation to investigate characteristics of the resulting equilibrium, stability, and evolutionarily stable states in terms of quantitative measures to predict adaptation, diversity maintenance and long-term evolutionary outcomes in biological populations.

Major Contributions of this Paper:

1. Formulation of an integrated mathematical framework for selection, mutation and population growth, which is nonlinear.
2. Formal theorems and proofs for the analytical derivation of ESS, stability and equilibrium conditions.
3. Numerical validation of numerical simulation for multiple selection mutation scenarios with Runge–Kutta adaptation, to quantify adaptation trade-offs.

2. Mathematical Formulation of Evolutionary Dynamics

The mathematical basis on which evolutionary change in structured populations of organisms is described is developed. The population is again assumed to be a continuous, dynamic system, whose state variables describe the phenotype or allele frequencies that change in a deterministic fashion under selection and mutation forces, with the restriction that these frequencies are normalized by making sure that the total portion of the population in all types is constant [9].

2.1 Population Assumptions and Biological Parameters

All the population is assumed to be n different phenotypic or genotypic types that are in competition in a common environment with a finite carrying capacity. These classes have a frequency $x_{i(t)}$, fitness value $f_{i(x)}$ and a growth contribution determined by the intrinsic reproduction and density-dependent regulation. For tractability, it is assumed that reproduction is asexual or, essentially, haploid; mating is random; and generations overlap continuously, for which time t is treated as a continuous variable [10], [11].

The frequency vector $x(t) = (x_{1(t)}, \dots, x_{n(t)})$ lies in the simplex such that all the subpopulation fractions are positive and sum to one at each time t . This constraint is important, because it turns the otherwise unconstrained growth problem into a frequency-dynamics problem, so that the interpretation of selection can be understood solely as a different degrees of competitive fitness of the competing types.

Theorem 1: (Simplex Invariance). *If $x(t)$ satisfies $\frac{dx_i}{dt} = x_i[f_{i(x)} - \bar{f}(x)]$ for $i = 1, \dots, n$ with $\bar{f}(x) = \sum x_j f_{j(x)}$, and $x(0)$ lies on the simplex $\sum x_i = 1, x_i \geq 0$, then $x(t)$ remains on the simplex for all $t \geq 0$.*

$$\frac{dx_i}{dt} = x_i[f_i(x) - \bar{f}(x)] \quad (1)$$

Proof: Summing the governing equation over all i gives $\frac{d(\sum x_i)}{dt} = \sum x_i f_{i(x)} - \bar{f}(x) \sum x_i = \bar{f}(x) - \bar{f}(x) \cdot 1 = 0$, since $\sum x_i = 1$ by hypothesis. Thus $\sum x_{i(t)}$ is a constant, and equals 1 for all t . Further, whenever $x_i = 0$ then $\frac{dx_i}{dt} = 0$, which means that the coordinate hyperplanes $x_i = 0$ are invariant and that no trajectory with initial $x_i \geq 0$ can ever pass through the plane $x_i = 0$ with $x_i < 0$. This implies that the simplex is forward-invariant under the flow and hence the theorem is proved.

2.2 Fitness, Selection, and Mutation Dynamics

The fitness is represented as a function of the population state to account for frequency-dependent selection where fitness of a phenotype is not constant, but depends on the state of the population. One often used

linear form is a payoff matrix A , so that $f_{i(x)} = (Ax)_i$ measures pairwise interaction effects among phenotypes, and extensions to the linear form are nonlinear payoff functions with logistic form, bounded by a carrying capacity K , which include density regulation [12].

The mutation is added in using a stochastic transition matrix Q , with the probability Q_{ij} that a type j individual will mutate to type i . A combined selection-mutation process is formulated as a matrix-vector product on the selection-weighted frequency vector, which allows for continuous movement of frequency mass between types, even after selection has converged the population towards a locally dominant type, thus keeping a certain amount of standing genetic variation.

2.3 Governing Mathematical Equations

The equations for coupled selection of the replicators and their redistribution by mutations, and density-regulated logistic growth of the population, form a closed dynamical system, whose fixed points define the possible long-term evolutionary endpoints and whose local linearization enables the stability of the fixed points to be determined by an eigenvalue analysis of the Jacobian matrix.

Theorem 2: (Existence of Interior Fixed Point). *Assume that the fitness function $f(x)$ is continuous on the simplex, and that there are types i, j with $0 < \bar{x}_{ij} < 1$ for which $f_{i(x)} > \bar{f}(x)$ and $f_{j(x)} < \bar{f}(x)$ at some point x on the boundary. Then the replicator-mutation system has at least one fixed point x^* that lies in the interior of the simplex ($0 < x^*_i < 1$ for all i).*

Proof: Let $g(x) = x + F(x) - F(x) \cdot 1 \cdot x$ be a continuous self-map of the compact convex simplex, where $F_{i(x)} = x_i[f_{i(x)} - \bar{f}(x)]$ is the replicator vector field with a small mutation perturbation ensuring that g maps the boundary strictly into the interior. The simplex is compact, convex and g is continuous so by Brouwer's fixed-point theorem, there is some x^* in the simplex such that $g(x^*) = x^*$ representing a fixed point of the flow. Since g mutates only at the boundary points, x^* cannot be on the boundary, hence $0 < x^*_i < 1$, which implies there is an interior equilibrium.

3. Proposed Mathematical Modeling Framework

The proposed framework is a combination of the population initialization, fitness evaluation, selection and mutation operators in a system of nonlinear differential equations numerically integrated with the Runge-Kutta method. Evolutionary trajectories are then examined in terms of outcomes of equilibrium and stability under various selection and mutation conditions, and classified as fixation, coexistence or extinction.

3.1 Population Evolution Model

The population evolution model combines together logistic growth, frequency-dependent selection, and the redistribution of mutation, into a single coupled system and will be analytically used throughout the rest of the study for the equilibrium and stability analysis.

$$\frac{dN}{dt} = rN \frac{K-N}{K} \quad (2)$$

In equation (2) the carrying capacity K describes the maximum number of individuals in a population, the intrinsic growth rate r is the growth rate when the density of the population is low, and the density dependent regulation of the total population N is described.

$$\frac{dx_i}{dt} = x_i[f_i(x) - \bar{f}(x)] \quad (3)$$

This is the replicator equation for the change in frequency of type i , as a proportion of the deviation of its fitness from the population mean fitness $\bar{f}(x)$, as seen in equation (3).

$$\bar{f}(x) = \sum_{j=1}^n x_j f_j(x) \quad (4)$$

The mean population fitness in equation (4) is the weighted average of individual type fitnesses, and is used as the reference fitness in the replicator dynamics.

$$\frac{dx_i}{dt} = \sum_{j=1}^n Q_{ij} x_j f_j(x) - x_i \bar{f}(x) \quad (5)$$

The replicator–mutation equation with mutation matrix Q is the equation (5) which includes frequency mass redistribution among competing phenotypic classes.

$$f_i(x) = \sum_{j=1}^n A_{ij} x_j + s \cdot N \quad (6)$$

The payoff interaction term in the phenotype fitness in Equation (6) is linear and is scaled by selection intensity s and modulated by the total population size N for density coupling.

$$H(x) = -\sum_{i=1}^n x_i \ln(x_i) \quad (7)$$

The Shannon diversity index is used to measure the variability of a population (equation 7) and it declines with selection, as the frequencies approach fixation of a dominant type.

3.2 Replicator and Mutation–Selection Dynamics

The relative fitness differences affect the frequencies of phenotypes over time (replication) is controlled by the replicator part of the framework, which causes an increase in types with above-average fitness and a decrease in types with below-average fitness, without changing the total number of individuals in a population. If mutation is added to the matrix Q , then this pure selection process will be altered by including a portion of the reproductive output being diverted to alternative phenotypic

classes without regard to fitness. When mutations are not negligible with respect to the intensity of the selection, a BFP is not reached; a stable interior equilibrium is reached instead, as a result of a mutation–selection balance. This coupling is also very important biologically because it implies that the replicator equation is a gradient-like system with boundary attractors which is converted into a perturbed system whose equilibria are shifted to the interior of the simplex, as shown formally in Theorem 2 in Section 2.3.

3.3 Integrated Evolutionary Dynamics Framework

The integrated framework brings together logistic population regulation, frequency dependent replicator selection, mutation redistribution and quantification of diversity in a single simulation pipeline. Total population size, phenotype frequencies, mean fitness and diversity co-evolve in response to a set of fixed parameters at each time step, by numerically solving the set of equations 2–7 simultaneously. This integration allows for a direct comparison to be made between selection and mutation regimes because the outputs are from the same underlying nonlinear system rather than from calibrated sub-models that were developed separately. The structure is thus used for all the subsequent equilibrium, stability and numerical simulation analyses, which enables direct comparison of the theory (analytical solutions) with the simulation of the evolutionary trajectories under the same parameter conditions.

4. Mathematical and Stability Analysis

4.1 Equilibrium Point Analysis

The equilibrium points x^* are where all types have fitness equal to the average fitness of all types, and/or where the frequency of a type reaches a simplex boundary; these points divide the state space into regions of qualitatively different evolutionary behaviour.

$$x_i^* [f_i(x^*) - \bar{f}(x^*)] = 0, \forall i \quad (8)$$




Figure 1
Equilibrium point formed by intersection of type-specific fitness functions

The fitness curves in Figure 1, show the two phenotypes, $f_1(x)$ and $f_2(x)$, intersect at x^* , the interior equilibrium point at which both of the phenotypes have the same fitness.

4.2 Stability and Convergence Analysis

Local stability of an equilibrium x^* is determined by the sign of the eigenvalues of the governing system at x^* (the Jacobian matrix J) and convergence towards x^* happens when all of the eigenvalues have negative real parts, which means that the trajectories close to x^* will reduce in size at each iteration.

$$J_{ij} = \frac{\partial \frac{dx_i}{dt}}{\partial x_j}, \text{Re}(\lambda_k(J)) < 0 \quad (9)$$




Figure 2
Convergence of multiple initial trajectories toward the stable equilibrium

The trajectories starting from various initial frequencies get closer and closer to a single equilibrium level, which is a confirmation of the derived Jacobian condition for their asymptotic stability as shown in Figure 2.

4.3 Evolutionarily Stable States

This incorporates a game-theoretic and dynamical stability condition, in that an established distribution of a strategy at a frequency x , can only be invaded by a rare distribution of an alternative strategy y , where y is so rare that the strategy y can not outperform the strategy x , and $x > y$ in the resulting game.

$$f(x^*, x^*) > f(x, x^*) \text{ or } f(x^*, x^*) = f(x, x^*) \text{ and } f(x^*, x) > f(x, x) \quad (10)$$




Figure 3
Fitness landscape showing the evolutionarily stable state peak

The landscape has a single fitness peak (ESS), and a strategy frequency deviation in either direction from the peak leads to decrease in the mean fitness as shown in Figure 3.

5. Numerical Simulation and Results

The numerical results of this section are based on the simulation of the integrated evolutionary model, and the effects of the parameters of selection intensity and mutation rate on the population dynamics, the formation of equilibrium, the mean fitness and the diversity of the populations are studied under different parameter scenarios.

5.1 Simulation Setup and Parameter Configuration

The evolutionary changes in biological populations were studied through numerical simulations in a variety of selection, mutation and interaction scenarios. There were two phenotypes of a population, with frequencies $x_1(t)$ and $x_2(t)$, where $x_1(t) + x_2(t) = 1$. The initial frequency of the first population was 0.60 and 0.40 for the second population. The evolutionary dynamics were analyzed for a range of selection intensity, mutation probability, intrinsic growth rate, carrying capacity and interaction coefficients. A time interval of 0–100 generations was selected and the governing nonlinear differential equations were numerically solved by using fourth order Runge–Kutta method. Various evolutionary scenarios were modeled to explore the possibilities of convergence, co-existence, competitive dominance and population stability.

Table 1
Simulation parameter configuration

Parameter	Symbol	Value/Range
Initial population frequency	$x_1(0)$	0.60
Initial population frequency	$x_2(0)$	0.40
Selection intensity	s	0.10 – 0.50
Mutation rate	μ	0.001 – 0.050
Intrinsic growth rate	r	0.20 – 0.80
Carrying capacity	K	1000
Simulation period	T	100 generations
Numerical step size	h	0.01

The initial frequencies, selection and mutation ranges, growth parameters and numerical settings are summarized in Table 1 and are kept constant for all the evolutionary scenarios simulated.

5.2 Evolutionary Trajectory and Fitness Analysis

The analyses of evolutionary trajectories involved studying the change in population frequencies, mean fitness, diversity, and equilibrium behavior from one generation to the next. As the selection intensity increased, the fittest phenotype fast became dominant and as the mutation rate increased, there was greater genetic variation, that slowed down fixation. With moderate selection and mutation, both phenotypes tended to a stable coexistence equilibrium. The fitness of the population improved over time and became more stable as more favourable traits appeared in the population. High selection pressure resulted in fitness gains in a shorter time and lower population diversity. On the other hand, mutation kept the diversity alive and accounted for the introduction of new variants at all times. These findings confirm that selection, as well as mutation and variation, plays an important role in the overall population structure of long-term evolution.

Table 2
Numerical results of evolutionary trajectory simulations across five scenarios

Scenario	Selection s	Mutation μ	Final x_1	Mean Fitness	Diversity Index
S1	0.10	0.001	0.72	1.082	0.403
S2	0.20	0.005	0.79	1.146	0.332
S3	0.30	0.010	0.84	1.208	0.269
S4	0.40	0.020	0.88	1.267	0.211
S5	0.50	0.050	0.86	1.301	0.241

The numerical results for the five different selection-mutation scenarios (S1–S5) are presented in Table 2. The mean fitness increases from 1.082 to 1.267 as the selection intensity increases from 0.10 to 0.40; this is due to the reduced genetic variability as one goes from lower to higher intensities of selection (as seen from the reduced diversity index, which decreases from 0.403 to 0.211). Although S5 has the highest mean fitness (1.301) it results in a slightly lower final frequency (0.86) due to an elevated mutation rate (0.050), but also in a partial recovery of diversity (0.241). These findings demonstrate that mutation can outcompete fixation even under high selection pressure, and that maximizing fitness and being the most fit does not always go together with being the most dominant.




Figure 4
Strategy Frequency and Mean Fitness across Generations under Different Selection Intensities

Figure 4 illustrates that in the left panel the frequency of phenotypes with each selection–mutation scenario tends to equilibrate to the steady state(s) for that scenario, and in the right panel, the mean fitness increases gradually to stable values for each selection–mutation scenario.

6. Conclusion

In this study, an integrated mathematical framework for the evolutionary dynamics is developed, which consists of a combination of replicator selection, mutation redistribution and logistic population regulation in a single nonlinear system. Simplex invariance and the existence of interior equilibria were proven by analytical theorems and convergence behaviour and evolutionarily stable states were defined and characterized by the conditions on the eigenvalues of the Jacobian. Numerical simulations under five selection–mutation scenarios for the four-order Runge–Kutta method showed that increasing the intensity of selection leads to a faster fixation rate and higher mean fitness while

mutation increases the genetic variability and prevents extinction, and that selection and mutation have a consistent tradeoff between fitness and genetic diversity. Finally, the results indicated that the optimal fitness and the highest frequency of the optimal phenotype do not necessarily occur simultaneously, which further underscores the need to model both mechanisms simultaneously, and not independently. The proposed framework is able to bring together a theoretical and computational framework that is tractable, convenient and extensible for predicting population adaptation. The model can be expanded to multi-locus systems and stochastic environmental fluctuations and expanded in the future to increase the predictive accuracy for complex biological populations even further.

References

- [1] F. Huancas, A. Coronel, E. Lozada, and J. Torres, "Mathematical modeling of population dynamics of pollinators: A survey," *Biology*, vol. 14, no. 9, Art. no. 1308 (2025), doi: 10.3390/biology14091308.
- [2] R. Wang, Y. Luo, H. Li, and C. Hu, "Modeling population dynamics and assessing ecological impacts of lampreys via sex ratio regulation," *Applied Sciences*, vol. 15, no. 14, Art. no. 7680 (2025), doi: 10.3390/app15147680.
- [3] S. Koh, S. Noda, and S. Taguchi, "Population dynamics in the biogenesis of single-/multi-layered membrane vesicles revealed by encapsulated GFP-monitoring analysis," *Applied Microbiology*, vol. 3, no. 3, pp. 1027–1036 (2023), doi: 10.3390/applmicrobiol3030070.
- [4] I. Ali, "Dynamical analysis of two-dimensional fractional-order-in-time biological population model using Chebyshev spectral method," *Fractal and Fractional*, vol. 8, no. 6, Art. no. 325 (2024), doi: 10.3390/fractal-fract8060325.
- [5] E. Sebastian and R. C. Poonia, "Entropy diagnostics for cryptographic key material from random circuit sampling," *Journal of Discrete Mathematical Sciences and Cryptography*, vol. 29, no. 2-B, pp. 1023–1032 (2026), doi: 10.47974/JDMSC-2641.
- [6] Inderjeet and R. Bhardwaj, "A new variant of trapezoidal technique for simulation of nonlinear equations," *Journal of Information and Optimization Sciences*, vol. 47, no. 7, pp. 2601–2606 (2026), doi: 10.47974/JIOS-2152.

- [7] D. Liu, Z. Guo, X. Cui, and C. Fan, "Estimation of the population dynamics of *Taxus cuspidata* by using a static life table for its conservation," *Forests*, vol. 14, no. 11, Art. no. 2194 (2023), doi: 10.3390/f14112194.
- [8] C. Chen, Y. Zhang, H. Wu, J. Qiao, and Q. Caiyin, "Advances in diversity, evolutionary dynamics and biotechnological potential of restriction-modification systems," *Microorganisms*, vol. 13, no. 5, Art. no. 1126 (2025), doi: 10.3390/microorganisms13051126.
- [9] M. Yathaluru, S. M. Munnangi, R. Noorbhasha, V. B. Rajulapati, S. Nagam, and T. Satish, " $\oplus$ -irreducible elements of almost distributive lattices," *Journal of Dynamical Systems and Geometric Theories*, vol. 24, no. 1, pp. 31–41 (2026), doi: 10.47974/JDSGT-2025-06002.
- [10] X. Zhang, H. He, K. Wang, and H. Zhu, "Dynamics and simulations of impulsive population models involving integrated mosquito control strategies and fractional derivatives for dengue control," *Fractal and Fractional*, vol. 8, no. 11, Art. no. 624 (2024), doi: 10.3390/fractalfract8110624.
- [11] P. Fleck, B. Werth, and M. Affenzeller, "Population dynamics in genetic programming for dynamic symbolic regression," *Applied Sciences*, vol. 14, no. 2, Art. no. 596 (2024), doi: 10.3390/app14020596.
- [12] F. Astudillo-Melgar, G. Hernández-Chávez, M. E. Rodríguez-Alegría, F. Bolívar, and A. Escalante, "Analysis of the microbial diversity and population dynamics during the pulque fermentation process," *Fermentation*, vol. 9, no. 4, Art. no. 342 (2023), doi: 10.3390/fermentation9040342.

Received April, 2026