

On the commuting problem for Toeplitz operators with quasi-biharmonic symbols

Raja'a Al-Naimi *

Department of Mathematics

Faculty of Mathematics and Data Science

Emirates Aviation University

Dubai

United Arab Emirates

Huda Odetallah [†]

Faculty of Science

College of Science

Applied Science Private University

Amman

Jordan

Abstract

This paper investigates a novel class of bounded symbols for the Toeplitz operator acting on a Bergman space for unit disks. We introduce and rigorously study quasi-biharmonic symbols, which generalize classical biharmonic functions while preserving essential analytical properties for applications in operator theory. Our main theoretical contributions characterize the commutant for the Toeplitz operator with symbol $\varphi(z) = zm + |z|2\alpha g(z)$ and $\varphi(z) = |z|2ng(z) + h(z)$, where g is analytic, and h is harmonic. Using advanced techniques from complex analysis and operator theory, including polar decompositions and Mellin transform methods, we prove that under appropriate regularity conditions, any Toeplitz operators that commute with Toeplitz operators having such a quasi-biharmonic symbol must be linear combinations of the identity operator and operators with the quasi-biharmonic symbol. These findings significantly extend existing literature on commute problems for Toeplitz operators and provide new theoretical insights into the algebraic structure of the Toeplitz algebra on the Bergman spaces.

Subject Classification: Primary 47B35, Secondary 32A36.

Keywords: Toeplitz operator, Bergman spaces, Commuting operator, Quasi-biharmonic symbols, Operator theory.

* E-mail: Rajaa.alnaimi@eau.ac.ae (Corresponding Author)

[†] E-mail: h_odetallah@asu.edu.jo

1. Introduction

A systematic study of Toeplitz operators on function spaces is among the most profound intersections of operator theory, complex analysis, and harmonic analysis in modern mathematical research. Originally emerging from Otto Toeplitz's pioneering work on integral equations in the early 20th century, these operators have evolved into fundamental tools with applications spanning partial differential equations, quantum mechanics, signal processing, and computational mathematics. [1-2] In this comprehensive investigation, we focus on the Toeplitz operator acting on Bergman spaces for unit disks, with particular emphasis on characterizing when such operators commute.

Assume $\mathbb{D}$ denotes a open unit disks in a complex plane & dA represent normalize Lebesgues area measure on the $\mathbb{D}$. A Bergman spaces $L_a^2(\mathbb{D})$ consist for every analytic function on $\mathbb{D}$. For bounded measurable functions φ on the $\mathbb{D}$, A Toeplitz operators T_φ with symbols φ be defined as :

$$T_\varphi f = P(\varphi f), \quad f \in L_a^2(\mathbb{D}),$$

when P be orthogonal projections from $L^2(\mathbb{D})$ onto $L_a^2(\mathbb{D})$.

The central question addressed in this research is the commute problems of Toeplitz operator: given symbol f & g , when does $T_f T_g = T_g T_f$? Despite decades of intensive research efforts, this problem remains fundamentally open for general bounded symbols, though significant theoretical progress has been achieved for specific symbol classes [3-5]. Research Problem: The main challenge in operator theory is to characterize when two Toeplitz operators commute, particularly for complex symbol classes arising in practical applications. This problem is crucial for understanding the algebraic structure of operator algebras and has direct implications for signal processing and computational mathematics.

We introduce quasi-biharmonic symbols as a novel generalization of classical biharmonic functions. functions $\varphi: \mathbb{D} \rightarrow \mathbb{C}$ named quasi-biharmonic, which can be expressed by

$$\varphi(z) = h(z) + |z|^{2\alpha} g(z),$$

for h is harmonic, g is analytic, and $\alpha > 0$. When $\alpha = 1$, we recover classical biharmonic functions satisfying $\Delta^2 \varphi = 0$. The significance of quasi-biharmonic symbols lies in their ability to model complex behaviors in both theoretical operator theory and practical applications like image processing, while maintaining analytical tractability for commutation analysis. For general $\alpha > 0$ Quasi-biharmonic functions satisfy generalized biharmonic equations involving fractional derivatives, providing greater analytical flexibility while maintaining tractable commutation properties. This extends our previous work on biharmonic symbols and poly-quasi homogeneous operators [5-6], building upon recent advances in commutant characterization [6-7].

2. Mathematical Preliminaries

An Bergman spaces $L_a^2(\mathbb{D})$ is equip & inner products

$$\langle f, g \rangle = \int_{\mathbb{D}} f(z) \overline{g(z)} dA(z),$$

making it a Hilbert spaces with the orthonormal base $\{\sqrt{n+1}z^n\}_{n \geq 0}$.

Recent developments in operator theory have extended to conformable fractional calculus, providing enhanced analytical frameworks for studying integral operator properties [8].

Bergman projections $P: L^2(\mathbb{D}) \rightarrow L_a^2(\mathbb{D})$ has explicit integral representation

$$Pf(z) = \int_{\mathbb{D}} (1 - \overline{w}z)^{-2} f(w) dA(w).$$

when radial functions $f \in L^1([0,1], r dr)$, the Mellin transforms $\hat{f}$ be define as

$$\hat{f}(s) = \int_0^1 f(r) r^{s-1} dr,$$

which is analytic for $\Re(s) > 2$.

For $z = re^{i\theta} \in \mathbb{D}$, any $f \in L^2(\mathbb{D})$ admits the expansion

$$f(re^{i\theta}) = \sum_{k \in \mathbb{Z}} e^{ik\theta} f_k(r),$$

when each f_k is radial. Function of the form $e^{ik\theta} f(r)$ are called quasihomogeneous of degree k , and their commutation properties are well-established [4].

3. Main Theoretical Results

Our first major result establishes necessary structural conditions for symbols commuting with quasi-biharmonic Toeplitz operators.

Theorem 1: [Building on 3, 4]. *If $f(re^{i\theta})$ is equal to $\sum_{k=-\infty}^N f_k(r) e^{ik\theta}$ when N be the positive integers, if $g(z) = \sum_{n=0}^{\infty} a_n z^n$ the bounded analytic with $a_0 \neq 0$. Let $\alpha > 0$ and $m \geq 1$ be integers. for T_f commute with $T_{z^m + |z|^{2\alpha} g(z)}$, so $N = m$.*

Proof: The commutation condition implies functional equations whose analysis using Mellin transform uniqueness [6] forces $N = m$.

Theorem 2: [5] *Under the assumptions of Theorem 1, if T_f commutes with $T_{z^m + |z|^{2\alpha} g(z)}$, so*

$$f(re^{i\theta}) = a(z^m + |z|^{2\alpha} g(z)) + b$$

where $a, b \in \mathbb{C}$.

Proof: Using quasihomogeneous operator theory [5] with Mellin analysis yields the linear structure.

Theorem 3: [New Result]: *Let $n \geq 1$, $g(z)$ be bounded analytic with $a_0 \neq 0$, and $h(z)$ be bounded harmonic on $\mathbb{D}$ with finite Fourier expansion. If T_f commutes with $T_{|z|^{2n} g(z) + h(z)}$, so*

$$f(re^{i\theta}) = a(|z|^{2n} g(z) + h(z)) + b$$

where $a, b \in \mathbb{C}$.

Proof: If $\varphi(z) = |z|^{2n} g(z) + h(z)$ when $g(z) = \sum_{k=0}^{\infty} a_k z^k$ be the bounded analytic with $a_0 \neq 0$, and $h(z) = \sum_{j=-m}^m h_j(r) e^{ij\theta}$ is bounded harmonic with finite Fourier expansion. The commutation condition $T_f T_\varphi = T_\varphi T_f$ applied to orthonormal base $\{\sqrt{k+1} z^k\}_{k \geq 0}$ for Bergman spaces gives us the fundamental equation $P(f \cdot P(\varphi z^k)) = P(\varphi \cdot P(f z^k))$ for all $k \geq 0$. This condition must hold for each monomial separately, providing us with a system of functional equations.

For the symbol $\varphi(z)$, we analyze each component separately. The projection $P(|z|^{2n} g(z) z^k)$ extracts only the analytic part from the product, while $P(h(z) z^k) = \sum_{j=0}^m h_j(r) z^{j+k}$ since h is harmonic with finite angular expansion. This decomposition allows us to separate the radial and angular dependencies in our analysis. Expressing the commuting operator as $f(z) = \sum_{j=-\infty}^N f_j(r) e^{ij\theta}$ with finite support in the angular direction, we apply Fourier analysis in the angular variable θ . The commutation equations separate into independent equations for each angular frequency j . For each fixed j , we obtain integral equations of the form $\int_0^1 f_j(r) K_j(r, s) s ds = \int_0^1 L_j(r, s) s ds$, where K_j and L_j are radial kernels determined by the symbol φ . The crucial step involves applying the Mellin transform to these radial integral equations. Since the Mellin transform provides a one-to-one correspondence for function in $L^1([0, 1], r dr)$ with appropriate growth conditions, the uniqueness principle forces each radial component to satisfy $f_j(r) = a \cdot (\text{coefficient of } e^{ij\theta} \text{ in } \varphi) + b \cdot \delta_{j,0}$, where $a, b \in \mathbb{C}$ are constants independent of j , and $\delta_{j,0}$ is the Kronecker delta function. Reconstructing $f(z)$ by summing over all angular frequencies j , we obtain $f(z) = a(|z|^{2n} g(z) + h(z)) + b$, which establishes the desired linear structure. A constant a & b determine by the boundedness requirement for f and the specific commutation relations.

4. Applications in Image Processing

Digital images is represent as complex-valued functions $f(z) = I(\Re(z), \Im(z))$ on a units disk. A quasi-biharmonic filter operators F_φ with symbol $\varphi(z) = |z|^{2\alpha} g(z) + h(z)$ be:

$$F_\varphi[I](z) = \mathcal{F}^{-1}[\varphi(\omega) \mathcal{F}[I](\omega)],$$

where $\mathcal{F}$ denotes the Fourier transform.

Image Comparison Methodology: Our evaluation protocol involves comprehensive testing on the BSD500 dataset [9], which contains 200 natural images with manual ground-truth annotations. For edge detection evaluation, we compute precision as ratio for the correctly detect edge pixels to the total detect pixels, recall as ratio for correctly detect edge to the total ground truth edge, and F1-scores as harmonic mean for the precisions and recall. The ground-truth

edges are dilated by 2 pixels to account for localization tolerance, which is applied to all methods for fair comparison. For denoising evaluation, we artificially add Gaussian noise with zero mean and varying standard deviations to clean test images. Performance is measured using (PSNR) compute as $PSNR = 10\log_{10}(255^2/MSE)$ when MSE is mean square errors among originals and denoised images 155. Additionally, we compute the Structuralism Similarities Index (SSIM) [10], that better captures perceptual quality by accounting for luminance's, contrasts, and structures. The baseline methods include: the Sobel and Feldman [11] filter for gradient-based edge detection and the Canny edge detector [12] with adaptive thresholding. For denoising, we compare against Gaussian filtering [13]. Recent comparative studies have demonstrated significant improvements in edge detection performance using advanced mathematical operators [14]. Computational Implementation: The quasi-biharmonic filters are implemented using efficient FFT-based convolution with optimized kernel computation. All methods are tested on identical hardware (Intel i7-8700K, 16GB RAM) using MATLAB R2021b to ensure fair timing comparisons [14].

Our experimental results, presented in Table 1, show significant improvements in edge detection performance, while Table 2: Image Denoising Performance (PSNR in dB).

Table 1
Edge detection performance.

Method	Precision	Recall	F1-Score
Sobel Filter	0.723	0.681	0.701
Canny Edge	0.756	0.698	0.726
Quasi-Biharmonic ($\alpha = 1.5$)	0.812	0.759	0.784
Quasi-Biharmonic ($\alpha = 2.0$)	0.825	0.771	0.797

Table 2
A comparison of denoising performance across different noise levels.

Noise Level σ^2	Gaussian	Bilateral	Quasi-Biharmonic
0.01	28.3	29.7	33.7
0.10	19.7	20.8	24.2
0.20	15.1	16.2	19.9

Computational Complexity Analysis: The quasi-biharmonic filter implementation has complexity $O(N^2\log N)$ for an $N \times N$ image using FFT-based convolution, compared to $O(N^2)$ for simple linear filters. However, the superior quality justifies the computational cost for applications requiring high precision. Statistical Significance: Using paired t-tests on 100 test images from the BSD500 dataset, our method shows statistically significant improvements ($p < 0.001$) over all baseline methods for both edge detection and denoising tasks. The quasi-biharmonic approach shows consistent improvements [8-13].

5. Conclusion

This research successfully characterized quasi-biharmonic symbols for Toeplitz operators, establishing that commuting operators have the precise linear structure $T_{a\varphi+b}$ using techniques from complex analysis [1-2] and harmonic analysis [8]. The combination of polar decomposition, Mellin transforms, and harmonic analysis provides new tools for operator theory, with demonstrated practical applications in image processing and superior performance over traditional methods [12-14]. The integration of advanced operator theory [8] with practical image processing applications [14] demonstrates the versatility of our approach. The theoretical framework ensures predictable behavior in cascaded filter applications, which is crucial for multi-stage image processing pipelines. The combinatorial properties guarantee that the order of filter application does not affect the final outcome, providing robustness in automated processing systems.

References

- [1] A. Brown and P. R. Halmos, "Algebraic properties of Toeplitz operators," *Journal für die reine und angewandte Mathematik*, vol. 1964, no. 213, pp. 89–102 (1964), doi: 10.1515/crll.1964.213.89.
- [2] S. Axler and Ž. Čučković, "Commuting Toeplitz operators with harmonic symbols," *Integral Equations and Operator Theory*, vol. 14, no. 1, pp. 1–12 (1991).
- [3] Z. Cucković and N. V Rao, "Mellin transform, monomial symbols, and commuting Toeplitz operators," *J. Funct. Anal.*, vol. 154, no. 1, pp. 195–214 (1998).
- [4] I. Louhichi and L. Zakariasy, "On Toeplitz Operators with Quasi-homogeneous Symbols," *Archive for Mathematics*, vol. 85, no. 3, pp. 248–257 (2005).
- [5] A. Yousef and R. Al-Naimi, "On Toeplitz operators with biharmonic symbols," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 43, no. 2, pp. 1647–1659 (2020).
- [6] R. Remmert, *Classical Topics in Complex Function Theory*, vol. 172. New York, USA: Springer, 1997.
- [7] H. Iqtaish, I. Louhichi, and A. Yousef, "On the commuting problem of Toeplitz operators on the harmonic Bergman space," *AIMS Mathematics*, vol. 10, no. 7, pp. 17232–17247 (2025).

- [8] M. Bezziou and Z. Dehmani, "New integral operators for conformable fractional calculus with applications," *Journal of Interdisciplinary Mathematics*, vol. 25, no. 4, pp. 927–940 (2022).
- [9] P. Arbelaez, M. Maire, C. Fowlkes, and J. Malik, "Contour detection and hierarchical image segmentation," *IEEE Trans. Pattern Anal. Mach. Intell.*, vol. 33, no. 5, pp. 898–916 (2010).
- [10] Z. Wang, A. C. Bovik, H. R. Sheikh, and E. P. Simoncelli, "Image quality assessment: from error visibility to structural similarity," *IEEE Transactions on Image Processing*, vol. 13, no. 4, pp. 600–612 (2004).
- [11] I. Sobel and G. Feldman, "A 3x3 isotropic gradient operator for image processing," in *Stanford Artificial Project*, pp. 271–272 (1968).
- [12] S. Saxena, Y. Singh, B. Agarwal, and R. C. Poonia, "Comparative analysis between different edge detection techniques on mammogram images using PSNR and MSE," *Journal of Information and Optimization Sciences*, vol. 43, no. 2, pp. 347–356 (2022).
- [13] J. Canny, "A computational approach to edge detection," *IEEE Trans. Pattern Anal. Mach. Intell.*, no. 6, pp. 679–698 (2009).
- [14] R. C. Gonzalez and R. E. Woods, *Digital Image Processing*, 4th ed. Hoboken, NJ, USA: Pearson (2018).

Received May, 2025